

REMARKS

The claims have been amended to further clarify the meaning of the term “associated values” so that the claim limitations relate more clearly to practical applications for analysis of data acquired from biological samples. As amended, the claims now clearly state that the associated values are acquired by a process where biological samples containing a plurality of genes are hybridized to one or more microarrays of probes, thus measuring the levels of mRNA or protein in the biological samples. This is how the associated values are acquired.

35 U.S.C. 101 REJECTION

Claims 1-22, 28-30, 33, 44, 46, 58 and 60 remain rejected under 35 U.S.C. 101. New Claims 65 and 66 are rejected as well as directed to non-statutory subject matter. The rejection is respectfully traversed.

The Examiner is of the opinion that the claimed invention of the rejected claims does not produce a useful, concrete and tangible result, but rather merely encompass combinations of groups of data about statistical differences in mRNA or protein levels, with no specific output that meets the concrete, tangible and useful criteria, or merely describes “functional descriptive material.” We disagree insofar as the rejection is applied to the claims as amended. We believe that one skilled in the art will think that the claimed invention of the Claims as amended produces a useful, concrete and tangible result.

As clearly explained in the present Application, one of the major problems in analysis of gene related data is that the process normally used in collecting information to find statistically significant biological phenomena in biological samples is by its very nature noisy. Thus, it is difficult to distinguish between measured variations in mRNA or protein levels due to the noise inherent in the process from variations in mRNA or protein levels caused by statistically significant biological phenomena. This is overcome in the claims as amended by deriving or providing an expected value of the parameter, where the expected value is indicative of extent of variations in the parameter introduced by the data collection process itself. Only when the observed phenomenon results in values of the parameter that are significant compared to such expected value will the gene exhibiting such characteristics be identified to be associated with statistically significant biological phenomena. This results in a better success rate of detecting *bona fide* changes in gene expression and a reduced false discovery rate in the examples

described in the present application. This limitation is present in all of the rejected Independent Claims. An embodiment of this feature is described in more detail on pages 12 –14 of the specification, and the improved results over conventional techniques on pages 15-18.

The above feature is taken one step further in claims 28, 46, 60 and 66. This is illustrated in the embodiment described on pages 12-14 of the present application. The sets of values of the relative difference $d(i)$ in expression of the genes are permuted to arrive at sets of relative difference values different from the original sets. The values in the new sets are then ranked, and an expected value of such relative difference for each rank is provided. Thus, comparing the largest relative difference among all the genes to the largest relative differences from the permutations provides one possible test for identifying genes to be of statistical significance. Therefore, the average of the largest relative differences from the permutations is the expected relative difference for such gene. A comparison of the relative difference of such gene with its expected value can be used as control as to whether statistical significance should be assigned to such gene. The same reasoning applies to the gene of the second highest relative difference and comparison to the second largest relative differences from the permutations, and so on for all the genes involved in the calculation. This process involving ranking the relative difference values further enhances the ability to identify biological phenomenon from noise inherent in the data collection and analysis.

Another difficulty in making use of microarray data is due to the fact that the expression levels of the genes have a wide range of values or scattered values. Another limitation present in some of the Claims amended solves this problem by adjusting the parameters of the plurality of genes so that variables related to the parameters are substantially independent or variations of scattered values or average associated values of the genes over the sets. The scattered values are defined by standard deviation of the associated values in the sets. In the embodiment described on pages 11 and 12 of the present specification, this is performed by adjusting the value of S_0 in equation 1 on page 10 so that the parameter $d(i)$ is substantially independent of the wide variations and scattered values or average associated values of the genes, so that all of the microarray data can be effectively used.

The above-described limitations in the Claims as amended are clearly described in the paper entitled "*Significance analysis of microarrays applied to the ionizing radiation response*," Virginia Goss Tusher, et al., Proceedings of the National Academy of Sciences of the United

States of America (PNAS), (published on line before print, April 17, 2001, 10.1073/pnas.091062498), PNAS, April 24, 2001, Volume 98, No. 9, pages 5116-5121. As will be noted, the content of this article is essentially captured in the present Application, and the inventors of this application are its authors. As referred to in the present Application and in this article, the various techniques described in the article are referred to as "SAM," and this article is referred to herein as the "SAM article."

A copy of the SAM article is attached, along with a printout of the references that cite or quote this article. Thus, there are roughly around 500 published articles that refer to the SAM paper by the inventors as of September of 2005. This makes the article one of the most cited or referred to published articles in its field of expertise. Enclosed are two of the articles that refer to the SAM article.

Attached is the article entitled "An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival," by Lance Miller et al., (published on line before print September 2, 2005, 10.1073/pnas.0506230102), PNAS, September 20, 2005, Volume 102, No. 38, pages 13550-13555. As stated on page 6 of this article: "Univariate analysis by statistical analysis of microarrays (SAM) (22) identified 6,545 Affymetrix probe sets representing $\approx 5,290$ distinct genes whose expression patterns distinguished P53 mt and wt tumors with a false discovery rate (q value) $< 1\%$ and d score (modified t statistic) > 2.0 ..., further illuminating the extensive nature of the molecular variation underlying p53 status." Reference to "(SAM)(22)" in the quote refers to the above SAM article.

SAM software that implements the features described in the present application and the SAM article has been widely licensed since 2001. Attached is a Bulletin of information available on the internet on licensing such software.

Attached is a declaration by Dr. Gilbert Chu, one of the inventors of the present application, stating that the SAM software that has been licensed to the public implements the two claim limitations in the claims as amended discussed above. Dr. Chu further states in his declaration that he believes that the analysis using SAM in the above quote from the Miller article employs the two claim limitations discussed above through the use of SAM software.

Thus, as can be seen from the above quote, the authors of the Miller, et al. article use features of the two claim limitations in the claims as amended to identify about 5290 distinct

genes whose expression patterns distinguished the p53 gene tumors. In other words, by making use of the above-described two claim limitations of the Claims as amended, without more, the authors of the Miller, et al. article identified around 5290 distinct genes whose associated values differ by an amount of statistical significance among the data, for distinguishing the p53 gene tumors. As described in this article by Miller, et al., certain correlations that are useful for analyzing human breast cancer for predicting mutation status transcriptional effects and patient survival are then developed. Thus, this article is direct evidence that to one of ordinary skill in the art, the two limitations in the Claims as amended produce a useful, concrete and tangible result. This contradicts the Examiner's positions if the rejection is applied to the claims as amended.

Another article citing or referring to the SAM article is "Ancestral antibiotic resistance in *Mycobacterium tuberculosis*," by Rowan P. Morris, et al., (published on line before print August 15, 2005, 10.1073/pans. 0505446102), PNAS, August 23, 2005, Volume 102, No. 34, pages 12200-12205. As described on page 5 of this article by Morris et al., monocytes were infected by mycobacterium and activated. Labeling of RNA and hybridizations were performed. Then, "data from each experimental condition was analyzed separately by using significance analysis of microarrays (22) with a false discovery ration $\leq 0.3\%$." The reference to "significance analysis of microarrays (22)" is to the SAM article. Dr. Chu's declaration states that he believes that the data analysis using SAM in the above quote from the Morris article uses the two claim limitations discussed above through the use of SAM software.

Thus, in each of the articles by Miller et al. and Morris et al., the two claim limitations in the claims as amended discussed above, apparently without more, allow genes whose associated values differ by an amount of statistical significance to be identified for very practical, tangible, useful and concrete applications, so that the invention in the claims as amended are recognized by those skilled in the art to produce a useful, concrete and tangible result.

The above two articles are merely two of the around 500 articles that refer to or use the techniques in the SAM article. More examples of useful, concrete and tangible results produced by application of the invention in the Claims as amended can undoubtedly be found in other articles in addition to the two described above. We therefore believe that there is ample evidence that the invention of the Claims as amended produces a useful, concrete and tangible result,

contrary to the opinion of the Examiner. If the Examiner disagrees, it is respectfully requested that the Examiner explain in detail the reasons why such rejection is still maintained in the face of overwhelming evidence that weighs against such position.

In responding to the Applicant's arguments, the Office Action stated that "in the instant Claims, the Claims as a whole, do not result in the physical transformation and as a whole do not constitute a practical application of an abstract idea" quoting *State Street Bank & Trust v. Signature Financial Group*, 47 USPQ 2d at 1600. The Office Action continues: "Thus, data transformation is not necessarily a physical transformation, as it is the result as a whole that is the focus. It is noted that the *Arrhythmia* case, for example, 'constituted a practical Application of an abstract idea because it corresponded to a useful, concrete, intangible thing – the condition of the patient's heart, which is not the case in the instant claimed invention."

We disagree with the above statement by the Examiner. The various claim elements in the rejected claims do not merely manipulate abstract concepts or data, but the associated values that are manipulated correspond to useful, concrete and tangible things – the levels of mRNA or levels of protein. As further clarified by the claim amendments herein, these levels are measured from biological samples containing the genes. These levels, may in turn, indicate certain significant biological characteristics or activity. As clearly described in the specification, such biological characteristics or activity may, for example, be caused by the effect of radiation on genes, by inducing or repressing the genes. The Examiner may again object on the grounds that there is inadequate nexus between such practical real world applications and the different claim elements in the rejected Claims. We believe, however, that there is no requirement under 35 U.S.C. § 101 that very specific applications such as gene inducement or repression need to be recited in the claims themselves. This is true in the *Arrhythmia* case as well. As will be noted from the *Arrhythmia* case, the condition of the patient's heart is also not directly recited in the Claims at issue. The Claim at issue in the *Arrhythmia* case merely recites the steps of converting QRS signals to time segments, applying a portion of the time segments in reverse time order to high pass filter means, determining an arithmetic value of the amplitude of the output of the filter and comparing the value with a pre-determined level. The patient's heart condition does not appear anywhere in the claim language. The different data that is manipulated by the method in *Arrhythmia* are an indication of the condition of the patient's heart, even though the condition of the patient's heart is not directly recited in the Claim. Analogous to the Claim in *Arrhythmia*, the

associated values that are manipulated in the Claims as amended indicate significant biological characteristics or activity. When the biological samples analyzed are samples that have been irradiated and control samples that have not irradiated, the invention of the claims as amended will reveal the genes that have been suppressed and those that have been induced. Analogous applications are also shown in the Miller and Morris articles. The claims as amended are no different from the *Arrhythmia* claims where the signals manipulated in the claimed method represent heart activity of a patient without actually reciting in the claim the condition of the patient's heart.

We therefore disagree with the Examiner's opinion quoted above.

UTILITY

Claims 1-22, 28-30, 33, 44, 46, 58, 60, 65 and 66 remain rejected under 35 U.S.C. 101 because the claimed invention lacks patentable utility. The rejection is respectfully traversed. The discussion above clearly describes many utilities of the invention in the rejected claims. Withdrawal of this rejection is respectfully requested.

In the amendment mailed January 19, 2005, Applicants set forth a number of specific Utilities on page 12 of the Amendment. These include the identification of genes whose DNA has been damaged by exposure to radiation, the identification of genes in tumors (page 18, line 24) or the identification of genes whose expression correlates with the survival time of patients (page 19, line 16), or with tumor stage (sentence bridging pages 19 and 20). These Utilities were rejected by the Examiner in the Office Action mailed April 19, 2005 on the ground that the Claims do not recite steps that are applicable to any of these uses and that there is no adequate nexus between the disclosed subject matter and these asserted Utilities. We disagree.

As will be evident from the present Application, as well as the articles described above, the utility of the invention of the rejected Claims is achieved by simply applying the steps involved to particular biological samples. In the present Application, the recited features were applied to samples that were grown with exposure to radiation and samples that were grown without exposure to radiation. The invention of the claims as amended then enables the identification of genes whose expression has been induced or repressed by radiation. The same is true in the case of the above articles involving tumor and breast cancer analysis. This is true

also in the case of the Claims in *Arrhythmia Research Technology, Inc. v. Corazonix Corp.*, 22 U.S.P.Q. 2d, 1033 at 1038. Claim 1 in the *Arrhythmia* case recites steps that merely manipulate QRS signals which represent signals obtained from the heart, but otherwise do not recite anything involving the patient's heart condition. The Court nevertheless deems such claim to have utility under 35 U.S.C. 101. In the same manner, it is believed that the Claims as amended have utility, without having to recite specific applications, which would unduly limit the claims.

By rejecting the Claims based on the inadequate nexus between the disclosed subject matter and the asserted utilities, the Examiner is in fact requiring Applicants to restrict the Claims to particular applications. This is against the rule articulated by the Court of Appeals for the Federal Circuit in *State Street Bank & Trust v. Signature Financial Group*, 47 USPQ 2d 1600 at 1604. In the ruling by the lower court in this case, the patent was found invalid "because the '056 patent is claimed [sic] sufficiently broadly to foreclose any computer-implemented accounting method necessary to manage this type of financial structure." In response, the Court of Appeals for the Federal Circuit stated as follows: "whether the patent's Claims are too broad to be patentable is not to be judged under Section 101, but rather under Sections 102, 103, and 112. Assuming the above statement to be correct, it has nothing to do with what is claimed is statutory subject matter."

Rejection under 35 U.S.C. 112 First Paragraph


Claims 1-22, 28-30, 33, 44, 46, 58, 60 and new Claims 65 and 66 are rejected under 35 U.S.C. 112 first paragraph for failing to comply with the written description requirement. Specifically, Claims 1, 28, 46, 58 and 60, 64 and 65 are rejected on the ground that it still includes the terms "protein." We disagree. As expressly stated in lines 1-5 on page 4 of the specification, one of the examples given for of the values associated with genes are the levels of protein encoded by the genes. This is also explicitly claimed in Claim 4 of the Claims as originally filed. As noted in MPEP 2163, page 2100-173, there is a strong presumption that an adequate written description of the claimed invention is present when the Application is filed. Further, the PTO has the initial burden of presenting evidence or reasons why persons skilled in the art would not recognize in the disclosure a description of the invention defined by the claims. The Examiner has failed to present any evidence why persons skilled in the art would not recognize a description of the invention originally present both in the Claims and the Summary

of the Invention and that the claims still fail to comply with 35 U.S.C. 112 for lacking an adequate written description. It is respectfully requested that this rejection be withdrawn.

CONCLUSION

In view of the amendments and remarks contained herein, it is believed that all claims are in condition for allowance and an indication of their allowance is requested. However, if the Examiner is aware of any additional matters that should be discussed, a call to the undersigned attorney at: (415) 318-1162 would be appreciated.

Respectfully submitted,



James S. Hsue
Reg. No. 29,545

July 20, 2006

Date

PARSONS HSUE & de RUNTZ LLP
595 Market Street, Suite 1900
San Francisco, California 94105
Telephone: 415.318.1160 (main)
Telephone: 415.318.1162 (direct)
Fax: 415.693.0194

Keystone Symposia

Sign up for PNAS Online Tools

Get notified by email when new content goes on line

[Info for Authors](#) | [Editorial Board](#) | [About](#) | [Subscribe](#) | [Advertise](#) | [Contact](#) | [Site Map](#)

PNAS

Proceedings of the National Academy of Sciences of the United States of America

[Current Issue](#)

[Archives](#)

[Online Submission](#)

GO

[advanced search >>](#)

**Published online before print April 17, 2001,
10.1073/pnas.091062498**

PNAS | April 24, 2001 | vol. 98 | no. 9 | 5116-5121

Statistics / Genetics

Significance analysis of microarrays applied to the ionizing radiation response

**Virginia Goss Tusher^{*}, Robert Tibshirani[†], and
Gilbert Chu^{*,†}**

^{*} Departments of Medicine and Biochemistry, Stanford University, 269 Campus Drive, Center for Clinical Sciences Research 1115, Stanford, CA 94305-5151; and [†] Department of Health Research and Policy and Department of Statistics, Stanford University, Stanford, CA 94305

Communicated by Bradley Efron, Stanford University, Stanford, CA, February 6, 2001 (received for review December 1, 2000)

This Article

- ▶ [Abstract](#) FREE
- ▶ [Full Text \(PDF\)](#)
- ▶ [An erratum has been published](#)
- ▶ [An erratum has been published](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)
- ▶ [Citation Map](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in ISI Web of Science](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Add to My File Cabinet](#)
- ▶ [Download to citation manager](#)
- ▶ [Cited by other online articles](#)
- ▶ [Search for citing articles in: ISI Web of Science \(1076\)](#)
- ▶ [Request Copyright Permission](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Tusher, V. G.](#)
- ▶ [Articles by Chu, G.](#)

▶ Abstract

▲ [Top](#)

Microarrays can measure the expression of thousands of genes to identify changes in expression between different biological states. Methods are needed to determine the significance of these changes while accounting for the enormous number of genes. We describe a method, Significance Analysis of Microarrays (SAM), that assigns a score to each gene on the basis of change in gene expression relative to the standard deviation of repeated measurements. For genes with scores greater than an adjustable threshold, SAM uses permutations of the repeated measurements to estimate the percentage of genes identified by chance, the false discovery rate (FDR). When the transcriptional response of human cells to ionizing radiation was measured by microarrays, SAM identified 34 genes that changed at least 1.5-fold with an estimated FDR of 12%, compared with FDRs of 60 and 84% by using conventional methods of analysis. Of the 34 genes, 19 were involved in cell cycle regulation and 3 in apoptosis. Surprisingly, four nucleotide excision repair genes were induced, suggesting that this repair pathway for UV-damaged DNA might play a previously unrecognized role in repairing DNA damaged by ionizing radiation.

- **Abstract**
- ▼ **Introduction**
- ▼ **Materials and Methods**
- ▼ **Results**
- ▼ **Discussion**
- ▼ **References**

► **Introduction**

DNA microarrays contain oligonucleotide or cDNA probes for measuring the expression of thousands of genes in a single hybridization experiment. Although massive amounts of data are generated, methods are needed to determine whether changes in gene expression are experimentally significant. Cluster analysis of microarray data can find coherent patterns of gene expression (1) but provides little information about statistical significance. Methods based on conventional t tests provide the probability (P) that a difference in gene expression occurred by chance (2, 3). Although

- ▲ **Top**
- ▲ **Abstract**
- **Introduction**
- ▼ **Materials and Methods**
- ▼ **Results**
- ▼ **Discussion**
- ▼ **References**

$P = 0.01$ is significant in the context of experiments designed to evaluate small numbers of genes, a microarray experiment for 10,000 genes would identify 100 genes by chance. This problem led us to develop a statistical method adapted specifically for microarrays, Significance Analysis of Microarrays (SAM).

SAM identifies genes with statistically significant changes in expression by assimilating a set of gene-specific t tests. Each gene is assigned a score on the basis of its change in gene expression relative to the standard deviation of repeated measurements for that gene. Genes with scores greater than a threshold are deemed potentially significant. The percentage of such genes identified by chance is the false discovery rate (FDR). To estimate the FDR, nonsense genes are identified by analyzing permutations of the measurements. The threshold can be adjusted to identify smaller or larger sets of genes, and FDRs are calculated for each set. To demonstrate its utility, SAM was used to analyze a biologically important problem: the transcriptional response of lymphoblastoid cells to ionizing radiation (IR).

► Materials and Methods

Preparation of RNA. Human lymphoblastoid cell lines GM14660 and GM08925 (Coriell Cell Repositories, Camden, NJ) were seeded at 2.5×10^5 cells/ml and exposed to IR 24 h later. RNA was isolated, labeled, and hybridized to the HUGENEFL GENECHIP microarray according to manufacturer's protocols (Affymetrix, Santa Clara, CA).

- ▲ [Top](#)
- ▲ [Abstract](#)
- ▲ [Introduction](#)
- [Materials and Methods](#)
- ▼ [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

Microarray Hybridization. Each gene in the microarray was represented by 20 oligonucleotide pairs, each pair consisting of an oligonucleotide perfectly matched to the cDNA sequence, and a second oligonucleotide containing a single base mismatch. Because gene expression was computed from differences in hybridization to the matched and mismatched probes, expression levels were sometimes reported by

the GENECHIP ANALYSIS SUITE software as negative numbers.

Northern Blot Hybridization. Total RNA (15 μ g) was resolved by agarose gel electrophoresis, transferred to a nylon membrane, and hybridized to specific radiolabeled DNA probes, which were prepared by PCR amplification.

► Results


RNA was harvested from wild-type human lymphoblastoid cell lines, designated 1 and 2, growing in an unirradiated state (U) or in an irradiated state (I) 4 h after exposure to a modest dose of 5 Gy of IR. RNA samples were labeled and divided into two identical aliquots for independent hybridizations, A and B. Thus, data for 6,800 genes on the microarray were generated from eight hybridizations (U1A, U1B, U2A, U2B, I1A, I1B, I2A, and I2B).

- ▲ [Top](#)
- ▲ [Abstract](#)
- ▲ [Introduction](#)
- ▲ [Materials and Methods](#)
- [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

We scaled the data from different hybridizations as follows. A reference data set was generated by averaging the expression of each gene over all eight hybridizations. The data for each hybridization were compared with the reference data set in a cube root scatter plot. We chose the cube root scatter plot because it resolved the vast majority of genes that are expressed at low levels and permitted the inclusion of negative levels of expression that are sometimes generated by the GENECHIP software. A linear least-squares fit to the cube root scatter plot was then used to calibrate each hybridization.

After scaling, a linear scatter plot was generated for average gene expression in the four A aliquots (U1A, I1A, U2A, and U2A) vs. the average in the four B aliquots (U1B, I1B, U2B, and U2B), a partitioning of the data that eliminates biological changes in gene expression (Fig. [1A](#)). The linear scatter plot confirmed that the data were generally reproducible but failed to resolve genes expressed at low levels. Better resolution of these genes was achieved by the cube root scatter plot (Fig. [1B](#)), which

revealed three salient features: the large percentage of genes (24%) assigned negative levels of expression, the large percentage of genes with low levels of expression, and the low signal-to-noise ratio at low levels of expression.



View larger version

(13K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 1. Gene expression measured by microarrays. (A) Linear scatter plot of gene expression. Each gene (i) in the microarray is represented by a point with coordinates consisting of average gene expression measured from the four A hybridizations ($\text{avg } x_A$) and the average gene expression in the four B hybridizations ($\text{avg } x_B$). (B) Cube root scatter plot of gene expression. The average gene expression from the A and B hybridizations have been plotted on a cube root scale to resolve genes expressed at low levels. (C) Cube root scatter plot of average gene expression from the four hybridizations with uninduced cells ($\text{avg } x_U$) and induced cells 4 h after exposure to 5 Gy of IR ($\text{avg } x_I$). Some of the genes that responded to IR are indicated by arrows.

To assess the biological effect of IR, a scatter plot was generated for average gene expression in the four irradiated states vs. the four unirradiated states (compare Fig. 1 B and C). A few of the potentially significant changes in gene expression are indicated by arrows in Fig. 1C, but the effect was not easily quantified, and a method was needed to identify changes with statistical confidence.

Our approach was based on analysis of random fluctuations in the data. In general, the signal-to-noise ratio decreased with decreasing gene expression (Fig. 1). However, even for a given level of expression, we found that fluctuations were gene specific. To account for gene-specific fluctuations, we defined a statistic based on the ratio of change in gene expression to standard deviation in the data for that gene. The

"relative difference" $d(i)$ in gene expression is:

$$d(i) = \frac{\bar{x}_I(i) - \bar{x}_U(i)}{s(i) + s_0} \quad [1]$$

where $\bar{x}_I(i)$ and $\bar{x}_U(i)$ are defined as the average levels of expression for gene (i) in states I and U, respectively. The "gene-specific scatter" $s(i)$ is the standard deviation of repeated expression measurements:


$$s(i) = \sqrt{a \left\{ \sum_m [x_m(i) - \bar{x}_I(i)]^2 + \sum_n [x_n(i) - \bar{x}_U(i)]^2 \right\}} \quad [2]$$

where \sum_m and \sum_n are summations of the expression measurements in states I and U, respectively, $a = (1/n_1 + 1/n_2)/(n_1 + n_2 - 2)$, and n_1 and n_2 are the numbers of measurements in states I and U (four in this experiment).

To compare values of $d(i)$ across all genes, the distribution of $d(i)$ should be independent of the level of gene expression. At low expression levels, variance in $d(i)$ can be high because of small values of $s(i)$. To ensure that the variance of $d(i)$ is independent of gene expression, we added a small positive constant s_0 to the denominator of Eq. 1. The coefficient of variation of $d(i)$ was computed as a function of $s(i)$ in moving windows across the data. The value for s_0 was chosen to minimize the coefficient of variation. For the data in this paper, this computation yielded $s_0 = 3.3$.

Scatter plots of $d(i)$ vs. $s(i)$ are shown in Fig. 2. The scatter plot for relative difference between states I and U is shown in Fig. 2A. By contrast, the scatter plot for relative difference between cell lines 1 and 2 shows more marked changes in Fig. 2B. These relative differences exceeded random fluctuations in the data, as measured by the relative difference between hybridizations A and B in Fig. 2C.

Fig. 2. Scatter plots of relative difference in gene



**View larger
version (32K):**

[\[in this window\]](#)

[\[in a new window\]](#)


expression $d(i)$ vs. gene-specific scatter $s(i)$. The data were partitioned to calculate $d(i)$, as indicated by the bar codes. The shaded and unshaded entries were used for the first and second terms in the numerator of $d(i)$ in Eq. 1. (A) Relative difference between irradiated and unirradiated states. The statistic $d(i)$ was computed from expression measurements partitioned between irradiated and unirradiated cells. (B) Relative difference between cell lines 1 and 2. The statistic $d(i)$ was computed from expression measurements partitioned between cell lines 1 and 2. (C) Relative difference between hybridizations A and B. The statistic $d(i)$ was computed from the permutation in which the expression measurements were partitioned between the equivalent hybridizations A and B. (D) Relative difference for a permutation of the data that was balanced between cell lines 1 and 2.

Although the relative difference computed from hybridizations A and B provided a control for random fluctuations, additional controls were needed to assign statistical significance to the biological effect of IR. Instead of performing more experiments, which are expensive and labor intensive, we generated a large number of controls by computing relative differences from permutations of the hybridizations for the four irradiated and four unirradiated states. To minimize potentially confounding effects from differences between the two cell lines, we analyzed the data by using the 36 permutations that were balanced for cell lines 1 and 2. Permutations were defined as balanced when each group of four experiments contained two experiments from cell line 1 and two experiments from cell line 2. Fig. 2 C and D are examples of balanced permutations.

To find significant changes in gene expression, genes were ranked by magnitude of their $d(i)$ values, so that $d(1)$ was the largest relative difference, $d(2)$ was the second largest relative difference, and $d(i)$ was the i th largest relative difference. For each of the 36 balanced permutations, relative differences $d_p(i)$ were also calculated, and the

genes were again ranked such that $d_p(i)$ was the i th largest relative difference for permutation p . The expected relative difference, $d_E(i)$, was defined as the average over the 36 balanced permutations, $d_E(i) = \sum_p d_p(i)/36$.

To identify potentially significant changes in expression, we used a scatter plot of the observed relative difference $d(i)$ vs. the expected relative difference $d_E(i)$ (Fig. 3A). For the vast majority of genes, $d(i) \approx d_E(i)$, but some genes are represented by points displaced from the $d(i) = d_E(i)$ line by a distance greater than a threshold Δ . For example, the threshold $\Delta = 1.2$ illustrated by the broken lines in Fig. 3A yielded 46 genes that were "called significant." These 46 genes are shown in the context of the scatter plot for $d(i)$ vs. $s(i)$ (Fig. 3B) and in the scatter plot for the cube root of gene expression $\bar{x}_I(i)$ vs. $\bar{x}_U(i)$ (Fig. 3C). Genes identified by $d(i)$ do not necessarily have the largest changes in gene expression.



View larger version

(21K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 3. Identification of genes with significant changes in expression. (A) Scatter plot of the observed relative difference $d(i)$ versus the expected relative difference $d_E(i)$. The solid line indicates the line for $d(i) = d_E(i)$, where the observed relative difference is identical to the expected relative difference. The dotted lines are drawn at a distance $\Delta = 1.2$ from the solid line. (B) Scatter plot of $d(i)$ vs. $s(i)$. (C) Cube root scatter plot of average gene expression in induced and uninduced cells. The cutoffs for 2-fold induction and repression are indicated by the dashed lines. In A-C, the 46 potentially significant genes for $\Delta = 1.2$ are indicated by the squares.

To determine the number of falsely significant genes generated by SAM, horizontal

cutoffs were defined as the smallest $d(i)$ among the genes called significantly induced and the least negative $d(i)$ among the genes called significantly repressed. The number of falsely significant genes corresponding to each permutation was computed by counting the number of genes that exceeded the horizontal cutoffs for induced and repressed genes. The estimated number of falsely significant genes was the average of the number of genes called significant from all 36 permutations. For $\Delta = 1.2$, the permuted data sets generated an average of 8.4 falsely significant genes, compared with 46 genes called significant, yielding an estimated FDR of 18% (Table 1). As Δ decreased, the number of genes called significant by SAM increased but at the cost of an increasing FDR. (Omitting s_0 from Eq. 1 produced higher FDRs of 45, 35, and 28% for $\Delta = 0.6, 0.9$, and 1.2.)

View this table: **Table 1.** Comparison of methods for identifying changes in gene expression
[\[in this window\]](#)
[\[in a new window\]](#)

Our method for setting thresholds provides asymmetric cutoffs for induced and repressed genes. The alternative is the standard t test, which imposes a symmetric horizontal cutoff, with $d(i) > c$ for induced genes and $d(i) < -c$ for repressed genes. However, the asymmetric cutoff is preferred because it allows for the possibility that $d(i)$ for induced and repressed genes may behave differently in some biological experiments.

SAM proved to be superior to conventional methods for analyzing microarrays (Table 1 and Fig. 4A). First, SAM was compared with the approach of identifying genes as significantly changed if an R -fold change was observed. In this "fold change" method, $r(i) = \bar{x}_I(i)/\bar{x}_U(i)$, and gene (i) was called significantly changed if $r(i) > R$ or $r(i) < 1/R$.

To permit computation of $r(i)$ from negative values for gene expression, $\bar{x}_I(i)$ and $\bar{x}_U(i)$

were converted to 10 when their values were negative or less than 10. The results of this procedure yielded unacceptably high FDRs of 73-84%.



View larger version

(13K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 4. Comparison of SAM to conventional methods for analyzing microarrays. (A) Falsely significant genes plotted against number of genes called significant. Of the 57 genes most highly ranked by the fold change method, 5 were included among the 46 genes most highly ranked by SAM. Of the 38 genes most highly ranked by the pairwise fold change method, 11 were included among the 46 genes most highly ranked by SAM. These results were consistent with the FDR of SAM compared to the FDRs of the fold change and pairwise fold change methods. (B) Northern blot validation for genes identified by the fold change method. Values of $r(i)$ are plotted for genes chosen at random from the 57 genes most highly ranked by the fold change method. (C) Validation for genes identified by SAM. Results are plotted for genes chosen at random from the 46 genes most highly ranked by SAM. Genes analyzed by Northern blot are represented by circles. $TNF-\alpha$ was validated by using a PreDeveloped TaqMan assay (PE Biosystems) and is represented by a square. The straight lines in B and C indicate the position of exact agreement between Northern blot and microarray results.

Another approach attempts to account for uncertainty in the data by identifying genes as significantly changed if an R -fold change is observed consistently between paired samples (4). To apply this "pairwise fold change" method to our four data sets before IR and four data sets after IR, changes in gene expression were declared significant if 12 of 16 pairings satisfied the criteria $r(i) > R$ or $r(i) < 1/R$. Despite the demand for consistent changes between paired samples, this method yielded FDRs of 60-71%.

To understand why fold-change methods fail, note that the vast majority of genes are expressed at low levels where the signal-to-noise ratio is very low (Fig. 3C). Thus, 2-fold changes in gene expression occur at random for a large number of genes.

Conversely, for higher levels of expression, smaller changes in gene expression may be real, but these changes are rejected by fold-change methods. The pairwise fold-change method provides modest improvement but remains inferior to SAM.

Of the 46 genes most highly ranked by SAM ($\Delta = 1.2$), 36 increased or decreased at least 1.5-fold ($R = 1.5$). The number of falsely significant genes that met these two criteria was 4.5, corresponding to a FDR of 12% (Table 1). Fas was identified three times as alternately spliced forms, leaving 34 independent genes (Table 2). As an indication of biological validity, 10 of the 34 genes have been reported in the literature as part of the transcriptional response to IR. $\text{TNF-}\alpha$ was reported to be induced by other investigators (5) but was repressed here. Quantitative reverse transcription-PCR confirmed this result.

View this table: **Table 2.** Genes with changes in expression called significant by SAM
[\[in this window\]](#)
[\[in a new window\]](#)

To test the validity of SAM directly, we performed Northern blots for genes that were randomly selected from the 46 and 57 genes most highly ranked by SAM ($\Delta = 1.2$) and the fold-change method (at least 3.6-fold change), respectively. Northern blots showed little correlation with the genes identified by the fold change method (Fig. 4B), but strong correlation with the genes identified by SAM (Fig. 4C). Indeed, Northern blots contradicted only 1 (maxiK) of 11 genes identified by SAM, consistent with our estimated FDR.

Nineteen of the 34 genes most highly ranked by SAM appear to be involved in the cell

cycle. Three are known to be induced in a p53-dependent manner: p21, cyclin G1, and mdm2 (6-8). Six cell cycle genes were repressed: E2-EPF, p55cdc, cyclin B, ckshs2, cdc25, and wee1 (9, 10). Five genes encoding the mitotic machinery were also repressed: PLK-1, MKLP-1, MCAK, C-TAK1, CENP-E (11-13). Three genes involved in cell proliferation were induced or repressed: PTP(CAAX1), LPAP, and c-myc (14-18). Some responses appeared paradoxical. For example, cdc25 phosphatase and wee1 kinase have antagonistic effects on the phosphorylation state of cdc2, but both genes were repressed. Repression of these genes together with the mitotic genes may represent a damage response that dismantles the cell cycle machinery until the cell has repaired the damaged DNA.

Four of the 34 genes play roles in DNA repair, but none are involved in the repair of IR-induced double-strand breaks. Instead, the genes (p48, XPC, gadd45, PCNA) have roles in nucleotide excision repair, a pathway conventionally associated with UV-induced damage (19-22). We confirmed the induction of these genes by Northern blot (23-25). Fornace *et al.* reported defective removal of base damage induced by IR in xeroderma pigmentosum cells (26). Leadon *et al.* reported that a novel DNA repair pathway involving long excision repair patches of at least 150 nucleotides is activated by IR but not UV (27). Our results suggest that this novel pathway might include p48, XPC, gadd45, and PCNA.

Four of the 34 genes play roles in apoptosis (Fas, bbc3, TNF- α , OX40 ligand). The remaining genes may have previously unsuspected roles in the DNA damage response or may be among the estimated set of four falsely detected genes.

The 34 genes most highly ranked by SAM are only a subset of all of the genes that change 1.5-fold with IR. Indeed, we calculated the difference between the number of genes called significant and the number of falsely significant genes for decreasing $\Delta = 0.3, 0.2$, and 0.1 , and found the differences to be 92, 170, and 184, respectively. Thus, SAM suggests that approximately 180 of the 6,800 genes on the microarray

were induced or repressed by 5 Gy IR.

► Discussion

SAM is a method for identifying genes on a microarray with statistically significant changes in expression, developed in the context of an actual biological experiment. SAM was successful in analyzing this experiment as well as several other experiments with oligonucleotide and cDNA microarrays (data not shown).

- ▲ **Top**
- ▲ **Abstract**
- ▲ **Introduction**
- ▲ **Materials and Methods**
- ▲ **Results**
- **Discussion**
- ▼ **References**

In the statistics of multiple testing (28-30), the family-wise error rate (FWER) is the probability of at least one false positive over the collection of tests. The Bonferroni method, the most basic method for bounding the FWER, assumes independence of the different tests. An acceptable FWER could be achieved for our microarray data only if the corresponding threshold was set so high that no genes were identified. The step-down correction method of Westfall and Young (29), adapted for microarrays by Dudoit *et al.* (<http://www.stat.berkeley.edu/users/terry/zarray/Html/matt.html>), allows for dependent tests but still remains too stringent, yielding no genes from our data.

Westfall and Young (29) define "weak control" to be control of the FWER when all of the null hypotheses are true (i.e., when there are no changes in gene expression). "Strong control" is control of the FWER when any subset of the null hypotheses is true. Under certain conditions, weak control implies strong control. In fact, the step-down correction method exerts both weak and strong control.

The method of Benjamini and Hochberg (31) assumes independent tests and guarantees an upper bound for the FDR (with both weak and strong control) by a step-up or step-down procedure applied to the individual P values. For our data, the P value for each gene is calculated from permutations of the eight experiments. Because of the limited number of permutations, the FDR is too "granular", and we identified

either zero or 300 significant genes, depending on how the P value was defined. A similar granular result was obtained for the adaptation to dependent tests by Benjamini *et al.* [*The Control of the False Discovery Rate in Multiple Testing Under Dependency* (Department of Statistics and Operations Research, Tel Aviv University, Tel Aviv). <http://www.math.tau.ac.il/~ybenja/>].

SAM does not have strong or weak control of the FWER. Instead, SAM provides an estimate of the FDR for each value of the tuning parameter Δ . The estimated FDR is computed from permutations of the data and hence assumes that all null hypotheses are true, allowing for the possibility of dependent tests. It seems plausible that this estimated FDR approximates the strongly controlled FDR when any subset of null hypotheses is true. However, we have not proven this in general. It is possible for SAM to give an estimate of the FDR that is greater than 1. However, this has not occurred in our experience. Indeed, SAM provides a reasonably accurate estimate for the true FDR. To confirm this, we constructed artificial data sets in which a subset of genes was induced over a background of noise. SAM successfully identified the induced genes and estimated the FDR with reasonable accuracy.

Although this paper analyzes a simple two-state experiment, SAM can be generalized to other types of experiments by defining $d(i)$ in a different way. Suppose the data includes gene expression $x_j(i)$ and a response parameter y_j in which $i = 1, 2, \dots, m$ genes, $j = 1, 2, \dots, n$ states. The generalized statistical parameter still takes the form $d(i) = r(i)/[s(i) + s_0]$, except that the definitions of $r(i)$ and $s(i)$ change.

To identify genes with changes in expression in an experiment with three or more states, the parameter $d(i)$ is defined in terms of the Fisher's linear discriminant. One goal might be to identify genes whose expression in one type of tumor is different from its expression in other types of tumors. Suppose that a set of n samples consists of K nonoverlapping subsets, such that the response parameter $y_{j \in \{1, \dots, K\}}$. Define C

$(k) = \{j: y_j = k\}$. Let n_k = number of observations in $C(k)$. The average gene expression in each subset is $\bar{x}_k(i) = \sum_{j \in C(k)} x_j(i)/n_k$ and the average gene expression for all n samples is $\bar{x}(i) = \sum_j x_j(i)/n$. Then define:

$$r(i) = \{ [\sum_k n_k / \Pi_k n_k] \sum_k n_k [\bar{x}_k(i) - \bar{x}(i)]^2 \}^{1/2} \quad [3]$$

$$s(i) = \{ [\sum_k (1/n_k) / \sum_k (n_k - 1)] \sum_k \sum_{j \in C(k)} [x_j(i) - \bar{x}_k(i)]^2 \}^{1/2} \quad [4]$$

SAM can be adapted for still other types of experimental data. For example, to identify genes whose expression correlates with survival time, $d(i)$ is defined in terms of Cox's proportional hazards function, in which some of the patients remain alive or are lost to follow-up at the time of the study. To identify genes whose expression correlates with a quantitative parameter, such as tumor stage, $d(i)$ can be defined in terms of the Pearson correlation coefficient. Another example includes the definition of $d(i)$ for paired data, such as gene expression in tumors before and after chemotherapy. In each case, the FDR is estimated by random permutation of the data for gene expression among the different experimental arms, i.e., permutations among the n arms of y_j . Thus, SAM is a robust and straightforward method that can be adapted to a broad range of experimental situations. SAM and the adaptations discussed above are available for use at <http://www-stat-class.stanford.edu/SAM/SAMServlet>.

► Acknowledgements

We thank Peter Jackson, Ron Davis, James Ferrell, Dean Felsher, Lisa DeFazio, Joe Budman, Jean Tang, Tom Tan, and Kerri Rieger for helpful discussions. This work was supported by the Burroughs Wellcome Clinical Scientist Award and by National Institutes of Health (NIH) Grant CA77302 to G.C., by NIH Small Business Technology Transfer grant CA75675 to G.C. and Affymetrix, and by the Stanford Genome Training Grant to V.T.

► Abbreviations

SAM, significance analysis of microarrays; FDR, false discovery rate; IR, ionizing radiation; FWER, family-wise error rate.

► Footnotes

† To whom reprint requests should be addressed. E-mail: chu@cmgm.stanford.edu .

► References

▲ [Top](#)
▲ [Abstract](#)
▲ [Introduction](#)
▲ [Materials and Methods](#)

▲ [Results](#)
▲ [Discussion](#)
• [References](#)

1. Eisen, M. , Spellman, P. , Brown, P. & Botstein, D. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 14863-14868[[Abstract/Free Full Text](#)].
2. Roberts, C. , Nelson, B. , Marton, M. , Stoughton, R. , Meyer, M. , Bennett, H. , He, Y. , Dai, H. , Walker, W. , Hughes, T. , Tyers, M. , Boone, C. & Friend, S. (2000) *Science* **287**, 873-880[[Abstract/Free Full Text](#)].
3. Galitski, T. , Saldanha, A. , Styles, C. , Lander, E. & Fink, G. (1999) *Science* **285**, 251-254[[Abstract/Free Full Text](#)].
4. Ly, D. , Lockhart, D. , Lerner, R. & Schultz, P. (2000) *Science* **287**, 2486-2492 [[Abstract/Free Full Text](#)].
5. Weill, D. , Gay, F. , Tovey, M. & Chouaib, S. (1996) *J. Interferon Cytokine Res.* **16**, 395-402[[ISI](#)][[Medline](#)] [[Order article via Infotrieve](#)].
6. Harper, J. W. , Adami, G. R. , Wei, N. , Keyomarsi, K. & Elledge, S. J. (1993) *Cell* **75**, 805-816[[ISI](#)][[Medline](#)] [[Order article via Infotrieve](#)].
7. Okamoto, K. & Beach, D. (1994) *EMBO J.* **13**, 4816-4822[[Abstract](#)].
8. Prives, C. (1998) *Cell* **95**, 5-8[[ISI](#)][[Medline](#)] [[Order article via Infotrieve](#)].

9. Furnari, B. , Rhind, N. & Russell, P. (1997) *Science* **277**, 1495-1497
[Abstract/Free Full Text].
10. Liu, Z. , Diaz, L. , Haas, A. & Giudice, G. (1992) *J. Biol. Chem.* **267**, 15829-15835
[Abstract/Free Full Text].
11. Lee, K. , Yuan, Y. , Kuriyama, R. & Erikson, R. (1995) *Mol. Cell. Biol.* **15**, 7143-7151[Abstract].
12. Maney, T. , Hunter, A. , Wagenbach, M. & Wordeman, L. (1998) *J. Cell. Biol.* **142**, 787-801[Abstract/Free Full Text].
13. Wood, K. , Sakowicz, R. , Goldstein, L. & Cleveland, D. (1997) *Cell* **91**, 357-366
[ISI][Medline] [Order article via Infotrieve].
14. Ding, I. , Bruyns, E. , Li, P. , Magada, D. , Paskind, M. , Rodman, L. , Seshadri, T. , Alexander, D. , Giese, T. & Schraven, B. (1999) *Eur. J. Immunol.* **29**, 3956-3961
[CrossRef][ISI][Medline] [Order article via Infotrieve].
15. Cates, C. , Michael, R. , Stayrook, K. , Harvey, K. , Burke, Y. , Randall, S. , Crowell, P. & Crowell, D. (1996) *Cancer Lett.* **110**, 49-55[CrossRef][ISI][Medline]
[Order article via Infotrieve].
16. Godfrey, W. , Fagnoni, R. , Harara, M. , Buck, D. & Engleman, E. (1994) *J. Exp. Med.* **180**, 757-762[Abstract].
17. Lord, J. , McIntosh, B. , Greenberg, P. & Nelson, B. (2000) *J. Immunol.* **164**, 2533-2541[Abstract/Free Full Text].
18. Prevot, D. , Voeltzel, T. , Birot, A. , Morel, A. , Rostan, M. , Magaud, J. & Corbo, L. (2000) *J. Biol. Chem.* **275**, 147-153[Abstract/Free Full Text].
19. Aboussekhra, A. , Biggerstaff, M. , Shivji, M. , Vilpo, J. , Moncollin, V. , Podust, V. , Protic, M. , Hubscher, U. , Egly, J. & Wood, R. (1995) *Cell* **80**, 859-868[ISI]
[Medline] [Order article via Infotrieve].
20. Smith, M. , Ford, J. , Hollander, M. , Bortnick, R. , Amundson, S. , Seo, Y. , Deng, C. , Hanawalt, P. & Fornace, A. J. (2000) *Mol. Cell. Biol.* **20**, 3705-3714
[Abstract/Free Full Text].
21. Sugasawa, K. , Ng, J. , Masutani, C. , Iwai, S. , van der Spek, P. , Eker, A. , Hanaoka, F. , Bootsma, D. & Hoeijmakers, J. (1998) *Mol. Cell* **2**, 223-232[ISI]
[Medline] [Order article via Infotrieve].
22. Tang, J. , Hwang, B. , Ford, J. , Hanawalt, P. & Chu, G. (2000) *Mol. Cell* **5**, 737-744[ISI][Medline] [Order article via Infotrieve].
23. Kastan, M. , Zhan, Q. , El-Deiry, F. , Jacks, T. , Walsh, W. , Plunkett, B. , Vogelstein, B. & Fornace, A. (1992) *Cell* **71**, 587-597[ISI][Medline]

[[Order article via Infotrieve](#)].

24. Hwang, B. J. , Ford, J. , Hanawalt, P. C. & Chu, G. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 424-428[[Abstract/Free Full Text](#)].
 25. Xu, J. & Morris, G. (1999) *Mol. Cell. Biol.* **19**, 12-20[[Abstract/Free Full Text](#)].
 26. Fornace, A. , Dobson, P. & Kinsella, T. (1986) *Radiat. Res.* **106**, 73-77[[ISI](#)][[Medline](#)] [[Order article via Infotrieve](#)].
 27. Leadon, S. , Dunn, A. & Ross, C. (1996) *Radiat. Res.* **146**, 123-130[[ISI](#)][[Medline](#)] [[Order article via Infotrieve](#)].
 28. Hochberg, Y. & Tamhane, A. (1987) *Multiple Comparisons Procedures* (Wiley, New York).
 29. Westfall, P. & Young, S. (1993) *Resampling-Based Multiple Testing* (Wiley, New York).
 30. Hsu, J. (1996) *Multiple Comparisons: Theory and Methods* (Chapman & Hall, London).
 31. Benjamini, Y. & Hochberg, Y. (1995) *J. R. Stat. Soc. B* **57**, 289-300[[ISI](#)].
- www.pnas.org/cgi/doi/10.1073/pnas.091062498

This article has been cited by other articles in [HighWire Press](#)-hosted journals:



Proceedings of the National Academy of Sciences

[HOME](#)

Z. Xie, E. Allen, A. Wilken, and J. C. Carrington
DICER-LIKE 4 functions in trans-acting small interfering RNA biogenesis and vegetative phase change in *Arabidopsis thaliana*

PNAS, September 6, 2005; 102(36): 12984 - 12989.

[[Abstract](#)] [[Full Text](#)] [[PDF](#)]



Proceedings of the National Academy of Sciences

[HOME](#)

A. I. Goranov, L. Katz, A. M. Breier, C. B. Burge, and A. D. Grossman
A transcriptional response to replication status mediated by the conserved bacterial replication protein DnaA

PNAS, September 6, 2005; 102(36): 12932 - 12937.

[[Abstract](#)] [[Full Text](#)] [[PDF](#)]



Proceedings of the National Academy of Sciences


► HOME

J. D. Storey, W. Xiao, J. T. Leek, R. G. Tompkins, and R. W. Davis

Significance analysis of time course microarray experiments

PNAS, September 6, 2005; 102(36): 12837 - 12842.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory and Critical Care Medicine

► HOME

A. C. Borczuk, H. K. Kim, H. A. Yegen, R. A. Friedman, and C. A. Powell

Lung Adenocarcinoma Global Profiling Identifies Type II Transforming Growth Factor- β Receptor as a Repressor of Invasiveness

Am. J. Respir. Crit. Care Med., September 15, 2005; 172(6): 729 - 737.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Clinical Investigation

► HOME

B. Kampmann, C. Hemingway, A. Stephens, R. Davidson, A. Goodsall, S. Anderson, M. Nicol, E. Scholvinck, D. Relman, S. Waddell, P. Langford, B. Sheehan, L. Semple, K. A. Wilkinson, R. J. Wilkinson, S. Ress, M. Hibberd, and M. Levin

Acquired predisposition to mycobacterial disease due to autoantibodies to IFN- γ

J. Clin. Invest., September 1, 2005; 115(9): 2480 - 2488.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

► HOME

H. Ji and W. H. Wong

TileMap: create chromosomal map of tiling array hybridizations

Bioinformatics, September 15, 2005; 21(18): 3629 - 3636.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Cell Science

► HOME

N. Bertani, P. Malatesta, G. Volpi, P. Sonogo, and R. Perris

Neurogenic potential of human mesenchymal stem cells revisited: analysis by immunostaining, time-lapse video and microarray

J. Cell Sci., September 1, 2005; 118(17): 3925 - 3936.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood


[HOME](#)

K. R. N. Baumforth, J. R. Flavell, G. M. Reynolds, G. Davies, T. R. Pettit, W. Wei, S. Morgan, T. Stankovic, Y. Kishi, H. Arai, M. Nowakova, G. Pratt, J. Aoki, M. J. O. Wakelam, L. S. Young, and P. G. Murray

Induction of autotaxin by the Epstein-Barr virus promotes the growth and survival of Hodgkin lymphoma cells

Blood, September 15, 2005; 106(6): 2138 - 2146.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

K.-Y. Jen and V. G. Cheung

Identification of Novel p53 Target Genes in Ionizing Radiation Response

Cancer Res., September 1, 2005; 65(17): 7666 - 7673.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

[HOME](#)

S. D. Barr, J. Leipzig, P. Shinn, J. R. Ecker, and F. D. Bushman

Integration Targeting by Avian Sarcoma-Leukosis Virus and Human Immunodeficiency Virus in the Chicken Genome

J. Virol., September 15, 2005; 79(18): 12035 - 12044.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Leukocyte Biology

[HOME](#)

D. McIlroy, S. Tanguy-Royer, N. Le Meur, I. Guisle, P.-J. Royer, J. Leger, K. Meflah, and M. Gregoire

Profiling dendritic cell maturation with dedicated microarrays

J. Leukoc. Biol., September 1, 2005; 78(3): 794 - 803.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology

[HOME](#)

G. Colombo, S. Gatti, F. Turcatti, A. Sordi, L. R. Fassati, F. Bonino, J. M. Lipton, and A. Catania

Gene Expression Profiling Reveals Multiple Protective Influences of the Peptide {alpha}-Melanocyte-Stimulating Hormone in Experimental Heart Transplantation

J. Immunol., September 1, 2005; 175(5): 3391 - 3401.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology

[HOME](#)

W. A. Altemeier, G. Matute-Bello, S. A. Gharib, R. W. Glenny, T. R. Martin, and W. C. Liles

Modulation of Lipopolysaccharide-Induced Gene Transcription and Promotion of Lung Injury by Mechanical Ventilation

J. Immunol., September 1, 2005; 175(5): 3369 - 3376.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

R. P. Morris, L. Nguyen, J. Gatfield, K. Visconti, K. Nguyen, D. Schnappinger, S. Ehrt, Y. Liu, L. Heifets, J. Pieters, G. Schoolnik, and C. J. Thompson

Ancestral antibiotic resistance in Mycobacterium tuberculosis

PNAS, August 23, 2005; 102(34): 12200 - 12205.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[HOME](#)

M. D. Hofer, T. J. Browne, L. He, R. I. Skotheim, R. A. Lothe, and M. A. Rubin

Identification of Two Molecular Groups of Seminomas by Using Expression and Tissue Microarrays

Clin. Cancer Res., August 15, 2005; 11(16): 5722 - 5729.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research


[HOME](#)

R. Rouzier, C. M. Perou, W. F. Symmans, N. Ibrahim, M. Cristofanilli, K. Anderson, K. R. Hess, J. Stec, M. Ayers, P. Wagner, P. Morandi, C. Fan, I. Rabiul, J. S. Ross, G. N. Hortobagyi, and L. Pusztai

Breast Cancer Molecular Subtypes Respond Differently to Preoperative Chemotherapy

Clin. Cancer Res., August 15, 2005; 11(16): 5678 - 5685.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Infection and Immunity

[HOME](#)

K. Schaecher, S. Kumar, A. Yadava, M. Vahey, and C. F. Ockenhouse

Genome-Wide Expression Profiling in Malaria Infection Reveals Transcriptional Changes Associated with Lethal and Nonlethal Outcomes

Infect. Immun., September 1, 2005; 73(9): 6091 - 6100.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

[▶ HOME](#)

K. Chan, C. C. Kim, and S. Falkow

Microarray-Based Detection of *Salmonella enterica* Seroovar Typhimurium Transposon Mutants That Cannot Survive in Macrophages and Mice

Infect. Immun., September 1, 2005; 73(9): 5438 - 5449.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



ANESTHESIA & ANALGESIA

[▶ HOME](#)

D. A. Lowes, H. F. Galley, P. R. Lowe, B. A. Rikke, T. E. Johnson, and N. R. Webster

A Microarray Analysis of Potential Genes Underlying the Neurosensitivity of Mice to Propofol

Anesth. Analg., September 1, 2005; 101(3): 697 - 704.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Development

[▶ HOME](#)

S. N. Sansom, J. M. Hebert, U. Thammongkol, J. Smith, G. Nisbet, M. A. Surani, S. K. McConnell, and F. J. Livesey

Genomic characterisation of a Fgf-regulated gradient-based neocortical protomap

Development, September 1, 2005; 132(17): 3947 - 3961.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Cancer Research

[▶ HOME](#)

J. F. Torres-Roca, S. Eschrich, H. Zhao, G. Bloom, J. Sung, S. McCarthy, A. B. Cantor, A. Scuto, C. Li, S. Zhang, R. Jove, and T. Yeatman

Prediction of Radiation Sensitivity Using a Gene Expression Classifier

Cancer Res., August 15, 2005; 65(16): 7169 - 7176.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am. J. Physiol. Lung, Cellular, and Molecular Physiology


[▶ HOME](#)

S.-F. Ma, D. N. Grigoryev, A. D. Taylor, S. Nonas, S. Sammani, S. Q. Ye, and J. G. N. Garcia

Bioinformatic identification of novel early stress response genes in rodent models of lung injury

Am J Physiol Lung Cell Mol Physiol, September 1, 2005; 289(3): L468 - L477.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

S. H. Ahmed, R. Lutjens, L. D. van der Stap, D. Lekic, V. Romano-Spica, M. Morales, G. F. Koob, V. Repunte-Canonigo, and P. P. Sanna

Gene expression evidence for remodeling of lateral hypothalamic circuitry in cocaine addiction

PNAS, August 9, 2005; 102(32): 11533 - 11538.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[▶ HOME](#)

S. Misra, M. K. Tripathi, and G. Chaudhuri

Down-regulation of 7SL RNA Expression and Impairment of Vesicular Protein Transport Pathways by Leishmania Infection of Macrophages

J. Biol. Chem., August 12, 2005; 280(32): 29364 - 29373.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY

[▶ HOME](#)

A. Vacca, C. Scavelli, V. Montefusco, G. Di Pietro, A. Neri, M. Mattioli, S. Biciato, B. Nico, D. Ribatti, F. Dammacco, and P. Corradini

Thalidomide Downregulates Angiogenic Genes in Bone Marrow Endothelial Cells of Patients With Active Multiple Myeloma

J. Clin. Oncol., August 10, 2005; 23(23): 5334 - 5346.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

H. Liu, J. Li, and L. Wong

Use of extreme patient samples for outcome prediction from gene expression data

Bioinformatics, August 15, 2005; 21(16): 3377 - 3384.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Development


[▶ HOME](#)

A. Klebes, A. Sustar, K. Kechris, H. Li, G. Schubiger, and T. B. Kornberg

Regulation of cellular plasticity in Drosophila imaginal disc cells by the Polycomb group, trithorax group and lama genes

Development, August 15, 2005; 132(16): 3753 - 3765.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENETICS

[▶ HOME](#)

D. Li, P. Bobrowicz, H. H. Wilkinson, and D. J. Ebbole
A Mitogen-Activated Protein Kinase Pathway Essential for Mating and Contributing to Vegetative Growth in *Neurospora crassa*

Genetics, July 1, 2005; 170(3): 1091 - 1104.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation

[▶ HOME](#)

N. Gaborit, M. Steenman, G. Lamirault, N. Le Meur, S. Le Bouter, G. Lande, J. Leger, F. Charpentier, T. Christ, D. Dobrev, D. Escande, S. Nattel, and S. Demolombe
Human Atrial Ion Channel and Transporter Subunit Gene-Expression Remodeling Associated With Valvular Heart Disease and Atrial Fibrillation

Circulation, July 26, 2005; 112(4): 471 - 481.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science

[▶ HOME](#)

V. Vittal, A. Rose, K. E. Gregory, M. J. Kelley, and T. S. Acott
Changes in Gene Expression by Trabecular Meshwork Cells in Response to Mechanical Stretching

Invest. Ophthalmol. Vis. Sci., August 1, 2005; 46(8): 2857 - 2868.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science

[▶ HOME](#)

W. Song, J. L. Barth, Y. Yu, K. Lu, A. Dashti, Y. Huang, C. K. Gittinger, W. S. Argraves, and T. J. Lyons
Effects of Oxidized and Glycated LDL on Gene Expression in Human Retinal Capillary Pericytes

Invest. Ophthalmol. Vis. Sci., August 1, 2005; 46(8): 2974 - 2982.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



diabetes


[▶ HOME](#)

R. Canello, C. Henegar, N. Viguerie, S. Taleb, C. Poitou, C. Rouault, M. Coupaye, V. Pelloux, D. Hugol, J.-L. Bouillot, A. Bouloumie, G. Barbatelli, S. Cinti, P.-A. Svensson, G. S. Barsh, J.-D. Zucker, A. Basdevant, D. Langin, and K. Clement
Reduction of Macrophage Infiltration and

Chemoattractant Gene Expression Changes in White Adipose Tissue of Morbidly Obese Subjects After Surgery-Induced Weight Loss

Diabetes, August 1, 2005; 54(8): 2277 - 2286.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of General Virology


[▶ HOME](#)

R. J. O. Dowling and D. Bienzle

Gene-expression changes induced by Feline immunodeficiency virus infection differ in epithelial cells and lymphocytes

J. Gen. Virol., August 1, 2005; 86(8): 2239 - 2248.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[▶ HOME](#)

M. Alcalay, E. Tiacci, R. Bergomas, B. Bigerna, E. Venturini, S. P. Minardi, N. Meani, D. Diverio, L. Bernard, L. Tizzoni, S. Volorio, L. Luzi, E. Colombo, F. Lo Coco, C. Mecucci, B. Falini, P. G. Pelicci, and for the Gruppo Italiano Malattie Ematologiche Mali

Acute myeloid leukemia bearing cytoplasmic nucleophosmin (NPMc+ AML) shows a distinct gene expression profile characterized by up-regulation of genes involved in stem-cell maintenance

Blood, August 1, 2005; 106(3): 899 - 902.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



HUMAN REPRODUCTION

[▶ HOME](#)

S. Mirkin, M. Arslan, D. Churikov, A. Corica, J.I. Diaz, S. Williams, S. Bocca, and S. Oehninger

In search of candidate genes critically expressed in the human endometrium during the window of implantation

Hum. Reprod., August 1, 2005; 20(8): 2104 - 2117.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics


[▶ HOME](#)

T. Y. Kostrominova, D. E. Dow, R. G. Dennis, R. A. Miller, and J. A. Faulkner

Comparison of gene expression of 2-mo denervated, 2-mo stimulated-denervated, and control rat skeletal muscles

Physiol Genomics, July 14, 2005; 22(2): 227 - 243.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

R. Tabibiazar, R. A. Wagner, E. A. Ashley, J. Y. King, R. Ferrara, J. M. Spin, D. A. Sanan, B. Narasimhan, R. Tibshirani, P. S. Tsao, B. Efron, and T. Quertermous
Signature patterns of gene expression in mouse atherosclerosis and their correlation to human coronary disease

Physiol Genomics, July 14, 2005; 22(2): 213 - 226.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[HOME](#)

J. Hu, F. Zou, and F. A. Wright
Practical FDR-based sample size calculations in microarray experiments

Bioinformatics, August 1, 2005; 21(15): 3264 - 3272.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J. E. Ippolito, J. Xu, S. Jain, K. Moulder, S. Mennerick, J. R. Crowley, R. R. Townsend, and J. I. Gordon
An integrated functional genomics and metabolomics approach for defining poor prognosis in human neuroendocrine cancers

PNAS, July 12, 2005; 102(28): 9901 - 9906.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Applied Physiology

[HOME](#)

M. Fluck, C. Dapp, S. Schmutz, E. Wit, and H. Hoppeler
Transcriptional profiling of tissue plasticity: role of shifts in gene expression and technical limitations

J Appl Physiol, August 1, 2005; 99(2): 397 - 413.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



British Journal of Anaesthesia

[HOME](#)

E. Lucchinetti, R. da Silva, T. Pasch, M. C. Schaub, and M. Zaugg

Anaesthetic preconditioning but not postconditioning prevents early activation of the deleterious cardiac remodelling programme: evidence of opposing genomic responses in cardioprotection by pre- and postconditioning

Br. J. Anaesth., August 1, 2005; 95(2): 140 - 152.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

G. Tonon, K.-K. Wong, G. Maulik, C. Brennan, B. Feng, Y. Zhang, D. B. Khatry, A. Protopopov, M. J. You, A. J. Aguirre, E. S. Martin, Z. Yang, H. Ji, L. Chin, and R. A. DePinho
High-resolution genomic profiles of human lung cancer
PNAS, July 5, 2005; 102(27): 9625 - 9630.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[▶ HOME](#)

E. Pan, M. Pellarin, E. Holmes, I. Smirnov, A. Misra, C. G. Eberhart, P. C. Burger, J. A. Biegel, and B. G. Feuerstein
Isochromosome 17q Is a Negative Prognostic Factor in Poor-Risk Childhood Medulloblastoma Patients
Clin. Cancer Res., July 1, 2005; 11(13): 4733 - 4740.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

Y. Lu, P.-Y. Liu, P. Xiao, and H.-W. Deng
Hotelling's T2 multivariate profiling for detecting differential expression in microarrays
Bioinformatics, July 15, 2005; 21(14): 3105 - 3113.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Reproduction

[▶ HOME](#)

C. A White and L. A Salamonsen
A guide to issues in microarray analysis: application to endometrial biology
Reproduction, July 1, 2005; 130(1): 1 - 13.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

X. J. Yang, M.-H. Tan, H. L. Kim, J. A. Ditlev, M. W. Betten, C. E. Png, E. J. Kort, K. Futami, K. A. Furge, M. Takahashi, H.-o. Kanayama, P. H. Tan, B. S. Teh, C. Luan, K. Wang, M. Pins, M. Tretiakova, J. Anema, R. Kahnoski, T. Nicol, W. Stadler, N. G. Vogelzang, R. Amato, D. Seligson, R. Figlin, A. Belldegrun, C. G. Rogers, and B. T. Teh
A Molecular Classification of Papillary Renal Cell Carcinoma
Cancer Res., July 1, 2005; 65(13): 5628 - 5637.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

R. I. Skotheim, G. E. Lind, O. Monni, J. M. Nesland, V. M. Abeler, S. D. Fossa, N. Duale, G. Brunborg, O. Kallioniemi, P. W. Andrews, and R. A. Lothe

Differentiation of Human Embryonal Carcinomas In vitro and In vivo Reveals Expression Profiles Relevant to Normal Development

Cancer Res., July 1, 2005; 65(13): 5588 - 5598.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[HOME](#)

C. Romualdi, N. Vitulo, M. D. Favero, and G. Lanfranchi

MIDAW: a web tool for statistical analysis of microarray data

Nucleic Acids Res., July 1, 2005; 33(suppl_2): W644 - W649.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[HOME](#)

A. Hsiao, T. Ideker, J. M. Olefsky, and S. Subramaniam

VAMPIRE microarray suite: a web-based platform for the interpretation of gene expression data

Nucleic Acids Res., July 1, 2005; 33(suppl_2): W627 - W632.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

M. S. S. Felipe, R. V. Andrade, F. B. M. Arraes, A. M. Nicola, A. Q. Maranhao, F. A. G. Torres, I. Silva-Pereira, M. J. Pocas-Fonseca, E. G. Campos, L. M. P. Moraes, P. A. Andrade, A. H. F. P. Tavares, S. S. Silva, C. M. Kyaw, D. P. Souza, P. Network, M. Pereira, R. S. A. Jesuino, E. V. Andrade, J. A. Parente, G. S. Oliveira, M. S. Barbosa, N. F. Martins, A. L. Fachin, R. S. Cardoso, G. A. S. Passos, N. F. Almeida, M. E. M. T. Walter, C. M. A. Soares, M. J. A. Carvalho, and M. M. Brigido

Transcriptional Profiles of the Human Pathogenic Fungus *Paracoccidioides brasiliensis* in Mycelium and Yeast Cells

J. Biol. Chem., July 1, 2005; 280(26): 24706 - 24714.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

I. Kalajzic, A. Staal, W.-P. Yang, Y. Wu, S. E. Johnson, J. H. M. Feyen, W. Krueger, P. Maye, F. Yu, Y. Zhao, L. Kuo, R. R. Gupta, L. E. K. Achenie, H.-W. Wang, D.-G. Shin, and D. W. Rowe

Expression Profile of Osteoblast Lineage at Defined Stages of Differentiation

J. Biol. Chem., July 1, 2005; 280(26): 24618 - 24626.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[HOME](#)

J. Qian, N. Esumi, Y. Chen, Q. Wang, I. Chowers, and D. J. Zack

Identification of regulatory targets of tissue-specific transcription factors: application to retina-specific gene regulation

Nucleic Acids Res., June 20, 2005; 33(11): 3479 - 3491.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

K.-H. Pan, C.-J. Lih, and S. N. Cohen

Effects of threshold choice on biological conclusions reached during analysis of gene expression by DNA microarrays

PNAS, June 21, 2005; 102(25): 8961 - 8965.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of Pathology

[HOME](#)

B. S. Knudsen, J. M. Lucas, L. Fazli, S. Hawley, S. Falcon, I. M. Coleman, D. B. Martin, C. Xu, L. D. True, M. E. Gleave, P. S. Nelson, and G. E. Ayala

Regulation of Hepatocyte Activator Inhibitor-1 Expression by Androgen and Oncogenic Transformation in the Prostate

Am. J. Pathol., July 1, 2005; 167(1): 255 - 266.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Carcinogenesis

[HOME](#)

S. Bloethner, B. Chen, K. Hemminki, J. Muller-Berghaus, S. Ugurel, D. Schadendorf, and R. Kumar

Effect of common B-RAF and N-RAS mutations on global gene expression in melanoma cell lines

Carcinogenesis, July 1, 2005; 26(7): 1224 - 1232.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

► HOME

Y. Pawitan, S. Michiels, S. Koscielny, A. Gusnanto, and A. Ploner

False discovery rate, sensitivity and sample size for microarray studies

Bioinformatics, July 1, 2005; 21(13): 3017 - 3024.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics


► HOME

S. A. Gharib, D. L. Luchtel, D. K. Madtes, and R. W. Glenny

Global gene annotation analysis and transcriptional profiling identify key biological modules in hypoxic pulmonary hypertension

Physiol Genomics, June 16, 2005; 22(1): 14 - 23.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

► HOME

R. Datta, M. L. deSchoolmeester, C. Hedeler, N. W. Paton, A. M. Brass, and K. J. Else

Identification of Novel Genes in Intestinal Tissue That Are Regulated after Infection with an Intestinal Nematode Parasite

Infect. Immun., July 1, 2005; 73(7): 4025 - 4033.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Plant Physiology

► HOME

K. Ramonell, M. Berrocal-Lobo, S. Koh, J. Wan, H. Edwards, G. Stacey, and S. Somerville

Loss-of-Function Mutations in Chitin Responsive Genes Show Increased Susceptibility to the Powdery Mildew Pathogen Erysiphe cichoracearum

Plant Physiology, June 1, 2005; 138(2): 1027 - 1036.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

► HOME

G. J. Gordon, G. N. Rockwell, P. A. Godfrey, R. V. Jensen, J. N. Glickman, B. Y. Yeap, W. G. Richards, D. J. Sugarbaker, and R. Bueno

Validation of Genomics-Based Prognostic Tests in Malignant Pleural Mesothelioma

Clin. Cancer Res., June 15, 2005; 11(12): 4406 - 4414.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

K. Camphausen, B. Purow, M. Sproull, T. Scott, T. Ozawa, D. F. Deen, and P. J. Tofilon

From The Cover: Influence of in vivo growth on human glioma cell line gene expression: Convergent profiles under orthotopic conditions

PNAS, June 7, 2005; 102(23): 8287 - 8292.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



TOXICOLOGICAL SCIENCES

► HOME

G. A. Boorman, P. E. Blackshear, J. S. Parker, E. K. Lobenhofer, D. E. Malarkey, M. K. Vallant, D. K. Gerken, and R. D. Irwin

Hepatic Gene Expression Changes throughout the Day in the Fischer Rat: Implications for Toxicogenomic Experiments

Toxicol. Sci., July 1, 2005; 86(1): 185 - 193.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of CLINICAL NUTRITION

► HOME

I. Dahlman, K. Linder, E. Arvidsson Nordstrom, I. Andersson, J. Liden, C. Verdich, T. I. Sorensen, P. Arner, and Nugenob

Changes in adipose tissue gene expression with energy-restricted diets in obese women

Am. J. Clinical Nutrition, June 1, 2005; 81(6): 1275 - 1285.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

► HOME

S. Garcia-Sanchez, A. L. Mavor, C. L. Russell, S. Argimon, P. Dennison, B. Enjalbert, and A. J.P. Brown

Global Roles of Ssn6 in Tup1- and Nrg1-dependent Gene Regulation in the Fungal Pathogen, Candida albicans

Mol. Biol. Cell, June 1, 2005; 16(6): 2913 - 2925.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

► HOME

D. Kadosh and A. D. Johnson

Induction of the Candida albicans Filamentous Growth Program by Relief of Transcriptional Repression: A Genome-wide Analysis

Mol. Biol. Cell, June 1, 2005; 16(6): 2903 - 2912.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

[HOME](#)

S. R. Green and A. D. Johnson

Genome-wide Analysis of the Functions of a Conserved Surface on the Corepressor Tup1

Mol. Biol. Cell, June 1, 2005; 16(6): 2605 - 2613.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology

[HOME](#)

Z. Vuralhan, M. A. H. Luttik, S. L. Tai, V. M. Boer, M. A. Morais, D. Schipper, M. J. H. Almering, P. Kotter, J. R. Dickinson, J.-M. Daran, and J. T. Pronk

Physiological Characterization of the ARO10-Dependent, Broad-Substrate-Specificity 2-Oxo Acid Decarboxylase Activity of *Saccharomyces cerevisiae*

Appl. Envir. Microbiol., June 1, 2005; 71(6): 3276 - 3284.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[HOME](#)

K. Ou, C. Ong, S. Y. Koh, F. Rodrigues, S. H. Sim, D. Wong, C. H. Ooi, K. C. Ng, H. Jikuya, C. C. Yau, S. Y. Soon, D. Kesuma, M. A. Lee, and P. Tan

Integrative Genomic, Transcriptional, and Proteomic Diversity in Natural Isolates of the Human Pathogen *Burkholderia pseudomallei*

J. Bacteriol., June 15, 2005; 187(12): 4276 - 4285.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



the FEBS Journal

[HOME](#)

C. C. Strom, M. Aplin, T. Ploug, T. E. H. Christoffersen, J. Langfort, M. Viese, H. Galbo, S. Haunso, and S. P. Sheikh

Expression profiling reveals differences in metabolic gene expression between exercise-induced cardiac effects and maladaptive cardiac hypertrophy

FEBS J., June 1, 2005; 272(11): 2684 - 2695.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

M. Thomas, Z. Yu, N. Dadgar, S. Varambally, J. Yu, A. M. Chinnaiyan, and A. P. Lieberman

The Unfolded Protein Response Modulates Toxicity of the Expanded Glutamine Androgen Receptor

J. Biol. Chem., June 3, 2005; 280(22): 21264 - 21271.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[▶ HOME](#)

E. Blaveri, J. P. Simko, J. E. Korkola, J. L. Brewer, F. Baehner, K. Mehta, S. DeVries, T. Koppie, S. Pejavar, P. Carroll, and F. M. Waldman

Bladder Cancer Outcome and Subtype Classification by Gene Expression

Clin. Cancer Res., June 1, 2005; 11(11): 4044 - 4055.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

G. R. Grant, J. Liu, and C. J. Stoeckert Jr.

A practical false discovery rate approach to identifying patterns of differential expression in microarray data

Bioinformatics, June 1, 2005; 21(11): 2684 - 2690.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

A. L. Tarca and J. E. K. Cooke

A robust neural networks approach for spatial and intensity-dependent normalization of cDNA microarray data

Bioinformatics, June 1, 2005; 21(11): 2674 - 2683.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of PATHOLOGY

[▶ HOME](#)

G. J. Gordon, G. N. Rockwell, R. V. Jensen, J. G. Rheinwald, J. N. Glickman, J. P. Aronson, B. J. Pottorf, M. D. Nitz, W. G. Richards, D. J. Sugarbaker, and R. Bueno

Identification of Novel Candidate Oncogenes and Tumor Suppressors in Malignant Pleural Mesothelioma Using Large-Scale Transcriptional Profiling

Am. J. Pathol., June 1, 2005; 166(6): 1827 - 1840.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[▶ HOME](#)

A. J. Alvi, B. Austen, V. J. Weston, C. Fegan, D. MacCallum, A. Gianella-Borradori, D. P. Lane, M. Hubank, J. E. Powell, W. Wei, A. M. R. Taylor, P. A. H. Moss, and T. Stankovic

A novel CDK inhibitor, CYC202 (R-roscovitine), overcomes the defect in p53-dependent apoptosis in B-

CLL by down-regulation of genes involved in transcription regulation and survival

Blood, June 1, 2005; 105(11): 4484 - 4491.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY

[▶ HOME](#)

S. Eschrich, I. Yang, G. Bloom, K. Y. Kwong, D. Boulware, A. Cantor, D. Coppola, M. Kruhoffer, L. Aaltonen, T. F. Orntoft, J. Quackenbush, and T. J. Yeatman

Molecular Staging for Survival Prediction of Colorectal Cancer Patients

J. Clin. Oncol., May 20, 2005; 23(15): 3526 - 3535.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity


[▶ HOME](#)

N. K. Viebig, U. Wulbrand, R. Forster, K. T. Andrews, M. Lanzer, and P. A. Knolle

Direct Activation of Human Endothelial Cells by Plasmodium falciparum-Infected Erythrocytes

Infect. Immun., June 1, 2005; 73(6): 3271 - 3277.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Experimental Botany

[▶ HOME](#)

C. J. Baxter, M. Sabar, W. P. Quick, and L. J. Sweetlove

Comparison of changes in fruit gene expression in tomato introgression lines provides evidence of genome-wide transcriptional changes and reveals links to mapped QTLs and described traits

J. Exp. Bot., June 1, 2005; 56(416): 1591 - 1604.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[▶ HOME](#)

M. Bonham, J. Posakony, I. Coleman, B. Montgomery, J. Simon, and P. S. Nelson

Characterization of Chemical Constituents in Scutellaria baicalensis with Antiandrogenic and Growth-Inhibitory Activities toward Prostate Carcinoma

Clin. Cancer Res., May 15, 2005; 11(10): 3905 - 3914.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

BIOINFORMATICS

[▶ HOME](#)

A. A. Sharov, D. B. Dudekula, and M. S. H. Ko



A web-based tool for principal component and significance analysis of microarray data

Bioinformatics, May 15, 2005; 21(10): 2548 - 2549.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS ▶ HOME

K. Y. Yeung, R. E. Bumgarner, and A. E. Raftery

Bayesian model averaging: development of an improved multi-class, gene selection and classification tool for microarray data

Bioinformatics, May 15, 2005; 21(10): 2394 - 2402.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics ▶ HOME

H. P. J. Buermans, E. M. Redout, A. E. Schiel, R. J. P. Musters, M. Zuidwijk, P. P. Eijk, C. van Hardeveld, S. Kasanmoentalib, F. C. Visser, B. Ylstra, and W. S. Simonides

Microarray analysis reveals pivotal divergent mRNA expression profiles early in the development of either compensated ventricular hypertrophy or heart failure

Physiol Genomics, May 11, 2005; 21(3): 314 - 323.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics ▶ HOME

M. M. Kittleson, K. M. Minhas, R. A. Irizarry, S. Q. Ye, G. Edness, E. Breton, J. V. Conte, G. Tomaselli, J. G. N. Garcia, and J. M. Hare

Gene expression analysis of ischemic and nonischemic cardiomyopathy: shared and distinct genes in the development of heart failure

Physiol Genomics, May 11, 2005; 21(3): 299 - 307.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular and Cellular Biology ▶ HOME

F. Miao and R. Natarajan


Mapping Global Histone Methylation Patterns in the Coding Regions of Human Genes

Mol. Cell. Biol., June 1, 2005; 25(11): 4650 - 4661.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

The Journal of THORACIC AND CARDIOVASCULAR SURGERY ▶ HOME


J. G. Coles, C. Boscarino, M. Takahashi, D. Grant, A. Chang, J. Ritter, X. Dai, C. Du, G. Musso, H. Yamabi, J. Goncalves,



A. S. Kumar, J. Woodgett, H. Lu, and G. Hannigan
Cardioprotective stress response in the human fetal heart

J. Thorac. Cardiovasc. Surg., May 1, 2005; 129(5): 1128 - 1136.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Microbiology [▶ HOME](#)

M. L. A. Jansen, J. A. Diderich, M. Mashego, A. Hassane, J. H. de Winde, P. Daran-Lapujade, and J. T. Pronk

Prolonged selection in aerobic, glucose-limited chemostat cultures of *Saccharomyces cerevisiae* causes a partial loss of glycolytic capacity

Microbiology, May 1, 2005; 151(5): 1657 - 1669.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE JOURNAL OF CLINICAL ENDOCRINOLOGY & METABOLISM [▶ HOME](#)

F. Weber, L. Shen, M. A. Aldred, C. D. Morrison, A. Frilling, M. Saji, F. Schuppert, C. E. Broelsch, M. D. Ringel, and C. Eng

Genetic Classification of Benign and Malignant Thyroid Follicular Neoplasia Based on a Three-Gene Combination

J. Clin. Endocrinol. Metab., May 1, 2005; 90(5): 2512 - 2521.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



EUKARYOTIC CELL [▶ HOME](#)

C. Wang and R. J. St. Leger

Developmental and Transcriptional Responses to Host and Nonhost Cuticles by the Specific Locust Pathogen *Metarhizium anisopliae* var. *acridum*

Eukaryot. Cell, May 1, 2005; 4(5): 937 - 947.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology [▶ HOME](#)

B. A. Methe, J. Webster, K. Nevin, J. Butler, and D. R. Lovley
DNA Microarray Analysis of Nitrogen Fixation and Fe (III) Reduction in *Geobacter sulfurreducens*

Appl. Envir. Microbiol., May 1, 2005; 71(5): 2530 - 2538.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

BIOINFORMATICS [▶ HOME](#)



S. Datta and S. Datta

Empirical Bayes screening of many p-values with applications to microarray studies

Bioinformatics, May 1, 2005; 21(9): 1987 - 1994.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

C. Lottaz and R. Spang

Molecular decomposition of complex clinical phenotypes using biologically structured analysis of microarray data

Bioinformatics, May 1, 2005; 21(9): 1971 - 1978.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

G. K. Smyth, J. Michaud, and H. S. Scott

Use of within-array replicate spots for assessing differential expression in microarray experiments

Bioinformatics, May 1, 2005; 21(9): 2067 - 2075.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[▶ HOME](#)

W. T. Barry, A. B. Nobel, and F. A. Wright

Significance analysis of functional categories in gene expression studies: a structured permutation approach

Bioinformatics, May 1, 2005; 21(9): 1943 - 1949.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS


[▶ HOME](#)

J. Toedling, S. Schmeier, M. Heinig, B. Georgi, and S. Roepcke

MACAT--microarray chromosome analysis tool

Bioinformatics, May 1, 2005; 21(9): 2112 - 2113.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of the American Society of Nephrology


[▶ HOME](#)

H. Reich, D. Tritchler, A. M. Herzenberg, Z. Kassiri, X. Zhou, W. Gao, and J. W. Scholey

Albumin Activates ERK Via EGF Receptor in Human Renal Epithelial Cells

J. Am. Soc. Nephrol., May 1, 2005; 16(5): 1266 - 1278.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation Research

► HOME

U. Flögel, T. Laussmann, A. Godecke, N. Abanador, M. Schafers, C. D. Fingas, S. Metzger, B. Levkau, C. Jacoby, and J. Schrader

Lack of Myoglobin Causes a Switch in Cardiac Substrate Selection

Circ. Res., April 29, 2005; 96(8): e68 - e75.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE FASEB JOURNAL

► HOME

J. A. Timmons, O. Larsson, E. Jansson, H. Fischer, T. Gustafsson, P. L. Greenhaff, J. Ridden, J. Rachman, M. Peyrard-Janvid, C. Wahlestedt, and C. J. Sundberg

Human muscle gene expression responses to endurance training provide a novel perspective on Duchenne muscular dystrophy

FASEB J, May 1, 2005; 19(7): 750 - 760.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JCB Online

► HOME

M. Kishi, T. T. Kummer, S. J. Eglén, and J. R. Sanes

LL5{beta}: a regulator of postsynaptic differentiation identified in a screen for synaptically enriched transcripts at the neuromuscular junction

J. Cell Biol., April 25, 2005; 169(2): 355 - 366.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Antimicrobial Agents and Chemotherapy

► HOME

Global Transcriptional Response of *Bacillus subtilis* to Treatment with Subinhibitory Concentrations of Antibiotics That Inhibit Protein Synthesis

Antimicrob. Agents Chemother., May 1, 2005; 49(5): 1915 - 1926.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



the Journal of Molecular Diagnostics

► HOME

A. N. Schuetz, Q. Yin-Goen, M. B. Amin, C. S. Moreno, C. Cohen, C. D. Hornsby, W. L. Yang, J. A. Petros, M. M. Issa, J. G. Pattaras, K. Ogan, F. F. Marshall, and A. N. Young

Molecular Classification of Renal Tumors by Gene Expression Profiling

Mol. Diagn., May 1, 2005; 7(2): 206 - 218.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research


► HOME

D. Magda, P. Lecane, R. A. Miller, C. Lepp, D. Miles, M. Mesfin, J. E. Biaglow, V. V. Ho, D. Chawannakul, S. Nagpal, M. W. Karaman, and J. G. Hacia

Motexafin Gadolinium Disrupts Zinc Metabolism in Human Cancer Cell Lines

Cancer Res., May 1, 2005; 65(9): 3837 - 3845.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



STEM CELLS

► HOME

C. M. Ferrell, S. T. Dorsam, H. Ohta, R. K. Humphries, M. K. Derynck, C. Haqq, C. Largman, and H. J. Lawrence

Activation of Stem-Cell Specific Genes by HOXA9 and HOXA10 Homeodomain Proteins in CD34+ Human Cord Blood Cells

Stem Cells, May 1, 2005; 23(5): 644 - 655.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology

► HOME

C. Huber, C. Thielen, H. Seeger, P. Schwarz, F. Montrasio, M. R. Wilson, E. Heinen, Y.-X. Fu, G. Miele, and A. Aguzzi

Lymphotoxin- β Receptor-Dependent Genes in Lymph Node and Follicular Dendritic Cell Transcriptomes

J. Immunol., May 1, 2005; 174(9): 5526 - 5536.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research


► HOME

A. Misra, M. Pellarin, J. Nigro, I. Smirnov, D. Moore, K. R. Lamborn, D. Pinkel, D. G. Albertson, and B. G. Feuerstein

Array Comparative Genomic Hybridization Identifies Genetic Subgroups in Grade 4 Human Astrocytoma

Clin. Cancer Res., April 15, 2005; 11(8): 2907 - 2918.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

► HOME

J. E. Eckel-Passow, A. Hoering, T. M. Therneau, and I. Ghobrial

Experimental Design and Analysis of Antibody Microarrays: Applying Methods from cDNA Arrays

Cancer Res., April 15, 2005; 65(8): 2985 - 2989.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

► HOME

E. T. Kimchi, M. C. Posner, J. O. Park, T. E. Darga, M. Kocherginsky, T. Karrison, J. Hart, K. D. Smith, J. J. Mezhir, R. R. Weichselbaum, and N. N. Khodarev

Progression of Barrett's Metaplasia to Adenocarcinoma Is Associated with the Suppression of the Transcriptional Programs of Epidermal Differentiation

Cancer Res., April 15, 2005; 65(8): 3146 - 3154.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

► HOME

Z. Chen and L. Liu

RealSpot: software validating results from DNA microarray data analysis with spot images

Physiol Genomics, April 14, 2005; 21(2): 284 - 291.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

► HOME

B. Groschel and F. Bushman

Cell Cycle Arrest in G2/M Promotes Early Steps of Infection by Human Immunodeficiency Virus

J. Virol., May 1, 2005; 79(9): 5695 - 5704.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

► HOME

B. Wu

Differential gene expression detection using penalized linear regression models: the improved SAM statistics

Bioinformatics, April 15, 2005; 21(8): 1565 - 1571.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

► HOME

V. Misra, N. Rapin, O. Akhova, M. Bainbridge, and P. Korchinski

Zhangfei Is a Potent and Specific Inhibitor of the Host Cell Factor-binding Transcription Factor Luman


J. Biol. Chem., April 15, 2005; 280(15): 15257 - 15266.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

The Journal of Immunology

► HOME

B. P.-L. Lee, E. Mansfield, S.-C. Hsieh, T. Hernandez-



Boussard, W. Chen, C. W. Thomson, M. S. Ford, S. E. Bosinger, S. Der, Z.-x. Zhang, M. Zhang, D. J. Kelvin, M. M. Sarwal, and L. Zhang

Expression Profiling of Murine Double-Negative Regulatory T Cells Suggest Mechanisms for Prolonged Cardiac Allograft Survival

J. Immunol., April 15, 2005; 174(8): 4535 - 4544.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[▶ HOME](#)

M. A. Fisher, D. Grimm, A. K. Henion, A. F. Elias, P. E. Stewart, P. A. Rosa, and F. C. Gherardini

From the Cover: *Borrelia burgdorferi* σ^{54} is required for mammalian infection and vector transmission but not for tick colonization

PNAS, April 5, 2005; 102(14): 5162 - 5167.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation Research

[▶ HOME](#)

R. L. Winslow and Z. Gao

Candidate Gene Discovery in Cardiovascular Disease

Circ. Res., April 1, 2005; 96(6): 605 - 606.

[\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

J. P. Cobb, M. N. Mindrinos, C. Miller-Graziano, S. E. Calvano, H. V. Baker, W. Xiao, K. Laudanski, B. H. Brownstein, C. M. Elson, D. L. Hayden, D. N. Herndon, S. F. Lowry, R. V. Maier, D. A. Schoenfeld, L. L. Moldawer, R. W. Davis, R. G. Tompkins, Inflammation and Host Response to Injury Large-Sca, H. V. Baker, P. Bankey, T. Billiar, B. H. Brownstein, S. E. Calvano, D. Camp, I. Chaudry, J. P. Cobb, R. W. Davis, C. M. Elson, B. Freeman, R. Gamelli, N. Gibran, B. Harbrecht, D. L. Hayden, W. Heagy, D. Heimbach, D. N. Herndon, J. Horton, J. Hunt, K. Laudanski, J. Lederer, S. F. Lowry, R. V. Maier, J. Mannick, B. McKinley, C. Miller-Graziano, M. N. Mindrinos, J. Minei, L. L. Moldawer, E. Moore, F. Moore, R. Munford, A. Nathens, G. O'Keefe, G. Purdue, L. Rahme, D. Remick, M. Sailors, D. A. Schoenfeld, M. Shapiro, G. Silver, R. Smith, G. Stephanopoulos, G. Stormo, R. G. Tompkins, M. Toner, S. Warren, M. West, S. Wolfe, W. Xiao, and V. Young

Application of genome-wide expression analysis to human health and disease

PNAS, March 29, 2005; 102(13): 4801 - 4806.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



LEARNING & MEMORY

[HOME](#)

M. A. Wood, M. P. Kaplan, A. Park, E. J. Blanchard, A. M.M. Oliveira, T. L. Lombardi, and T. Abel

Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits in hippocampal synaptic plasticity and memory storage

Learn. Mem., March 1, 2005; 12(2): 111 - 119.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[HOME](#)

Y. H. Yang, Y. Xiao, and M. R. Segal

Identifying differentially expressed genes from microarray experiments via statistic synthesis

Bioinformatics, April 1, 2005; 21(7): 1084 - 1093.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



HUMAN REPRODUCTION

[HOME](#)

A. A. Arslan, L. I. Gold, K. Mittal, T.-C. Suen, I. Belitskaya-Levy, M.-S. Tang, and P. Toniolo

Gene expression studies provide clues to the pathogenesis of uterine leiomyoma: new evidence and a systematic review

Hum. Reprod., April 1, 2005; 20(4): 852 - 863.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

J. Li, M. L. Spletter, and J. A. Johnson

Dissecting tBHQ induced ARE-driven gene expression through long and short oligonucleotide arrays

Physiol Genomics, March 21, 2005; 21(1): 43 - 58.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



RNA

[HOME](#)

K. KIM GUISBERT, K. DUNCAN, H. LI, and C. GUTHRIE

Functional specificity of shuttling hnRNPs revealed by genome-wide analysis of their RNA binding profiles


RNA, April 1, 2005; 11(4): 383 - 393.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Proceedings of the National Academy of Sciences

[HOME](#)

J. C. Mulloy, V. Jankovic, M. Wunderlich, R. Delwel, J. Cammenga, O. Krejci, H. Zhao, P. J. M. Valk, B. Lowenberg,



and S. D. Nimer

AML1-ETO fusion protein up-regulates TRKA mRNA expression in human CD34+ cells, allowing nerve growth factor-induced expansion

PNAS, March 15, 2005; 102(11): 4016 - 4021.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[▶ HOME](#)

D. K. Richardson, S. Kashyap, M. Bajaj, K. Cusi, S. J. Mandarino, J. Finlayson, R. A. DeFronzo, C. P. Jenkinson, and L. J. Mandarino

Lipid Infusion Decreases the Expression of Nuclear Encoded Mitochondrial Genes and Increases the Expression of Extracellular Matrix Genes in Human Skeletal Muscle

J. Biol. Chem., March 18, 2005; 280(11): 10290 - 10297.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Plant Physiology

[▶ HOME](#)

F. Liu, T. VanToai, L. P. Moy, G. Bock, L. D. Linford, and J. Quackenbush

Global Transcription Profiling Reveals Comprehensive Insights into Hypoxic Response in Arabidopsis

Plant Physiology, March 1, 2005; 137(3): 1115 - 1129.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[▶ HOME](#)

K. R. Mehta, K. Nakao, M. B. Zuraek, D. T. Ruan, E. K. Bergsland, A. P. Venook, D. H. Moore, T. A. Tokuyasu, A. N. Jain, R. S. Warren, J. P. Terdiman, and F. M. Waldman

Fractional Genomic Alteration Detected by Array-Based Comparative Genomic Hybridization Independently Predicts Survival after Hepatic Resection for Metastatic Colorectal Cancer

Clin. Cancer Res., March 1, 2005; 11(5): 1791 - 1797.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

J. M. Nigro, A. Misra, L. Zhang, I. Smirnov, H. Colman, C. Griffin, N. Ozburn, M. Chen, E. Pan, D. Koul, W.K. A. Yung, B. G. Feuerstein, and K. D. Aldape

Integrated Array-Comparative Genomic Hybridization and Expression Array Profiles Identify Clinically Relevant Molecular Subtypes of Glioblastoma

Cancer Res., March 1, 2005; 65(5): 1678 - 1686.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Neuroscience

[HOME](#)

R. T. Kerns, A. Ravindranathan, S. Hassan, M. P. Cage, T. York, J. M. Sikela, R. W. Williams, and M. F. Miles

Ethanol-Responsive Brain Region Expression Networks: Implications for Behavioral Responses to Acute Ethanol in DBA/2J versus C57BL/6J Mice

J. Neurosci., March 2, 2005; 25(9): 2255 - 2266.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[HOME](#)

I. Zwir, D. Shin, A. Kato, K. Nishino, T. Latifi, F. Solomon, J. M. Hare, H. Huang, and E. A. Groisman

Dissecting the PhoP regulatory network of Escherichia coli and Salmonella enterica

PNAS, February 22, 2005; 102(8): 2862 - 2867.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

[HOME](#)

A. Stintzi, D. Marlow, K. Palyada, H. Naikare, R. Panciera, L. Whitworth, and C. Clarke

Use of Genome-Wide Expression Profiling and Mutagenesis To Study the Intestinal Lifestyle of Campylobacter jejuni

Infect. Immun., March 1, 2005; 73(3): 1797 - 1810.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



MOLECULAR ENDOCRINOLOGY

[HOME](#)

E. Rico-Bautista, C. J. Greenhalgh, P. Tollet-Egnell, D. J. Hilton, W. S. Alexander, G. Norstedt, and A. Flores-Morales

Suppressor of Cytokine Signaling-2 Deficiency Induces Molecular and Metabolic Changes that Partially Overlap with Growth Hormone-Dependent Effects

Mol. Endocrinol., March 1, 2005; 19(3): 781 - 793.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

N. Le, R. Nagarajan, J. Y. T. Wang, T. Araki, R. E. Schmidt, and J. Milbrandt

Analysis of congenital hypomyelinating Egr2Lo/Lo nerves identifies Sox2 as an inhibitor of Schwann cell

differentiation and myelination

PNAS, February 15, 2005; 102(7): 2596 - 2601.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



ARCHIVES OF DERMATOLOGY [▶ HOME](#)

S. Nambiar, A. Mirmohammadsadeh, R. Doroudi, A. Gustrau, A. Marini, G. Roeder, T. Ruzicka, and U. R. Hengge
Signaling Networks in Cutaneous Melanoma Metastasis Identified by Complementary DNA Microarrays

Arch Dermatol, February 1, 2005; 141(2): 165 - 173.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics [▶ HOME](#)

C. Moreilhon, D. Gras, C. Hologne, O. Bajolet, F. Cottrez, V. Magnone, M. Merten, H. Groux, E. Puchelle, and P. Barbry
Live Staphylococcus aureus and bacterial soluble factors induce different transcriptional responses in human airway cells

Physiol Genomics, February 10, 2005; 20(3): 244 - 255.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Endocrinology [▶ HOME](#)

X. Luo, L. Ding, J. Xu, R. S. Williams, and N. Chegini
Leiomyoma and Myometrial Gene Expression Profiles and Their Responses to Gonadotropin-Releasing Hormone Analog Therapy

Endocrinology, March 1, 2005; 146(3): 1074 - 1096.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS [▶ HOME](#)

Z. Guan and H. Zhao
A semiparametric approach for marker gene selection based on gene expression data

Bioinformatics, February 15, 2005; 21(4): 529 - 536.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS [▶ HOME](#)

B. S. Kim, I. Kim, S. Lee, S. Kim, S. Y. Rha, and H. C. Chung
Statistical methods of translating microarray data into clinically relevant diagnostic information in colorectal cancer

Bioinformatics, February 15, 2005; 21(4): 517 - 528.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOINFORMATICS

[HOME](#)

P. Delmar, S. 屠. Robin, and J. J. Daudin

VarMixt: efficient variance modelling for the differential analysis of replicated gene expression data

Bioinformatics, February 15, 2005; 21(4): 502 - 508.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

S. Desagher, D. Severac, A. Lipkin, C. Bernis, W. Ritchie, A. Le Digarcher, and L. Journot

Genes Regulated in Neurons Undergoing Transcription-dependent Apoptosis Belong to Signaling Pathways Rather than the Apoptotic Machinery

J. Biol. Chem., February 18, 2005; 280(7): 5693 - 5702.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

R. Sandberg and I. Ernberg

Assessment of tumor characteristic gene expression in cell lines using a tissue similarity index (TSI)

PNAS, February 8, 2005; 102(6): 2052 - 2057.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[HOME](#)

N. Wong, K. Y-Y. Chan, P. F. Macgregor, P. B-S. Lai, J. A. Squire, B. Beheshti, M. Albert, and T. W-T. Leung

Transcriptional Profiling Identifies Gene Expression Changes Associated with IFN- α Tolerance in Hepatitis C-Related Hepatocellular Carcinoma Cells

Clin. Cancer Res., February 1, 2005; 11(3): 1319 - 1326.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research


[HOME](#)

V. N. Kristensen, T. Sorlie, J. Geisler, A. Langerod, N. Yoshimura, R. Karesen, N. Harada, P.E. Lonning, and A.-L. Borresen-Dale

Gene Expression Profiling of Breast Cancer in Relation to Estrogen Receptor Status and Estrogen-Metabolizing Enzymes: Clinical Implications

Clin. Cancer Res., January 15, 2005; 11(2): 878s - 883s.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

C. Kristensen, M. Morant, C. E. Olsen, C. T. Ekstrom, D. W. Galbraith, B. Lindberg Moller, and S. Bak

Metabolic engineering of dhurrin in transgenic Arabidopsis plants with marginal inadvertent effects on the metabolome and transcriptome

PNAS, February 1, 2005; 102(5): 1779 - 1784.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Clinical Investigation

▶ HOME

C. J. Greenhalgh, E. Rico-Bautista, M. Lorentzon, A. L. Thaus, P. O. Morgan, T. A. Willson, P. Zervoudakis, D. Metcalf, I. Street, N. A. Nicola, A. D. Nash, L. J. Fabri, G. Norstedt, C. Ohlsson, A. Flores-Morales, W. S. Alexander, and D. J. Hilton
SOCS2 negatively regulates growth hormone action in vitro and in vivo

J. Clin. Invest., February 1, 2005; 115(2): 397 - 406.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

▶ HOME

H. Sultmann, A. v. Heydebreck, W. Huber, R. Kuner, A. Buness, M. Vogt, B. Gunawan, M. Vingron, L. Fuzesi, and A. Poustka

Gene Expression in Kidney Cancer Is Associated with Cytogenetic Abnormalities, Metastasis Formation, and Patient Survival

Clin. Cancer Res., January 15, 2005; 11(2): 646 - 655.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

▶ HOME

J.-M. Kim, H.-Y. Sohn, S. Y. Yoon, J.-H. Oh, J. O. Yang, J. H. Kim, K. S. Song, S.-M. Rho, H. S. Yoo, Y. S. Kim, J.-G. Kim, and N.-S. Kim

Identification of Gastric Cancer-Related Genes Using a cDNA Microarray Containing Novel Expressed Sequence Tags Expressed in Gastric Cancer Cells

Clin. Cancer Res., January 15, 2005; 11(2): 473 - 482.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Chemistry

▶ HOME


T. C.W. Poon, A. Y. Hui, H. L.Y. Chan, I. L. Ang, S. M. Chow, N. Wong, and J. J.Y. Sung

Prediction of Liver Fibrosis and Cirrhosis in Chronic

Hepatitis B Infection by Serum Proteomic Fingerprinting: A Pilot Study

Clin. Chem., February 1, 2005; 51(2): 328 - 335.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Arteriosclerosis, Thrombosis, and Vascular Biology ▶ HOME

R. Tabibiazar, R. A. Wagner, J. M. Spin, E. A. Ashley, B. Narasimhan, E. M. Rubin, B. Efron, P. S. Tsao, R. Tibshirani, and T. Quertermous

Mouse Strain-Specific Differences in Vascular Wall Gene Expression and Their Relationship to Vascular Disease

Arterioscler. Thromb. Vasc. Biol., February 1, 2005; 25(2): 302 - 308.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell ▶ HOME

K. Hardy, L. Mansfield, A. Mackay, S. Benvenuti, S. Ismail, P. Arora, M. J. O'Hare, and P. S. Jat

Transcriptional Networks and Cellular Senescence in Human Mammary Fibroblasts

Mol. Biol. Cell, February 1, 2005; 16(2): 943 - 953.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology ▶ HOME

A. Dorn, H. Zhao, F. Granberg, M. Hosel, D. Webb, C. Svensson, U. Pettersson, and W. Doerfler

Identification of Specific Cellular Genes Up-Regulated Late in Adenovirus Type 12 Infection

J. Virol., February 15, 2005; 79(4): 2404 - 2412.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics ▶ HOME

S. Xia, J. Yang, Y. Su, J. Qian, E. Ma, and G. G. Haddad

Identification of new targets of Drosophila pre-mRNA adenosine deaminase

Physiol Genomics, January 20, 2005; 20(2): 195 - 202.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology ▶ HOME

P. Gyaneshwar, O. Paliy, J. McAuliffe, D. L. Popham, M. I. Jordan, and S. Kustu

Sulfur and Nitrogen Limitation in Escherichia coli K-12:

Specific Homeostatic Responses

J. Bacteriol., February 1, 2005; 187(3): 1074 - 1090.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology [▶ HOME](#)

G. Astua-Monge, J. Freitas-Astua, G. Bacocina, J. Roncoletta, S. A. Carvalho, and M. A. Machado

Expression Profiling of Virulence and Pathogenicity Genes of *Xanthomonas axonopodis* pv. *citri*

J. Bacteriol., February 1, 2005; 187(3): 1201 - 1205.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research [▶ HOME](#)

T. M. Kim, H. J. Jeong, M. Y. Seo, S. C. Kim, G. Cho, C. H. Park, T. S. Kim, K. H. Park, H. C. Chung, and S. Y. Rha

Determination of Genes Related to Gastrointestinal Tract Origin Cancer Cells Using a cDNA Microarray

Clin. Cancer Res., January 1, 2005; 11(1): 79 - 86.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research [▶ HOME](#)

B. M. Fine, G. J.L. Kaspers, M. Ho, A. H. Loonen, and L. M. Boxer

A Genome-Wide View of the In vitro Response to L-Asparaginase in Acute Lymphoblastic Leukemia

Cancer Res., January 1, 2005; 65(1): 291 - 299.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research [▶ HOME](#)

K.-K. Wong, Y.-M. Chang, Y. T.M. Tsang, L. Perlaky, J. Su, A. Adesina, D. L. Armstrong, M. Bhattacharjee, R. Dauser, S. M. Blaney, M. Chintagumpala, and C. C. Lau

Expression Analysis of Juvenile Pilocytic Astrocytomas by Oligonucleotide Microarray Reveals Two Potential Subgroups

Cancer Res., January 1, 2005; 65(1): 76 - 84.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Nucleic Acids Research [▶ HOME](#)

Y. Tan, L. Shi, W. Tong, and C. Wang

Multi-class cancer classification by total principal component regression (TPCR) using microarray gene expression data

Nucleic Acids Res., January 7, 2005; 33(1): 56 - 65.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[▶ HOME](#)

G. Cario, M. Stanulla, B. M. Fine, O. Teuffel, N. v. Neuhoff, A. Schrauder, T. Flohr, B. W. Schafer, C. R. Bartram, K. Welte, B. Schlegelberger, and M. Schrappe

Distinct gene expression profiles determine molecular treatment response in childhood acute lymphoblastic leukemia

Blood, January 15, 2005; 105(2): 821 - 826.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[▶ HOME](#)

S. L. Tai, V. M. Boer, P. Daran-Lapujade, M. C. Walsh, J. H. de Winde, J.-M. Daran, and J. T. Pronk

Two-dimensional Transcriptome Analysis in Chemostat Cultures: COMBINATORIAL EFFECTS OF OXYGEN AVAILABILITY AND MACRONUTRIENT LIMITATION IN SACCHAROMYCES CEREVISIAE

J. Biol. Chem., January 7, 2005; 280(1): 437 - 447.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[▶ HOME](#)

K. Iwamoto, M. Bundo, and T. Kato

Altered expression of mitochondria-related genes in postmortem brains of patients with bipolar disorder or schizophrenia, as revealed by large-scale DNA microarray analysis

Hum. Mol. Genet., January 15, 2005; 14(2): 241 - 253.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY

[▶ HOME](#)

M. Port, H.U. Schmelz, M. Stockinger, C. Sparwasser, P. Albers, T. Pottek, and M. Abend

Gene Expression Profiling in Seminoma and Nonseminoma

J. Clin. Oncol., January 1, 2005; 23(1): 58 - 69.


[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Journal of Nutrition

[▶ HOME](#)

S. Kim, I. Sohn, Y. S. Lee, and Y. S. Lee

Hepatic Gene Expression Profiles Are Altered by



Genistein Supplementation in Mice with Diet-Induced Obesity

J. Nutr., January 1, 2005; 135(1): 33 - 41.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation

[HOME](#)

P. A. Horwitz, E. J. Tsai, M. E. Putt, J. M. Gilmore, J. J. Lepore, M. S. Parmacek, A. C. Kao, S. S. Desai, L. R. Goldberg, S. C. Brozena, M. L. Jessup, J. A. Epstein, and T. P. Cappola

Detection of Cardiac Allograft Rejection and Response to Immunosuppressive Therapy With Peripheral Blood Gene Expression

Circulation, December 21, 2004; 110(25): 3815 - 3821.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

C. Dapp, S. Schmutz, H. Hoppeler, and M. Fluck
Transcriptional reprogramming and ultrastructure during atrophy and recovery of mouse soleus muscle

Physiol Genomics, December 15, 2004; 20(1): 97 - 107.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

R. M. Adam, S. H. Eaton, C. Estrada, A. Nimgaonkar, S.-C. Shih, L. E. H. Smith, I. S. Kohane, D. Bagli, and M. R. Freeman

Mechanical stretch is a highly selective regulator of gene expression in human bladder smooth muscle cells

Physiol Genomics, December 15, 2004; 20(1): 36 - 44.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

R. da Silva, E. Lucchinetti, T. Pasch, M. C. Schaub, and M. Zaugg

Ischemic but not pharmacological preconditioning elicits a gene expression profile similar to unprotected myocardium

Physiol Genomics, December 15, 2004; 20(1): 117 - 130.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

JBC Online

[HOME](#)

M. H. Malone, Z. Wang, and C. W. Distelhorst



The Glucocorticoid-induced Gene tdag8 Encodes a Pro-apoptotic G Protein-coupled Receptor Whose Activation Promotes Glucocorticoid-induced Apoptosis

J. Biol. Chem., December 17, 2004; 279(51): 52850 - 52859.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology


[HOME](#)

X. Zhu, R. Hart, M. S. Chang, J.-W. Kim, S. Y. Lee, Y. A. Cao, D. Mock, E. Ke, B. Saunders, A. Alexander, J. Grosseohme, K.-M. Lin, Z. Yan, R. Hsueh, J. Lee, R. H. Scheuermann, D. A. Fruman, W. Seaman, S. Subramaniam, P. Sternweis, M. I. Simon, and S. Choi

Analysis of the Major Patterns of B Cell Gene Expression Changes in Response to Short-Term Stimulation with 33 Single Ligands

J. Immunol., December 15, 2004; 173(12): 7141 - 7149.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular and Cellular Biology


[HOME](#)

M. J. Walter, J. S. Park, S. K. M. Lau, X. Li, A. A. Lane, R. Nagarajan, W. D. Shannon, and T. J. Ley

Expression Profiling of Murine Acute Promyelocytic Leukemia Cells Reveals Multiple Model-Dependent Progression Signatures

Mol. Cell. Biol., December 15, 2004; 24(24): 10882 - 10893.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

A. Y. Gracey, E. J. Fraser, W. Li, Y. Fang, R. R. Taylor, J. Rogers, A. Brass, and A. R. Cossins

Coping with cold: An integrative, multitissue analysis of the transcriptome of a poikilothermic vertebrate

PNAS, November 30, 2004; 101(48): 16970 - 16975.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

J. B. Geigl, S. Langer, S. Barwisch, K. Pfleghaar, G. Lederer, and M. R. Speicher

Analysis of Gene Expression Patterns and Chromosomal Changes Associated with Aging

Cancer Res., December 1, 2004; 64(23): 8550 - 8557.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Cancer Research

[HOME](#)

W. Wang, S. Goswami, K. Lapidus, A. L. Wells, J. B. Wyckoff, E. Sahai, R. H. Singer, J. E. Segall, and J. S. Condeelis
Identification and Testing of a Gene Expression Signature of Invasive Carcinoma Cells within Primary Mammary Tumors

Cancer Res., December 1, 2004; 64(23): 8585 - 8594.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Annals of SURGICAL ONCOLOGY

[HOME](#)

T. Kan, Y. Shimada, F. Sato, T. Ito, K. Kondo, G. Watanabe, M. Maeda, S. Yamasaki, S. J. Meltzer, and M. Imamura
Prediction of Lymph Node Metastasis with Use of Artificial Neural Networks Based on Gene Expression Profiles in Esophageal Squamous Cell Carcinoma

Ann. Surg. Oncol., December 1, 2004; 11(12): 1070 - 1078.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood


[HOME](#)

M. E. Ross, R. Mahfouz, M. Onciu, H.-C. Liu, X. Zhou, G. Song, S. A. Shurtleff, S. Pounds, C. Cheng, J. Ma, R. C. Ribeiro, J. E. Rubnitz, K. Girtman, W. K. Williams, S. C. Raimondi, D.-C. Liang, L.-Y. Shih, C.-H. Pui, and J. R. Downing

Gene expression profiling of pediatric acute myelogenous leukemia

Blood, December 1, 2004; 104(12): 3679 - 3687.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

[HOME](#)

S. Jeyaseelan, H. W. Chu, S. K. Young, and G. S. Worthen
Transcriptional Profiling of Lipopolysaccharide-Induced Acute Lung Injury

Infect. Immun., December 1, 2004; 72(12): 7247 - 7256.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

C.C. dos Santos, B. Han, C.F. Andrade, X. Bai, S. Uhlig, R. Hubmayr, M. Tsang, M. Lodyga, S. Keshavjee, A.S. Slutsky, and M. Liu

DNA microarray analysis of gene expression in alveolar epithelial cells in response to TNF{alpha}, LPS, and

cyclic stretch

Physiol Genomics, November 12, 2004; 19(3): 331 - 342.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics


[▶ HOME](#)

J. M. Spin, S. Nallamshetty, R. Tabibiazar, E. A. Ashley, J. Y. King, M. Chen, P. S. Tsao, and T. Quertermous

Transcriptional profiling of in vitro smooth muscle cell differentiation identifies specific patterns of gene and pathway activation

Physiol Genomics, November 12, 2004; 19(3): 292 - 302.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Animal Science

[▶ HOME](#)

A. Reverter, Y. H. Wang, K. A. Byrne, S. H. Tan, G. S. Harper, and S. A. Lehnert

Joint analysis of multiple cDNA microarray studies via multivariate mixed models applied to genetic improvement of beef cattle

J Anim Sci, December 1, 2004; 82(12): 3430 - 3439.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am. J. Physiol. Endocrinol. and Metabolism

[▶ HOME](#)

B. Glenmark, M. Nilsson, H. Gao, J.-A. Gustafsson, K. Dahlman-Wright, and H. Westerblad

Difference in skeletal muscle function in males vs. females: role of estrogen receptor- β

Am J Physiol Endocrinol Metab, December 1, 2004; 287(6): E1125 - E1131.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology

[▶ HOME](#)

Y. Xie, L.-s. Chou, A. Cutler, and B. Weimer

DNA Macroarray Profiling of *Lactococcus lactis* subsp. *lactis* IL1403 Gene Expression during Environmental Stresses


Appl. Envir. Microbiol., November 1, 2004; 70(11): 6738 - 6747.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Arteriosclerosis, Thrombosis, and Vascular Biology

[▶ HOME](#)

F. Morello, T. W.A. de Bruin, J. I. Rotter, R. E. Pratt, C. J.H. van der Kallen, G. A. Hladik, V. J. Dzau, C.-C. Liew, and Y.-D.



I. Chen

Differential Gene Expression of Blood-Derived Cell Lines in Familial Combined Hyperlipidemia

Arterioscler. Thromb. Vasc. Biol., November 1, 2004; 24(11): 2149 - 2154.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE FASEB JOURNAL

[▶ HOME](#)


K. CLEMENT, N. VIGUERIE, C. POITOU, C. CARETTE, V. PELLOUX, C. A. CURAT, A. SICARD, S. ROME, A. BENIS, J.-D. ZUCKER, H. VIDAL, M. LAVILLE, G. S. BARSH, A.

BASDEVANT, V. STICH, R. CANCELLO, and D. LANGIN

Weight loss regulates inflammation-related genes in white adipose tissue of obese subjects

FASEB J, November 1, 2004; 18(14): 1657 - 1669.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Nutrition

[▶ HOME](#)

C. Fu, L. Xi, Y. Wu, R. McCarter, A. Richardson, M. Hickey, and E.-S. Han

Hepatic Genes Altered in Expression by Food Restriction Are Not Influenced by the Low Plasma Glucose Level in Young Male GLUT4 Transgenic Mice

J. Nutr., November 1, 2004; 134(11): 2965 - 2974.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology


[▶ HOME](#)

W. Wu, H. Badrane, S. Arora, H. V. Baker, and S. Jin

MucA-Mediated Coordination of Type III Secretion and Alginate Synthesis in Pseudomonas aeruginosa

J. Bacteriol., November 15, 2004; 186(22): 7575 - 7585.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

S. Goswami, W. Wang, J. B. Wyckoff, and J. S. Condeelis

Breast Cancer Cells Isolated by Chemotaxis from Primary Tumors Show Increased Survival and Resistance to Chemotherapy

Cancer Res., November 1, 2004; 64(21): 7664 - 7667.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Molecular Biology of the Cell

[▶ HOME](#)

F. F. Assaad, J.-L. Qiu, H. Youngs, D. Ehrhardt, L. Zimmerli,




M. Kalde, G. Wanner, S. C. Peck, H. Edwards, K. Ramonell, C. R. Somerville, and H. Thordal-Christensen

The PEN1 Syntaxin Defines a Novel Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of Papillae

Mol. Biol. Cell, November 1, 2004; 15(11): 5118 - 5129.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood [HOME](#)

N. J. Lacayo, S. Meshinchi, P. Kinnunen, R. Yu, Y. Wang, C. M. Stuber, L. Douglas, R. Wahab, D. L. Becton, H. Weinstein, M. N. Chang, C. L. Willman, J. P. Radich, R. Tibshirani, Y. Ravindranath, B. I. Sikic, and G. V. Dahl

Gene expression profiles at diagnosis in de novo childhood AML patients identify FLT3 mutations with good clinical outcomes

Blood, November 1, 2004; 104(9): 2646 - 2654.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Chemistry [HOME](#)

C. I. Dumur, S. Nasim, A. M. Best, K. J. Archer, A. C. Ladd, V. R. Mas, D. S. Wilkinson, C. T. Garrett, and A. Ferreira-Gonzalez

Evaluation of Quality-Control Criteria for Microarray Gene Expression Analysis

Clin. Chem., November 1, 2004; 50(11): 1994 - 2002.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science [Science of Aging Knowledge Environment](#) [HOME](#)

S. Melov and A. Hubbard

Microarrays as a Tool to Investigate the Biology of Aging: A Retrospective and a Look to the Future

Sci. Aging Knowl. Environ., October 20, 2004; 2004(42): re7 - re7.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



OBESITY RESEARCH [HOME](#)

N. C. Jerez-Timaure, F. Kearney, E. B. Simpson, E. J. Eisen, and D. Pomp

Characterization of QTL with Major Effects on Fatness and Growth on Mouse Chromosome 2

Obes. Res., September 1, 2004; 12(9): 1408 - 1420.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research


[HOME](#)

L.-X. Qin, K. F. Kerr, and Contributing Members of the
Toxicogenomics Research

**Empirical evaluation of data transformations and
ranking statistics for microarray analysis**

Nucleic Acids Res., October 12, 2004; 32(18): 5471 - 5479.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of General Virology

[HOME](#)

S. Booth, C. Bowman, R. Baumgartner, G. Sorensen, C.
Robertson, M. Coulthart, C. Phillipson, and R. L. Somorjai

**Identification of central nervous system genes
involved in the host response to the scrapie agent
during preclinical and clinical infection**

J. Gen. Virol., November 1, 2004; 85(11): 3459 - 3471.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

M. J. Scian, K. E. R. Stagliano, M. A. Ellis, S. Hassan, M.
Bowman, M. F. Miles, S. P. Deb, and S. Deb

**Modulation of Gene Expression by Tumor-Derived p53
Mutants**

Cancer Res., October 15, 2004; 64(20): 7447 - 7454.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[HOME](#)

N. Le Meur, G. Lamirault, A. Bihouee, M. Steenman, H.
Bedrine-Ferran, R. Teusan, G. Ramstein, and J. J. Leger

**A dynamic, web-accessible resource to process raw
microarray scan data into consolidated gene
expression values: importance of replication**

Nucleic Acids Res., October 8, 2004; 32(18): 5349 - 5358.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology


[HOME](#)

L. Hertel and E. S. Mocarski

**Global Analysis of Host Cell Gene Expression Late
during Cytomegalovirus Infection Reveals Extensive
Dysregulation of Cell Cycle Gene Expression and
Induction of Pseudomitosis Independent of US28
Function**

J. Virol., November 1, 2004; 78(21): 11988 - 12011.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

S. Ogawa, J. Lozach, K. Jepsen, D. Sawka-Verhelle, V. Perissi, R. Sasik, D. W. Rose, R. S. Johnson, M. G. Rosenfeld, and C. K. Glass

A nuclear receptor corepressor transcriptional checkpoint controlling activator protein 1-dependent gene networks required for macrophage activation

PNAS, October 5, 2004; 101(40): 14461 - 14466.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



EUKARYOTIC CELL

► HOME

Z. Zhang, J. Shrager, M. Jain, C.-W. Chang, O. Vallon, and A. R. Grossman

Insights into the Survival of Chlamydomonas reinhardtii during Sulfur Starvation Based on Microarray Analysis of Gene Expression

Eukaryot. Cell, October 1, 2004; 3(5): 1331 - 1348.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



EUKARYOTIC CELL

► HOME

S. Nicholls, M. Straffon, B. Enjalbert, A. Nantel, S. Macaskill, M. Whiteway, and A. J. P. Brown

Msn2- and Msn4-Like Transcription Factors Play No Obvious Roles in the Stress Responses of the Fungal Pathogen Candida albicans

Eukaryot. Cell, October 1, 2004; 3(5): 1111 - 1123.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Neuroscience

► HOME

M. J. Velardo, C. Burger, P. R. Williams, H. V. Baker, M. C. Lopez, T. H. Mareci, T. E. White, N. Muzyczka, and P. J. Reier

Patterns of Gene Expression Reveal a Temporally Orchestrated Wound Healing Response in the Injured Spinal Cord

J. Neurosci., September 29, 2004; 24(39): 8562 - 8576.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

► HOME

C. S. Moreno, S. Ramachandran, D. G. Ashby, N. Laycock, C. A. Plattner, W. Chen, W. C. Hahn, and D. C. Pallas

Signaling and Transcriptional Changes Critical for Transformation of Human Cells by Simian Virus 40 Small Tumor Antigen or Protein Phosphatase 2A B56

{gamma} Knockdown

Cancer Res., October 1, 2004; 64(19): 6978 - 6988.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

[▶ HOME](#)

D. A. Abate, S. Watanabe, and E. S. Mocarski

Major Human Cytomegalovirus Structural Protein pp65 (ppUL83) Prevents Interferon Response Factor 3 Activation in the Interferon Response

J. Virol., October 15, 2004; 78(20): 10995 - 11006.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

[▶ HOME](#)

W. Xiang, O. Windl, G. Wunsch, M. Dugas, A. Kohlmann, N. Dierkes, I. M. Westner, and H. A. Kretzschmar

Identification of Differentially Expressed Genes in Scrapie-Infected Mouse Brains by Using Global Gene Expression Technology

J. Virol., October 15, 2004; 78(20): 11051 - 11060.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[▶ HOME](#)

Y. Kurokawa, R. Matoba, H. Nagano, M. Sakon, I. Takemasa, S. Nakamori, K. Dono, K. Umeshita, N. Ueno, S. Ishii, K. Kato, and M. Monden

Molecular Prediction of Response to 5-Fluorouracil and Interferon-{alpha} Combination Chemotherapy in Advanced Hepatocellular Carcinoma

Clin. Cancer Res., September 15, 2004; 10(18): 6029 - 6038.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

J. Grimm, A. Mueller, F. Hefti, and A. Rosenthal

Molecular basis for catecholaminergic neuron diversity

PNAS, September 21, 2004; 101(38): 13891 - 13896.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[▶ HOME](#)

D. G. Romero, M. Plonczynski, G. R. Vergara, E. P. Gomez-Sanchez, and C. E. Gomez-Sanchez

Angiotensin II early regulated genes in H295R human adrenocortical cells

Physiol Genomics, September 16, 2004; 19(1): 106 - 116.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Plant Physiology

[▶ HOME](#)

P. Armengaud, R. Breitling, and A. Amtmann

The Potassium-Dependent Transcriptome of Arabidopsis Reveals a Prominent Role of Jasmonic Acid in Nutrient Signaling

Plant Physiology, September 1, 2004; 136(1): 2556 - 2576.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[▶ HOME](#)

K. E. Rieger and G. Chu

Portrait of transcriptional responses to ultraviolet and ionizing radiation in human cells

Nucleic Acids Res., September 8, 2004; 32(16): 4786 - 4803.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

T. H. Qiu, G. V. R. Chandramouli, K. W. Hunter, N. W. Alkharouf, J. E. Green, and E. T. Liu

Global Expression Profiling Identifies Signatures of Tumor Virulence in MMTV-PyMT-Transgenic Mice: Correlation to Human Disease

Cancer Res., September 1, 2004; 64(17): 5973 - 5981.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[▶ HOME](#)

C. L. Yauk, M. L. Berndt, A. Williams, and G. R. Douglas

Comprehensive comparison of six microarray technologies

Nucleic Acids Res., August 27, 2004; 32(15): e124 - e124.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE FASEB JOURNAL

[▶ HOME](#)

S.-T. J. HUANG, K. C. T. VO, D. J. LYELL, G. H. FAESSEN, S. TULAC, R. TIBSHIRANI, A. J. GIACCIA, and L. C. GIUDICE

Developmental response to hypoxia

FASEB J, September 1, 2004; 18(12): 1348 - 1365.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

[HOME](#)

S. R. Green and A. D. Johnson

Promoter-dependent Roles for the Srb10 Cyclin-dependent Kinase and the Hda1 Deacetylase in Tup1-mediated Repression in *Saccharomyces cerevisiae*

Mol. Biol. Cell, September 1, 2004; 15(9): 4191 - 4202.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY

[HOME](#)

M. A. Aldred, Y. Huang, S. Liyanarachchi, N. S. Pellegata, O. Gimm, S. Jhiang, R. V. Davuluri, A. de La Chapelle, and C. Eng

Papillary and Follicular Thyroid Carcinomas Show Distinctly Different Microarray Expression Profiles and Can Be Distinguished by a Minimum of Five Genes

J. Clin. Oncol., September 1, 2004; 22(17): 3531 - 3539.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research

[HOME](#)

K. Yu, C. H. Lee, P. H. Tan, and P. Tan

Conservation of Breast Cancer Molecular Subtypes and Transcriptional Patterns of Tumor Progression Across Distinct Ethnic Populations

Clin. Cancer Res., August 15, 2004; 10(16): 5508 - 5517.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Cancer Research

[HOME](#)

R. Namba, J. E. Maglione, L. J.T. Young, A. D. Borowsky, R. D. Cardiff, C. L. MacLeod, and J. P. Gregg

Molecular Characterization of the Transition to Malignancy in a Genetically Engineered Mouse-Based Model of Ductal Carcinoma In situ

Mol. Cancer Res., August 1, 2004; 2(8): 453 - 463.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science


[HOME](#)

K. Ishibashi, J. Tian, and J. T. Handa

Similarity of mRNA Phenotypes of Morphologically Normal Macular and Peripheral Retinal Pigment Epithelial Cells in Older Human Eyes

Invest. Ophthalmol. Vis. Sci., September 1, 2004; 45(9): 3291 - 3301.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science

[HOME](#)

A. S. Hackam, R. Strom, D. Liu, J. Qian, C. Wang, D. Otteson, T. Gunatilaka, R. H. Farkas, I. Chowers, M. Kageyama, T. Leveillard, J.-A. Sahel, P. A. Campochiaro, G. Parmigiani, and D. J. Zack

Identification of Gene Expression Changes Associated with the Progression of Retinal Degeneration in the rd1 Mouse

Invest. Ophthalmol. Vis. Sci., September 1, 2004; 45(9): 2929 - 2942.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Carcinogenesis

[HOME](#)

L. McArdle, M. McDermott, R. Purcell, D. Grehan, A. O'Meara, F. Breatnach, D. Catchpoole, A. C. Culhane, I. Jeffery, W. M. Gallagher, and R. L. Stallings

Oligonucleotide microarray analysis of gene expression in neuroblastoma displaying loss of chromosome 11q

Carcinogenesis, September 1, 2004; 25(9): 1599 - 1609.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Carcinogenesis

[HOME](#)

P. J. Moos, K. Edes, J. E. Mullally, and F. A. Fitzpatrick

Curcumin impairs tumor suppressor p53 function in colon cancer cells

Carcinogenesis, September 1, 2004; 25(9): 1611 - 1617.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

S. Diederichs, E. Bulk, B. Steffen, P. Ji, L. Tickenbrock, K. Lang, K. S. Zanker, R. Metzger, P. M. Schneider, V. Gerke, M. Thomas, W. E. Berdel, H. Serve, and C. Muller-Tidow

S100 Family Members and Trypsinogens Are Predictors of Distant Metastasis and Survival in Early-Stage Non-Small Cell Lung Cancer

Cancer Res., August 15, 2004; 64(16): 5564 - 5569.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

R. van Doorn, R. Dijkman, M. H. Vermeer, J. J. Out-Luiting, E. M. H. van der Raaij-Helmer, R. Willemze, and C. P. Tensen

Aberrant Expression of the Tyrosine Kinase Receptor EphA4 and the Transcription Factor Twist in Sezary

Syndrome Identified by Gene Expression Analysis

Cancer Res., August 15, 2004; 64(16): 5578 - 5586.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Experimental Botany

[HOME](#)

M. Y. Hirai and K. Saito

Post-genomics approaches for the elucidation of plant adaptive mechanisms to sulphur deficiency

J. Exp. Bot., August 1, 2004; 55(404): 1871 - 1879.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

R. L. McCaffrey, P. Fawcett, M. O'Riordan, K.-D. Lee, E. A. Havell, P. O. Brown, and D. A. Portnoy

From the Cover: A specific gene expression program triggered by Gram-positive bacteria in the cytosol

PNAS, August 3, 2004; 101(31): 11386 - 11391.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



European Respiratory Journal

[HOME](#)

C. Schulz, V. Petrig, K. Wolf, K. Kratzel, M. Kohler, B. Becker, and M. Pfeifer

Upregulation of MCAM in primary bronchial epithelial cells from patients with COPD

Eur. Respir. J., September 1, 2003; 22(3): 450 - 456.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Chemistry

[HOME](#)

T. C.W. Poon, K.C. A. Chan, P.-C. Ng, R. W.K. Chiu, I. L. Ang, Y.-k. Tong, E. K.O. Ng, F. W.T. Cheng, A. M. Li, E. K.L. Hon, T.-F. Fok, and Y.M. D. Lo

Serial Analysis of Plasma Proteomic Signatures in Pediatric Patients with Severe Acute Respiratory Syndrome and Correlation with Viral Load

Clin. Chem., August 1, 2004; 50(8): 1452 - 1455.

[\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

Y. Peng, Q. Kang, Q. Luo, W. Jiang, W. Si, B. A. Liu, H. H. Luu, J. K. Park, X. Li, J. Luo, A. G. Montag, R. C. Haydon, and T.-C. He

Inhibitor of DNA Binding/Differentiation Helix-Loop-Helix Proteins Mediate Bone Morphogenetic Protein-

induced Osteoblast Differentiation of Mesenchymal Stem Cells

J. Biol. Chem., July 30, 2004; 279(31): 32941 - 32949.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Experimental Botany

[HOME](#)

J. Melkonian, L.-X. Yu, and T. L. Setter

Chilling responses of maize (*Zea mays* L.) seedlings: root hydraulic conductance, abscisic acid, and stomatal conductance

J. Exp. Bot., August 1, 2004; 55(403): 1751 - 1760.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Plant Physiology

[HOME](#)

J. A. Lopez-Valenzuela, B. C. Gibbon, D. R. Holding, and B. A. Larkins

Cytoskeletal Proteins Are Coordinately Increased in Maize Genotypes with High Levels of eEF1A

Plant Physiology, July 1, 2004; 135(3): 1784 - 1797.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology

[HOME](#)

J. A. Whitsett, C. J. Bachurski, K. C. Barnes, P. A. Bunn Jr., L. M. Case, D. N. Cook, D. Crooks, M. W. Duncan, L. Dwyer-Nield, R. C. Elston, M. B. Fessler, W. A. Franklin, N. Friedman, J. G. N. Garcia, M. W. Geraci, C. Glasgow, S. W. Glasser, W. D. Hardie, L. M. Henning, G. L. Johnson, K. Kawkitinarong, R. L. Keith, T. R. Korfhagen, G. D. Leikauf, S. B. Liggett, K. C. Malcolm, A. M. Malkinson, T. J. Mariani, S. A. McDowell, D. W. McGraw, M. Medvedovic, J. Moss, L. M. Noguee, S. Nonas, G. Pacheco-Rodriguez, L. J. Palmer, D. G. Peters, D. R. Prows, S. Redline, A. Regev, M. A. Sartor, D. A. Schwartz, E. K. Silverman, W. K. Steagall, R. S. Stearman, A. Taveira-DaSilva, J. W. Tichelaar, C. R. Tomlinson, K. Tsukada, T. E. Weaver, S. E. Wert, S. C. Wesselkamper, G. S. Worthen, Y. Xu, L. Zerbe, Y. Zhang, Y. Zhang, A. M. K. Choi, and N. Kaminski

Functional Genomics of Lung Disease

Am. J. Respir. Cell Mol. Biol., August 1, 2004; 31(2/S1): S1 - S81.

[\[Full Text\]](#) [\[PDF\]](#)

blood

[HOME](#)

W. Wagner, A. Ansorge, U. Wirkner, V. Eckstein, C.




Schwager, J. Blake, K. Miesala, J. Selig, R. Saffrich, W. Ansorge, and A. D. Ho

Molecular evidence for stem cell function of the slow-dividing fraction among human hematopoietic progenitor cells by genome-wide analysis

Blood, August 1, 2004; 104(3): 675 - 686.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Microbiology [▶ HOME](#)

V. Kapatral, J. W. Campbell, S. A. Minnich, N. R. Thomson, P. Matsumura, and B. M. Pruss

Gene array analysis of *Yersinia enterocolitica* FlhD and FlhC: regulation of enzymes affecting synthesis and degradation of carbamoylphosphate

Microbiology, July 1, 2004; 150(7): 2289 - 2300.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics [▶ HOME](#)

J. E. Larkin, B. C. Frank, R. M. Gaspard, I. Duka, H. Gavras, and J. Quackenbush

Cardiac transcriptional response to acute and chronic angiotensin II treatments

Physiol Genomics, July 8, 2004; 18(2): 152 - 166.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Arteriosclerosis, Thrombosis, and Vascular Biology [▶ HOME](#)

E. R. Mulvihill, J. Jaeger, R. Sengupta, W. L. Ruzzo, C. Reimer, S. Lukito, and S. M. Schwartz

Atherosclerotic Plaque Smooth Muscle Cells Have a Distinct Phenotype

Arterioscler. Thromb. Vasc. Biol., July 1, 2004; 24(7): 1283 - 1289.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences [▶ HOME](#)


R. M. Mitra, S. L. Shaw, and S. R. Long

Six nonnodulating plant mutants defective for Nod factor-induced transcriptional changes associated with the legume-rhizobia symbiosis

PNAS, July 6, 2004; 101(27): 10217 - 10222.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Proceedings of the National Academy of Sciences [▶ HOME](#)



M. Y. Hirai, M. Yano, D. B. Goodenowe, S. Kanaya, T. Kimura, M. Awazuhara, M. Arita, T. Fujiwara, and K. Saito
From The Cover: Integration of transcriptomics and metabolomics for understanding of global responses to nutritional stresses in *Arabidopsis thaliana*
PNAS, July 6, 2004; 101(27): 10205 - 10210.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




JBC Online [HOME](#)
D. M. Sadlier, S. B. Connolly, N. E. Kieran, S. Roxburgh, D. P. Brazil, L. Kairaitis, Y. Wang, D. C. H. Harris, P. Doran, and H. R. Brady
Sequential Extracellular Matrix-focused and Baited-global Cluster Analysis of Serial Transcriptomic Profiles Identifies Candidate Modulators of Renal Tubulointerstitial Fibrosis in Murine Adriamycin-induced Nephropathy
J. Biol. Chem., July 9, 2004; 279(28): 29670 - 29680.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences [HOME](#)
D. Manna, A. M. Breier, and N. P. Higgins
Microarray analysis of transposition targets in *Escherichia coli*: The impact of transcription
PNAS, June 29, 2004; 101(26): 9780 - 9785.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science [HOME](#)
I. Chowers, Y. Kim, R. H. Farkas, T. L. Gunatilaka, A. S. Hackam, P. A. Campochiaro, S. C. Finnemann, and D. J. Zack
Changes in Retinal Pigment Epithelial Gene Expression Induced by Rod Outer Segment Uptake
Invest. Ophthalmol. Vis. Sci., July 1, 2004; 45(7): 2098 - 2106.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology [HOME](#)
T. N. Nguyen, A. D. Ejaz, M. A. Brancieri, A. M. Mikula, K. E. Nelson, S. R. Gill, and K. M. Noll
Whole-Genome Expression Profiling of *Thermotoga maritima* in Response to Growth on Sugars in a Chemostat
J. Bacteriol., July 15, 2004; 186(14): 4824 - 4828.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

► HOME

C. C. Kim and S. Falkow

Delineation of Upstream Signaling Events in the Salmonella Pathogenicity Island 2 Transcriptional Activation Pathway

J. Bacteriol., July 15, 2004; 186(14): 4694 - 4704.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

► HOME

S. Volinia, R. Evangelisti, F. Francioso, D. Arcelli, M. Carella, and P. Gasparini

GOAL: automated Gene Ontology analysis of expression profiles

Nucleic Acids Res., July 1, 2004; 32(suppl_2): W492 - W499.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research


► HOME

S. Karanam and C. S. Moreno

CONFAC: automated application of comparative genomic promoter analysis to DNA microarray datasets

Nucleic Acids Res., July 1, 2004; 32(suppl_2): W475 - W484.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



ANNALS of the New York Academy of Sciences


► HOME

B. T. GREER and J. KHAN

Diagnostic Classification of Cancer Using DNA Microarrays and Artificial Intelligence

Ann. N.Y. Acad. Sci., May 1, 2004; 1020(1): 49 - 66.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



ANNALS of the New York Academy of Sciences

► HOME

I. M. KAPETANOVIC, S. ROSENFELD, and G. IZMIRLIAN

Overview of Commonly Used Bioinformatics Methods and Their Applications


Ann. N.Y. Acad. Sci., May 1, 2004; 1020(1): 10 - 21.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Human Molecular Genetics

► HOME

A. T. Dobson, R. Raja, M. J. Abeyta, T. Taylor, S. Shen, C.




Haqq, and R. A. R. Pera

The unique transcriptome through day 3 of human preimplantation development

Hum. Mol. Genet., July 15, 2004; 13(14): 1461 - 1470.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

[HOME](#)

T. L. Nicholson, K. Chiu, and R. S. Stephens

Chlamydia trachomatis Lacks an Adaptive Response to Changes in Carbon Source Availability

Infect. Immun., July 1, 2004; 72(7): 4286 - 4289.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of PATHOLOGY

[HOME](#)

J. Iqbal, W. G. Sanger, D. E. Horsman, A. Rosenwald, D. L. Pickering, B. Dave, S. Dave, L. Xiao, K. Cao, Q. Zhu, S. Sherman, C. P. Hans, D. D. Weisenburger, T. C. Greiner, R. D. Gascoyne, G. Ott, H. K. Muller-Hermelink, J. Delabie, R. M. Braziel, E. S. Jaffe, E. Campo, J. C. Lynch, J. M. Connors, J. M. Vose, J. O. Armitage, T. M. Grogan, L. M. Staudt, and W. C. Chan

BCL2 Translocation Defines a Unique Tumor Subset within the Germinal Center B-Cell-Like Diffuse Large B-Cell Lymphoma

Am. J. Pathol., July 1, 2004; 165(1): 159 - 166.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Genes & Development

[HOME](#)

O. J. Sansom, K. R. Reed, A. J. Hayes, H. Ireland, H. Brinkmann, I. P. Newton, E. Batlle, P. Simon-Assmann, H. Clevers, I. S. Nathke, A. R. Clarke, and D. J. Winton

Loss of Apc in vivo immediately perturbs Wnt signaling, differentiation, and migration

Genes & Dev., June 15, 2004; 18(12): 1385 - 1390.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

M. A. Troester, K. A. Hoadley, T. Sorlie, B.-S. Herbert, A.-L. Borresen-Dale, P. E. Lonning, J. W. Shay, W. K. Kaufmann, and C. M. Perou

Cell-Type-Specific Responses to Chemotherapeutics in Breast Cancer

Cancer Res., June 15, 2004; 64(12): 4218 - 4226.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

B. G. Masayeva, P. Ha, E. Garrett-Mayer, T. Pilkington, R. Mao, J. Pevsner, T. Speed, N. Benoit, C.-S. Moon, D. Sidransky, W. H. Westra, and J. Califano

Gene expression alterations over large chromosomal regions in cancers include multiple genes unrelated to malignant progression

PNAS, June 8, 2004; 101(23): 8715 - 8720.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science

[HOME](#)

D. C. Bergmann, W. Lukowitz, and C. R. Somerville

Stomatal Development and Pattern Controlled by a MAPKK Kinase

Science, June 4, 2004; 304(5676): 1494 - 1497.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH

[HOME](#)

C. Conrad, H. Erfle, P. Warnat, N. Daigle, T. Lorch, J. Ellenberg, R. Pepperkok, and R. Eils

Automatic Identification of Subcellular Phenotypes on Human Cell Arrays

Genome Res., June 1, 2004; 14(6): 1130 - 1136.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

[HOME](#)

H. Zhao, A. Langerod, Y. Ji, K. W. Nowels, J. M. Nesland, R. Tibshirani, I. K. Bukholm, R. Karsen, D. Botstein, A.-L. Borresen-Dale, and S. S. Jeffrey

Different Gene Expression Patterns in Invasive Lobular and Ductal Carcinomas of the Breast

Mol. Biol. Cell, June 1, 2004; 15(6): 2523 - 2536.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

L. Dyrskjot, M. Kruhoffer, T. Thykjaer, N. Marcussen, J. L. Jensen, K. Moller, and T. F. Orntoft

Gene Expression in the Urinary Bladder: A Common Carcinoma in Situ Gene Expression Signature Exists Disregarding Histopathological Classification

Cancer Res., June 1, 2004; 64(11): 4040 - 4048.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research


► HOME

S. Jain, M. A. Watson, M. K. DeBenedetti, Y. Hiraki, J. F. Moley, and J. Milbrandt

Expression Profiles Provide Insights into Early Malignant Potential and Skeletal Abnormalities in Multiple Endocrine Neoplasia Type 2B Syndrome Tumors

Cancer Res., June 1, 2004; 64(11): 3907 - 3913.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

E. Glawischnig, B. G. Hansen, C. E. Olsen, and B. A. Halkier
Camalexin is synthesized from indole-3-acetaldoxime, a key branching point between primary and secondary metabolism in Arabidopsis

PNAS, May 25, 2004; 101(21): 8245 - 8250.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

► HOME

J. L. Hall, S. Grindle, X. Han, D. Fermin, S. Park, Y. Chen, R. J. Bache, A. Mariash, Z. Guan, S. Ormaza, J. Thompson, J. Graziano, S. E. de Sam Lazaro, S. Pan, R. D. Simari, and L. W. Miller

Genomic profiling of the human heart before and after mechanical support with a ventricular assist device reveals alterations in vascular signaling networks

Physiol Genomics, May 19, 2004; 17(3): 283 - 291.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

► HOME

J. Morrison, K. Knoll, M. J. Hessner, and M. Liang
Effect of high glucose on gene expression in mesangial cells: upregulation of the thiol pathway is an adaptational response

Physiol Genomics, May 19, 2004; 17(3): 271 - 282.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

G. N. Landis, D. Abdueva, D. Skvortsov, J. Yang, B. E. Rabin, J. Carrick, S. Tavare, and J. Tower

Similar gene expression patterns characterize aging and oxidative stress in Drosophila melanogaster

PNAS, May 18, 2004; 101(20): 7663 - 7668.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology


[▶ HOME](#)

R. J. Feezor, H. V. Baker, W. Xiao, W. A. Lee, T. S. Huber, M. Mindrinos, R. A. Kim, L. Ruiz-Taylor, L. L. Moldawer, R. W. Davis, and J. M. Seeger

Genomic and Proteomic Determinants of Outcome in Patients Undergoing Thoracoabdominal Aortic Aneurysm Repair

J. Immunol., June 1, 2004; 172(11): 7103 - 7109.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

K.-L. Schaefer, K. Brachwitz, D. H. Wai, Y. Braun, R. Diallo, E. Korsching, M. Eisenacher, R. Voss, F. van Valen, C. Baer, B. Selle, L. Spahn, S.-K. Liao, K. A. W. Lee, P. C. W. Hogendoorn, G. Reifemberger, H. E. Gabbert, and C. Poremba
Expression Profiling of t(12;22) Positive Clear Cell Sarcoma of Soft Tissue Cell Lines Reveals Characteristic Up-Regulation of Potential New Marker Genes Including ERBB3

Cancer Res., May 15, 2004; 64(10): 3395 - 3405.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE PLANT CELL


[▶ HOME](#)

F. Wellmer, J. L. Riechmann, M. Alves-Ferreira, and E. M. Meyerowitz

Genome-Wide Analysis of Spatial Gene Expression in Arabidopsis Flowers

PLANT CELL, May 1, 2004; 16(5): 1314 - 1326.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Microbiology

[▶ HOME](#)

S. Arous, C. Buchrieser, P. Folio, P. Glaser, A. Namane, M. Hebraud, and Y. Hechard

Global analysis of gene expression in an rpoN mutant of Listeria monocytogenes

Microbiology, May 1, 2004; 150(5): 1581 - 1590.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

THE JOURNAL OF CLINICAL ENDOCRINOLOGY & METABOLISM


[▶ HOME](#)

N. Viguerie, K. Clement, P. Barbe, M. Courtine, A. Benis, D.



Larrouy, B. Hanczar, V. Pelloux, C. Poitou, Y. Khalfallah, G. S. Barsh, C. Thalamas, J.-D. Zucker, and D. Langin
In Vivo Epinephrine-Mediated Regulation of Gene Expression in Human Skeletal Muscle
J. Clin. Endocrinol. Metab., May 1, 2004; 89(5): 2000 - 2014.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research [▶ HOME](#)

K. Yu, C. H. Lee, P. H. Tan, G. S. Hong, S. B. Wee, C. Y. Wong, and P. Tan
A Molecular Signature of the Nottingham Prognostic Index in Breast Cancer
Cancer Res., May 1, 2004; 64(9): 2962 - 2968.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research [▶ HOME](#)

M. H. Asyali, M. M. Shoukri, O. Demirkaya, and K. S. A. Khabar
Assessment of reliability of microarray data and estimation of signal thresholds using mixture modeling
Nucleic Acids Res., April 27, 2004; 32(8): 2323 - 2335.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences [▶ HOME](#)

K. E. Rieger, W.-J. Hong, V. G. Tusher, J. Tang, R. Tibshirani, and G. Chu
Toxicity from radiation therapy associated with abnormal transcriptional responses to DNA damage
PNAS, April 27, 2004; 101(17): 6635 - 6640.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Nucleic Acids Research [▶ HOME](#)

G. Wang, L. Chuang, X. Zhang, S. Colton, A. Dombkowski, J. Reiners, A. Diakiw, and X. S. Xu
The initiative role of XPC protein in cisplatin DNA damaging treatment-mediated cell cycle regulation
Nucleic Acids Res., April 23, 2004; 32(7): 2231 - 2240.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

The NEW ENGLAND JOURNAL of MEDICINE [▶ HOME](#)

I. S. Lossos, D. K. Czerwinski, A. A. Alizadeh, M. A. Wechsler, R. Tibshirani, D. Botstein, and R. Levy



Prediction of Survival in Diffuse Large-B-Cell Lymphoma Based on the Expression of Six Genes

N. Engl. J. Med., April 29, 2004; 350(18): 1828 - 1837.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

J. Z. Pan, R. Jornsten, and R. P. Hart

Screening anti-inflammatory compounds in injured spinal cord with microarrays: a comparison of bioinformatics analysis approaches

Physiol Genomics, April 13, 2004; 17(2): 201 - 214.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

J. Tian, K. Ishibashi, and J. T. Handa

The expression of native and cultured RPE grown on different matrices

Physiol Genomics, April 13, 2004; 17(2): 170 - 182.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics


[HOME](#)

L. T. Vaszar, T. Nishimura, J. D. Storey, G. Zhao, D. Qiu, J. L. Faul, R. G. Pearl, and P. N. Kao

Longitudinal transcriptional analysis of developing neointimal vascular occlusion and pulmonary hypertension in rats

Physiol Genomics, April 13, 2004; 17(2): 150 - 156.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE NEW ENGLAND JOURNAL OF MEDICINE

[HOME](#)

P. J.M. Valk, R. G.W. Verhaak, M. A. Beijen, C. A.J. Erpelinck, S. B. v. W. van Doorn-Khosrovani, J. M. Boer, H. B. Beverloo, M. J. Moorhouse, P. J. van der Spek, B. Lowenberg, and R. Delwel

Prognostically Useful Gene-Expression Profiles in Acute Myeloid Leukemia

N. Engl. J. Med., April 15, 2004; 350(16): 1617 - 1628.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE NEW ENGLAND JOURNAL OF MEDICINE

[HOME](#)


L. Bullinger, K. Dohner, E. Bair, S. Frohling, R. F. Schlenk, R. Tibshirani, H. Dohner, and J. R. Pollack

Use of Gene-Expression Profiling to Identify Prognostic

Subclasses in Adult Acute Myeloid Leukemia

N. Engl. J. Med., April 15, 2004; 350(16): 1605 - 1616.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Microbiology

[▶ HOME](#)

A. M. Prouty, I. E. Brodsky, S. Falkow, and J. S. Gunn
Bile-salt-mediated induction of antimicrobial and bile resistance in *Salmonella typhimurium*

Microbiology, April 1, 2004; 150(4): 775 - 783.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



EUKARYOTIC CELL

[▶ HOME](#)

S. Garcia-Sanchez, S. Aubert, I. Iraqui, G. Janbon, J.-M. Ghigo, and C. d'Enfert

***Candida albicans* Biofilms: a Developmental State Associated With Specific and Stable Gene Expression Patterns**

Eukaryot. Cell, April 1, 2004; 3(2): 536 - 545.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[▶ HOME](#)

Y. H. Huang, D. Li, A. Winoto, and E. A. Robey

Distinct transcriptional programs in thymocytes responding to T cell receptor, Notch, and positive selection signals

PNAS, April 6, 2004; 101(14): 4936 - 4941.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology


[▶ HOME](#)

M. L. A. Jansen, P. Daran-Lapujade, J. H. de Winde, M. D. W. Piper, and J. T. Pronk

Prolonged Maltose-Limited Cultivation of *Saccharomyces cerevisiae* Selects for Cells with Improved Maltose Affinity and Hypersensitivity

Appl. Envir. Microbiol., April 1, 2004; 70(4): 1956 - 1963.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology

[▶ HOME](#)

M. Sonderegger, M. Jeppsson, B. Hahn-Hagerdal, and U. Sauer

Molecular Basis for Anaerobic Growth of *Saccharomyces cerevisiae* on Xylose, Investigated by Global Gene Expression and Metabolic Flux Analysis

Appl. Envir. Microbiol., April 1, 2004; 70(4): 2307 - 2317.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am J Physiol Renal Physiology [▶ HOME](#)

R. Mishra, S. N. Emancipator, C. Miller, T. Kern, and M. S. Simonson

Adipose differentiation-related protein and regulators of lipid homeostasis identified by gene expression profiling in the murine db/db diabetic kidney

Am J Physiol Renal Physiol, May 1, 2004; 286(5): F913 - F921.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology [▶ HOME](#)

E. D. Cahir-McFarland, K. Carter, A. Rosenwald, J. M. Giltneane, S. E. Henrickson, L. M. Staudt, and E. Kieff

Role of NF- κ B in Cell Survival and Transcription of Latent Membrane Protein 1-Expressing or Epstein-Barr Virus Latency III-Infected Cells

J. Virol., April 15, 2004; 78(8): 4108 - 4119.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of THORACIC AND CARDIOVASCULAR SURGERY [▶ HOME](#)

I. E. Konstantinov, J. G. Coles, C. Boscarino, M. Takahashi, J. Goncalves, J. Ritter, and G. S. Van Arsdell

Gene expression profiles in children undergoing cardiac surgery for right heart obstructive lesions

J. Thorac. Cardiovasc. Surg., March 1, 2004; 127(3): 746 - 754.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Endocrinology [▶ HOME](#)

N. Stahlberg, E. Rico-Bautista, R. M. Fisher, X. Wu, L. Cheung, A. Flores-Morales, G. Tybring, G. Norstedt, and P. Tollet-Egnell

Female-Predominant Expression of Fatty Acid Translocase/CD36 in Rat and Human Liver

Endocrinology, April 1, 2004; 145(4): 1972 - 1979.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Molecular and Cellular Biology [▶ HOME](#)

C. Muller-Tidow, B. Steffen, T. Cauvet, L. Tickenbrock, P. Ji, S. Diederichs, B. Sargin, G. Kohler, M. Stelljes, E. Puccetti,



M. Ruthardt, S. deVos, S. W. Hiebert, H. P. Koeffler, W. E. Berdel, and H. Serve

Translocation Products in Acute Myeloid Leukemia Activate the Wnt Signaling Pathway in Hematopoietic Cells

Mol. Cell. Biol., April 1, 2004; 24(7): 2890 - 2904.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research


[HOME](#)

F. M. Selaru, J. Yin, A. Olaru, Y. Mori, Y. Xu, S. H. Epstein, F. Sato, E. Deacu, S. Wang, A. Sterian, A. Fulton, J. M. Abraham, D. Shibata, C. Baquet, S. A. Stass, and S. J. Meltzer

An Unsupervised Approach to Identify Molecular Phenotypic Components Influencing Breast Cancer Features

Cancer Res., March 1, 2004; 64(5): 1584 - 1588.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology

[HOME](#)

A. Gravesen, B. Kallipolitis, K. Holmstrom, P. E. Hoiby, M. Ramnath, and S. Knochel

pbp2229-Mediated Nisin Resistance Mechanism in *Listeria monocytogenes* Confers Cross-Protection to Class IIa Bacteriocins and Affects Virulence Gene Expression

Appl. Envir. Microbiol., March 1, 2004; 70(3): 1669 - 1679.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J. A. Bernstein, P.-H. Lin, S. N. Cohen, and S. Lin-Chao

Global analysis of *Escherichia coli* RNA degradosome function using DNA microarrays

PNAS, March 2, 2004; 101(9): 2758 - 2763.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Clinical Investigation

[HOME](#)

E. Quaglino, S. Rolla, M. Iezzi, M. Spadaro, P. Musiani, C. De Giovanni, P. L. Lollini, S. Lanzardo, G. Forni, R. Sanges, S. Crispi, P. De Luca, R. Calogero, and F. Cavallo

Concordant morphologic and gene expression data show that a vaccine halts HER-2/neu preneoplastic lesions

J. Clin. Invest., March 1, 2004; 113(5): 709 - 717.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

P. Daran-Lapujade, M. L. A. Jansen, J.-M. Daran, W. van Gulik, J. H. de Winde, and J. T. Pronk

Role of Transcriptional Regulation in Controlling Fluxes in Central Carbon Metabolism of *Saccharomyces cerevisiae*: A CHEMOSTAT CULTURE STUDY

J. Biol. Chem., March 5, 2004; 279(10): 9125 - 9138.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

K. L. Meibom, X. B. Li, A. T. Nielsen, C.-Y. Wu, S. Roseman, and G. K. Schoolnik

The *Vibrio cholerae* chitin utilization program

PNAS, February 24, 2004; 101(8): 2524 - 2529.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

A. G. Passerini, D. C. Polacek, C. Shi, N. M. Francesco, E. Manduchi, G. R. Grant, W. F. Pritchard, S. Powell, G. Y. Chang, C. J. Stoeckert Jr., and P. F. Davies

Coexisting proinflammatory and antioxidative endothelial transcription profiles in a disturbed flow region of the adult porcine aorta

PNAS, February 24, 2004; 101(8): 2482 - 2487.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



diabetes

[HOME](#)

K. Susztak, E. Bottinger, A. Novetsky, D. Liang, Y. Zhu, E. Ciccone, D. Wu, S. Dunn, P. McCue, and K. Sharma

Molecular Profiling of Diabetic Mouse Kidney Reveals Novel Genes Linked to Glomerular Disease

Diabetes, March 1, 2004; 53(3): 784 - 794.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[HOME](#)

J. Z. Li, M. P. Vawter, D. M. Walsh, H. Tomita, S. J. Evans, P. V. Choudary, J. F. Lopez, A. Avelar, V. Shokoohi, T. Chung, O. Mesarwi, E. G. Jones, S. J. Watson, H. Akil, W. E. Bunney Jr, and R. M. Myers

Systematic changes in gene expression in postmortem human brains associated with tissue pH and terminal

medical conditions

Hum. Mol. Genet., March 15, 2004; 13(6): 609 - 616.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research


[HOME](#)

D. Patel, A. Incassati, N. Wang, and D. J. McCance

Human Papillomavirus Type 16 E6 and E7 Cause Polyploidy in Human Keratinocytes and Up-Regulation of G2-M-Phase Proteins

Cancer Res., February 15, 2004; 64(4): 1299 - 1306.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity

[HOME](#)

B. Raju, Y. Hoshino, K. Kuwabara, I. Belitskaya, S.

Prabhakar, A. Canova, J. A. Gold, R. Condos, R. I. Pine, S.

Brown, W. N. Rom, and M. D. Weiden

Aerosolized Gamma Interferon (IFN- γ) Induces Expression of the Genes Encoding the IFN- γ -Inducible 10-Kilodalton Protein but Not Inducible Nitric Oxide Synthase in the Lung during Tuberculosis

Infect. Immun., March 1, 2004; 72(3): 1275 - 1283.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

R. J. Feezor, H. N. Paddock, H. V. Baker, J. C. Varela, J.

Barreda, L. L. Moldawer, G. S. Schultz, and D. W. Mozingo

Temporal patterns of gene expression in murine cutaneous burn wound healing

Physiol Genomics, February 13, 2004; 16(3): 341 - 348.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[HOME](#)

S. T. Dorsam, C. M. Ferrell, G. P. Dorsam, M. K. Derynck, U.

Vijapurkar, D. Khodabakhsh, B. Pau, H. Bernstein, C. M.

Haqq, C. Largman, and H. J. Lawrence

The transcriptome of the leukemogenic homeoprotein HOXA9 in human hematopoietic cells

Blood, March 1, 2004; 103(5): 1676 - 1684.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Clinical Cancer Research

[HOME](#)


Z. Kote-Jarai, R. D. Williams, N. Cattini, M. Copeland, I.



Giddings, R. Wooster, R. H. tePoele, P. Workman, B. Gusterson, J. Peacock, G. Gui, C. Campbell, and R. Eeles
Gene Expression Profiling after Radiation-Induced DNA Damage Is Strongly Predictive of BRCA1 Mutation Carrier Status

Clin. Cancer Res., February 1, 2004; 10(3): 958 - 963.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

F. J. Livesey, T. L. Young, and C. L. Cepko
An analysis of the gene expression program of mammalian neural progenitor cells

PNAS, February 3, 2004; 101(5): 1374 - 1379.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

J. Fan, P. Tam, G. V. Woude, and Y. Ren
Normalization and analysis of cDNA microarrays using within-array replications applied to neuroblastoma cell response to a cytokine

PNAS, February 3, 2004; 101(5): 1135 - 1140.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Leukocyte Biology

[▶ HOME](#)

X. Zhang, Y. Kluger, Y. Nakayama, R. Poddar, C. Whitney, A. DeTora, S. M. Weissman, and P. E. Newburger

Gene expression in mature neutrophils: early responses to inflammatory stimuli

J. Leukoc. Biol., February 1, 2004; 75(2): 358 - 372.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

[▶ HOME](#)

J. P.T. Higgins, L. Wang, N. Kambham, K. Montgomery, V. Mason, S. U. Vogelmann, K. V. Lemley, P. O. Brown, J. D. Brooks, and M. van de Rijn

Gene Expression in the Normal Adult Human Kidney Assessed by Complementary DNA Microarray

Mol. Biol. Cell, February 1, 2004; 15(2): 649 - 656.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Molecular and Cellular Biology

[▶ HOME](#)

L. A. Scott, J. K. Vass, E. K. Parkinson, D. A. F. Gillespie, J. N. Winnie, and B. W. Ozanne



Invasion of Normal Human Fibroblasts Induced by v-Fos Is Independent of Proliferation, Immortalization, and the Tumor Suppressors p16INK4a and p53

Mol. Cell. Biol., February 15, 2004; 24(4): 1540 - 1559.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

M. Diarra-Mehrpour, S. Arrabal, A. Jalil, X. Pinson, C. Gaudin, G. Pietu, A. Pitaval, H. Ripoche, M. Eloit, D. Dormont, and S. Chouaib

Prion Protein Prevents Human Breast Carcinoma Cell Line from Tumor Necrosis Factor {alpha}-Induced Cell Death

Cancer Res., January 15, 2004; 64(2): 719 - 727.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J. Lapointe, C. Li, J. P. Higgins, M. van de Rijn, E. Bair, K. Montgomery, M. Ferrari, L. Egevad, W. Rayford, U. Bergerheim, P. Ekman, A. M. DeMarzo, R. Tibshirani, D. Botstein, P. O. Brown, J. D. Brooks, and J. R. Pollack

Gene expression profiling identifies clinically relevant subtypes of prostate cancer

PNAS, January 20, 2004; 101(3): 811 - 816.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[HOME](#)

B. M. Fine, M. Stanulla, M. Schrappe, M. Ho, S. Viehmann, J. Harbott, and L. M. Boxer

Gene expression patterns associated with recurrent chromosomal translocations in acute lymphoblastic leukemia

Blood, February 1, 2004; 103(3): 1043 - 1049.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

[HOME](#)

G. Mercier, N. Berthault, J. Mary, J. Peyre, A. Antoniadis, J.-P. Comet, A. Cornuejols, C. Froidevaux, and M. Dutreix

Biological detection of low radiation doses by combining results of two microarray analysis methods

Nucleic Acids Res., January 13, 2004; 32(1): e12 - 12.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

► HOME

D. Walker, M. Rolfe, A. Thompson, G. R. Moore, R. James, J. C. D. Hinton, and C. Kleanthous

Transcriptional Profiling of Colicin-Induced Cell Death of *Escherichia coli* MG1655 Identifies Potential Mechanisms by Which Bacteriocins Promote Bacterial Diversity

J. Bacteriol., February 1, 2004; 186(3): 866 - 869.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

R. O. Stuart, W. Wachsman, C. C. Berry, J. Wang-Rodriguez, L. Wasserman, I. Klacansky, D. Masys, K. Arden, S.

Goodison, M. McClelland, Y. Wang, A. Sawyers, I. Kalcheva, D. Tarin, and D. Mercola

In silico dissection of cell-type-associated patterns of gene expression in prostate cancer

PNAS, January 13, 2004; 101(2): 615 - 620.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

► HOME

P. Tollet-Egnell, P. Parini, N. Stahlberg, I. Lonnstedt, N. H. Lee, M. Rudling, A. Flores-Morales, and G. Norstedt

Growth hormone-mediated alteration of fuel metabolism in the aged rat as determined from transcript profiles

Physiol Genomics, January 15, 2004; 16(2): 261 - 267.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics


► HOME

D. Sanoudou, P. B. Kang, J. N. Haslett, M. Han, L. M. Kunkel, and A. H. Beggs

Transcriptional profile of postmortem skeletal muscle

Physiol Genomics, January 15, 2004; 16(2): 222 - 228.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology

► HOME

A. J. A. van Maris, J.-M. A. Geertman, A. Vermeulen, M. K. Groothuizen, A. A. Winkler, M. D. W. Piper, J. P. van Dijken, and J. T. Pronk

Directed Evolution of Pyruvate Decarboxylase-Negative *Saccharomyces cerevisiae*, Yielding a C2-Independent, Glucose-Tolerant, and Pyruvate-Hyperproducing Yeast

Appl. Envir. Microbiol., January 1, 2004; 70(1): 159 - 166.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am J Physiol Cell Physiol [▶ HOME](#)

I. A. Barash, L. Mathew, A. F. Ryan, J. Chen, and R. L. Lieber
Rapid muscle-specific gene expression changes after a single bout of eccentric contractions in the mouse

Am J Physiol Cell Physiol, February 1, 2004; 286(2): C355 - 364.

[\[Abstract\]](#) [\[Full Text\]](#)



JBC Online [▶ HOME](#)

A. Grechez-Cassiau, S. Panda, S. Lacoche, M. Teboul, S. Azmi, V. Laudet, J. B. Hogenesch, R. Taneja, and F. Delaunay
The Transcriptional Repressor STRA13 Regulates a Subset of Peripheral Circadian Outputs

J. Biol. Chem., January 9, 2004; 279(2): 1141 - 1150.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology [▶ HOME](#)

F. J. T. Staal, F. Weerkamp, M. R. M. Baert, C. M. M. van den Burg, M. van Noort, E. F. E. de Haas, and J. J. M. van Dongen

Wnt Target Genes Identified by DNA Microarrays in Immature CD34+ Thymocytes Regulate Proliferation and Cell Adhesion

J. Immunol., January 15, 2004; 172(2): 1099 - 1108.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Journal of Bacteriology [▶ HOME](#)

E. C. Gaynor, S. Cawthraw, G. Manning, J. K. MacKichan, S. Falkow, and D. G. Newell

The Genome-Sequenced Variant of Campylobacter jejuni NCTC 11168 and the Original Clonal Clinical Isolate Differ Markedly in Colonization, Gene Expression, and Virulence-Associated Phenotypes

J. Bacteriol., January 15, 2004; 186(2): 503 - 517.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular and Cellular Biology [▶ HOME](#)

L. Bruno, R. Hoffmann, F. McBlane, J. Brown, R. Gupta, C. Joshi, S. Pearson, T. Seidl, C. Heyworth, and T. Enver

Molecular Signatures of Self-Renewal, Differentiation,

and Lineage Choice in Multipotential Hemopoietic Progenitor Cells In Vitro

Mol. Cell. Biol., January 15, 2004; 24(2): 741 - 756.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

T. Kelly, H.-Q. Miao, Y. Yang, E. Navarro, P. Kussie, Y. Huang, V. MacLeod, J. Casciano, L. Joseph, F. Zhan, M. Zangari, B. Barlogie, J. Shaughnessy, and R. D. Sanderson

High Heparanase Activity in Multiple Myeloma Is Associated with Elevated Microvessel Density

Cancer Res., December 15, 2003; 63(24): 8749 - 8756.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

B. Kim, S. Bang, S. Lee, S. Kim, Y. Jung, C. Lee, K. Choi, S.-G. Lee, K. Lee, Y. Lee, S.-S. Kim, Y.-I. Yeom, Y.-S. Kim, H.-S. Yoo, K. Song, and I. Lee

Expression Profiling and Subtype-Specific Expression of Stomach Cancer

Cancer Res., December 1, 2003; 63(23): 8248 - 8255.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology


[HOME](#)

S. K. Banerjee, H. W. J. Young, A. Barczak, D. J. Erle, and M. R. Blackburn

Abnormal Alveolar Development Associated with Elevated Adenine Nucleosides

Am. J. Respir. Cell Mol. Biol., January 1, 2004; 30(1): 38 - 50.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation Research

[HOME](#)

R. Tabibiazar, R. A. Wagner, A. Liao, and T. Quertermous

Transcriptional Profiling of the Heart Reveals Chamber-Specific Gene Expression Patterns

Circ. Res., December 12, 2003; 93(12): 1193 - 1201.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY


[HOME](#)

R. I. Skotheim, A. Kallioniemi, B. Bjerkhagen, F. Mertens, H. R. Brekke, O. Monni, S. Mousses, N. Mandahl, G. Soeter, J. M. Nesland, S. Smeland, O.-P. Kallioniemi, and R. A. Lothe

Topoisomerase-II{alpha} Is Upregulated in Malignant Peripheral Nerve Sheath Tumors and Associated With Clinical Outcome

J. Clin. Oncol., December 15, 2003; 21(24): 4586 - 4591.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The EMBO Journal


[▶ HOME](#)

A. Rajpal, Y. A. Cho, B. Yelent, P. H. Koza-Taylor, D. Li, E. Chen, M. Whang, C. Kang, T. G. Turi, and A. Winoto

Transcriptional activation of known and novel apoptotic pathways by Nur77 orphan steroid receptor

EMBO J., December 15, 2003; 22(24): 6526 - 6536.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology


[▶ HOME](#)

K. Kretschmer, A. Jungebloud, J. Stopkiewicz, T. Kleinke, R. Hoffmann, and S. Weiss

The Selection of Marginal Zone B Cells Differs from That of B-1a Cells

J. Immunol., December 15, 2003; 171(12): 6495 - 6501.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[▶ HOME](#)

Y. Kun, L. C. How, T. P. Hoon, V. B. Bajic, T. S. Lam, A. Aggarwal, H. G. Sze, W. S. Bok, W. C. Yin, and P. Tan

Classifying the estrogen receptor status of breast cancers by expression profiles reveals a poor prognosis subpopulation exhibiting high expression of the ERBB2 receptor

Hum. Mol. Genet., December 15, 2003; 12(24): 3245 - 3258.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

I. Rogatsky, J.-C. Wang, M. K. Derynck, D. F. Nonaka, D. B. Khodabakhsh, C. M. Haqq, B. D. Darimont, M. J. Garabedian, and K. R. Yamamoto

Target-specific utilization of transcriptional regulatory surfaces by the glucocorticoid receptor

PNAS, November 25, 2003; 100(24): 13845 - 13850.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Nucleic Acids Research

[▶ HOME](#)



C. Romualdi, S. Trevisan, B. Celegato, G. Costa, and G. Lanfranchi

Improved detection of differentially expressed genes in microarray experiments through multiple scanning and image integration

Nucleic Acids Res., December 1, 2003; 31(23): e149 - 149.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of PATHOLOGY [▶ HOME](#)

S. C. Linn, R. B. West, J. R. Pollack, S. Zhu, T. Hernandez-Boussard, T. O. Nielsen, B. P. Rubin, R. Patel, J. R. Goldblum, D. Siegmund, D. Botstein, P. O. Brown, C. B. Gilks, and M. van de Rijn

Gene Expression Patterns and Gene Copy Number Changes in Dermatofibrosarcoma Protuberans

Am. J. Pathol., December 1, 2003; 163(6): 2383 - 2395.

[\[Abstract\]](#) [\[Full Text\]](#)



Development [▶ HOME](#)

R. Artero, E. E. Furlong, K. Beckett, M. P. Scott, and M. Baylies

Notch and Ras signaling pathway effector genes expressed in fusion competent and founder cells during Drosophila myogenesis

Development, December 22, 2003; 130(25): 6257 - 6272.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology [▶ HOME](#)

D. E. Dimcheff, S. Askovic, A. H. Baker, C. Johnson-Fowler, and J. L. Portis

Endoplasmic Reticulum Stress Is a Determinant of Retrovirus-Induced Spongiform Neurodegeneration

J. Virol., December 1, 2003; 77(23): 12617 - 12629.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences [▶ HOME](#)

J. M. Sperger, X. Chen, J. S. Draper, J. E. Antosiewicz, C. H. Chon, S. B. Jones, J. D. Brooks, P. W. Andrews, P. O. Brown, and J. A. Thomson

Gene expression patterns in human embryonic stem cells and human pluripotent germ cell tumors

PNAS, November 11, 2003; 100(23): 13350 - 13355.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



HEMATOLOGY

► HOME

W. L. Carroll, D. Bhojwani, D.-J. Min, E. Raetz, M. Relling, S. Davies, J. R. Downing, C. L. Willman, and J. C. Reed

Pediatric Acute Lymphoblastic Leukemia

Hematology, January 1, 2003; 2003(1): 102 - 131.

[\[Abstract\]](#) [\[Full Text\]](#)



Physiological Genomics

► HOME

H. Ida, S. A. Boylan, A. L. Weigel, and L. M. Hjelmeland

Age-related changes in the transcriptional profile of mouse RPE/choroid

Physiol Genomics, November 11, 2003; 15(3): 258 - 262.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory and Critical Care Medicine

► HOME

V. J. Gimino, J. D. Lande, T. R. Berryman, R. A. King, and M. I. Hertz

Gene Expression Profiling of Bronchoalveolar Lavage Cells in Acute Lung Rejection

Am. J. Respir. Crit. Care Med., November 15, 2003; 168(10): 1237 - 1242.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

► HOME

J. E. Korkola, S. DeVries, J. Fridlyand, E. S. Hwang, A. L. H. Estep, Y.-Y. Chen, K. L. Chew, S. H. Dairkee, R. M. Jensen, and F. M. Waldman

Differentiation of Lobular versus Ductal Breast Carcinomas by Expression Microarray Analysis

Cancer Res., November 1, 2003; 63(21): 7167 - 7175.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



MOLECULAR BIOLOGY AND EVOLUTION

► HOME

Z. Bochdanovits, H. van der Klis, and G. de Jong

Covariation of Larval Gene Expression and Adult Body Size in Natural Populations of *Drosophila melanogaster*

Mol. Biol. Evol., November 1, 2003; 20(11): 1760 - 1766.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

► HOME

M. E. Schaner, D. T. Ross, G. Ciaravino, T. Sorlie, O.

Troyanskaya, M. Diehn, Y. C. Wang, G. E. Duran, T. L. Sikic,

S. Caldeira, H. Skomedal, I-P. Tu, T. Hernandez-Boussard, S.

W. Johnson, P. J. O'Dwyer, M. J. Fero, G. B. Kristensen, A.-L. Borresen-Dale, T. Hastie, R. Tibshirani, M. van de Rijn, N. N. Teng, T. A. Longacre, D. Botstein, P. O. Brown, and B. I. Sikic

Gene Expression Patterns in Ovarian Carcinomas

Mol. Biol. Cell, November 1, 2003; 14(11): 4376 - 4386.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[▶ HOME](#)

C. Hemsley, E. Joyce, D. L. Hava, A. Kawale, and A. Camilli
MgrA, an Orthologue of Mga, Acts as a Transcriptional Repressor of the Genes within the rlrA Pathogenicity Islet in Streptococcus pneumoniae

J. Bacteriol., November 15, 2003; 185(22): 6640 - 6647.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory and Critical Care Medicine

[▶ HOME](#)

I. B. Copland, B. P. Kavanagh, D. Engelberts, C. McKerlie, J. Belik, and M. Post

Early Changes in Lung Gene Expression due to High Tidal Volume

Am. J. Respir. Crit. Care Med., November 1, 2003; 168(9): 1051 - 1059.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[▶ HOME](#)

I. Chowers, D. Liu, R. H. Farkas, T. L. Gunatilaka, A. S. Hackam, S. L. Bernstein, P. A. Campochiaro, G. Parmigiani, and D. J. Zack

Gene expression variation in the adult human retina

Hum. Mol. Genet., November 15, 2003; 12(22): 2881 - 2893.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

P. A. Clarke, M. L. George, S. Easdale, D. Cunningham, R. I. Swift, M. E. Hill, D. M. Tait, and P. Workman

Molecular Pharmacology of Cancer Therapy in Human Colorectal Cancer by Gene Expression Profiling

Cancer Res., October 15, 2003; 63(20): 6855 - 6863.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Clinical Cancer Research

[▶ HOME](#)



N. Yamagata, Y. Shyr, K. Yanagisawa, M. Edgerton, T. P. Dang, A. Gonzalez, S. Nadaf, P. Larsen, J. R. Roberts, J. C. Nesbitt, R. Jensen, S. Levy, J. H. Moore, J. D. Minna, and D. P. Carbone

A Training-Testing Approach to the Molecular Classification of Resected Non-Small Cell Lung Cancer

Clin. Cancer Res., October 15, 2003; 9(13): 4695 - 4704.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Endocrinology


[▶ HOME](#)

K. Dillner, J. Kindblom, A. Flores-Morales, R. Shao, J. Tornell, G. Norstedt, and H. Wennbo

Gene Expression Analysis of Prostate Hyperplasia in Mice Overexpressing the Prolactin Gene Specifically in the Prostate

Endocrinology, November 1, 2003; 144(11): 4955 - 4966.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENETICS

[▶ HOME](#)

T. Fukushige, B. Goszczynski, H. Tian, and J. D. McGhee

The Evolutionary Duplication and Probable Demise of an Endodermal GATA Factor in *Caenorhabditis elegans*

Genetics, October 1, 2003; 165(2): 575 - 588.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[▶ HOME](#)

M. Mirotsoy, C. M.H. Watanabe, P. G. Schultz, R. E. Pratt, and V. J. Dzau

Elucidating the molecular mechanism of cardiac remodeling using a comparative genomic approach

Physiol Genomics, October 17, 2003; 15(2): 115 - 126.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[▶ HOME](#)

B. C. Blaxall, R. Spang, H. A. Rockman, and W. J. Koch

Differential myocardial gene expression in the development and rescue of murine heart failure

Physiol Genomics, October 17, 2003; 15(2): 105 - 114.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

JBC Online

[▶ HOME](#)

G. Pandini, E. Medico, E. Conte, L. Sciacca, R. Vigneri, and A. Belfiore



Differential Gene Expression Induced by Insulin and Insulin-like Growth Factor-II through the Insulin Receptor Isoform A

J. Biol. Chem., October 24, 2003; 278(43): 42178 - 42189.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science

[▶ HOME](#)

N. O. Fortunel, H. H. Otu, H.-H. Ng, J. Chen, X. Mu, T. Chevassut, X. Li, M. Joseph, C. Bailey, J. A. Hatzfeld, A. Hatzfeld, F. Usta, V. B. Vega, P. M. Long, T. A. Libermann, and B. Lim

Comment on " 'Stemness': Transcriptional Profiling of Embryonic and Adult Stem Cells" and "A Stem Cell Molecular Signature" (I)

Science, October 17, 2003; 302(5644): 393b - 393.

[\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

M. L. Whitfield, D. R. Finlay, J. I. Murray, O. G. Troyanskaya, J.-T. Chi, A. Pergamenschikov, T. H. McCalmont, P. O. Brown, D. Botstein, and M. K. Connolly

Systemic and cell type-specific gene expression patterns in scleroderma skin

PNAS, October 14, 2003; 100(21): 12319 - 12324.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

A. Mueller, J. O'Rourke, P. Chu, C. C. Kim, P. Sutton, A. Lee, and S. Falkow

Protective immunity against Helicobacter is characterized by a unique transcriptional signature

PNAS, October 14, 2003; 100(21): 12289 - 12294.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am J Physiol Cell Physiol

[▶ HOME](#)

R. Mishra, P. Leahy, and M. S. Simonson

Gene expression profile of endothelin-1-induced growth in glomerular mesangial cells

Am J Physiol Cell Physiol, November 1, 2003; 285(5): C1109 - 1115.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Physiological Genomics

[▶ HOME](#)



M. Fornage, M. W. Swank, E. Boerwinkle, and P. A. Doris
Gene expression profiling and functional proteomic analysis reveal perturbed kinase-mediated signaling in genetic stroke susceptibility

Physiol Genomics, September 29, 2003; 15(1): 75 - 83.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

J. S. Pattison, L. C. Folk, R. W. Madsen, T. E. Childs, and F. W. Booth

Transcriptional profiling identifies extensive downregulation of extracellular matrix gene expression in sarcopenic rat soleus muscle

Physiol Genomics, September 29, 2003; 15(1): 34 - 43.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

[HOME](#)

H. Zhao, F. Granberg, L. Elfineh, U. Pettersson, and C. Svensson

Strategic Attack on Host Cell Gene Expression during Adenovirus Infection

J. Virol., October 15, 2003; 77(20): 11006 - 11015.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

[HOME](#)

H. E. Neuman de Vegvar, R. R. Amara, L. Steinman, P. J. Utz, H. L. Robinson, and W. H. Robinson

Microarray Profiling of Antibody Responses against Simian-Human Immunodeficiency Virus: Postchallenge Convergence of Reactivities Independent of Host Histocompatibility Type and Vaccine Regimen

J. Virol., October 15, 2003; 77(20): 11125 - 11138.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

S. Hassan, B. Duong, K.-S. Kim, and M. F. Miles

Pharmacogenomic Analysis of Mechanisms Mediating Ethanol Regulation of Dopamine {beta}-Hydroxylase

J. Biol. Chem., October 3, 2003; 278(40): 38860 - 38869.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Cancer Research

[HOME](#)

U. Woelfle, J. Cloos, G. Sauter, L. Riethdorf, F. Janicke, P.



van Diest, R. Brakenhoff, and K. Pantel

Molecular Signature Associated with Bone Marrow Micrometastasis in Human Breast Cancer

Cancer Res., September 15, 2003; 63(18): 5679 - 5684.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



BIOLOGY of REPRODUCTION

[HOME](#)

X. Wu, S.-T. Pang, L. Sahlin, A. Blanck, G. Norstedt, and A. Flores-Morales

Gene Expression Profiling of the Effects of Castration and Estrogen Treatment in the Rat Uterus

Biol Reprod, October 1, 2003; 69(4): 1308 - 1317.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J.-T. Chi, H. Y. Chang, G. Haraldsen, F. L. Jahnsen, O. G. Troyanskaya, D. S. Chang, Z. Wang, S. G. Rockson, M. van de Rijn, D. Botstein, and P. O. Brown

Endothelial cell diversity revealed by global expression profiling

PNAS, September 16, 2003; 100(19): 10623 - 10628.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[HOME](#)

M. J. Kazmierczak, S. C. Mithoe, K. J. Boor, and M. Wiedmann

Listeria monocytogenes {sigma}B Regulates Stress Response and Virulence Functions

J. Bacteriol., October 1, 2003; 185(19): 5722 - 5734.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[HOME](#)

C. R. Scherzer, R. V. Jensen, S. R. Gullans, and M. B. Feany
Gene expression changes presage neurodegeneration in a Drosophila model of Parkinson's disease

Hum. Mol. Genet., October 1, 2003; 12(19): 2457 - 2466.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH

[HOME](#)

K.-Y. Jen and V. G. Cheung

Transcriptional Response of Lymphoblastoid Cells to Ionizing Radiation

Genome Res., September 1, 2003; 13(9): 2092 - 2100.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

Y.-C. Yang, E. Piek, J. Zavadil, D. Liang, D. Xie, J. Heyer, P. Pavlidis, R. Kucherlapati, A. B. Roberts, and E. P. Bottinger
Hierarchical model of gene regulation by transforming growth factor {beta}

PNAS, September 2, 2003; 100(18): 10269 - 10274.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Experimental Medicine

[HOME](#)

D. Schnappinger, S. Ehrt, M. I. Voskuil, Y. Liu, J. A. Mangan, I. M. Monahan, G. Dolganov, B. Efron, P. D. Butcher, C. Nathan, and G. K. Schoolnik

Transcriptional Adaptation of Mycobacterium tuberculosis within Macrophages: Insights into the Phagosomal Environment

J. Exp. Med., September 1, 2003; 198(5): 693 - 704.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science

Science of Aging Knowledge Environment

[HOME](#)

K. G. Becker

Deciphering the Gene Expression Profile of Long-Lived Snell Mice

Sci. Aging Knowl. Environ., March 20, 2002; 2002(11): pe4 - 4.

[\[Abstract\]](#) [\[Full Text\]](#)



Investigative Ophthalmology & Visual Science

[HOME](#)

I. Chowers, T. L. Gunatilaka, R. H. Farkas, J. Qian, A. S. Hackam, E. Duh, M. Kageyama, C. Wang, A. Vora, P. A. Campochiaro, and D. J. Zack

Identification of Novel Genes Preferentially Expressed in the Retina Using a Custom Human Retina cDNA Microarray

Invest. Ophthalmol. Vis. Sci., September 1, 2003; 44(9): 3732 - 3741.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science

[HOME](#)

M. T. Nishimura, M. Stein, B.-H. Hou, J. P. Vogel, H. Edwards, and S. C. Somerville

Loss of a Callose Synthase Results in Salicylic Acid-

Dependent Disease Resistance

Science, August 15, 2003; 301(5635): 969 - 972.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research [▶ HOME](#)

S. Singhal, R. Wiewrodt, L. D. Malden, K. M. Amin, K. Matzie, J. Friedberg, J. C. Kucharczuk, L. A. Litzky, S. W. Johnson, L. R. Kaiser, and S. M. Albelda

Gene Expression Profiling of Malignant Mesothelioma

Clin. Cancer Res., August 1, 2003; 9(8): 3080 - 3097.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Cancer Research [▶ HOME](#)

P. K. Ha, N. E. Benoit, R. Yochem, J. Sciubba, M. Zahurak, D. Sidransky, J. Pevsner, W. H. Westra, and J. Califano

A Transcriptional Progression Model for Head and Neck Cancer

Clin. Cancer Res., August 1, 2003; 9(8): 3058 - 3064.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Proceedings of the National Academy of Sciences [▶ HOME](#)

J. D. Storey and R. Tibshirani

Statistical significance for genomewide studies

PNAS, August 5, 2003; 100(16): 9440 - 9445.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Applied and Environmental Microbiology [▶ HOME](#)

Z. Vuralhan, M. A. Morais, S.-L. Tai, M. D. W. Piper, and J. T. Pronk

Identification and Characterization of Phenylpyruvate Decarboxylase Genes in *Saccharomyces cerevisiae*

Appl. Envir. Microbiol., August 1, 2003; 69(8): 4534 - 4541.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE FASEB JOURNAL [▶ HOME](#)

N. F. MARKO, P. B. DIEFFENBACH, G. YAN, S. CERYAK, R. W. HOWELL, T. A. MCCAFFREY, and V. W. HU

Does metabolic radiolabeling stimulate the stress response? Gene expression profiling reveals differential cellular responses to internal beta vs. external gamma radiation

FASEB J, August 1, 2003; 17(11): 1470 - 1486.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology

[HOME](#)

C. A. Powell, A. Spira, A. Derti, C. DeLisi, G. Liu, A. Borczuk, S. Busch, S. Sahasrabudhe, Y. Chen, D. Sugarbaker, R. Bueno, W. G. Richards, and J. S. Brody

Gene Expression in Lung Adenocarcinomas of Smokers and Nonsmokers

Am. J. Respir. Cell Mol. Biol., August 1, 2003; 29(2): 157 - 162.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology


[HOME](#)

K. Kretschmer, A. Jungebloud, J. Stopkiewicz, B. Stoermann, R. Hoffmann, and S. Weiss

Antibody Repertoire and Gene Expression Profile: Implications for Different Developmental and Functional Traits of Splenic and Peritoneal B-1 Lymphocytes

J. Immunol., August 1, 2003; 171(3): 1192 - 1201.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of the American Society of Nephrology


[HOME](#)

K. Susztak, K. Sharma, M. Schiffer, P. McCue, E. Ciccone, and E. P. Bottinger

Genomic Strategies for Diabetic Nephropathy

J. Am. Soc. Nephrol., August 1, 2003; 14(90003): S271 - 278.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Animal Science

[HOME](#)

A. Reverter, K. A. Byrne, H. L. Bruce, Y. H. Wang, B. P. Dalrymple, and S. A. Lehnert

A mixture model-based cluster analysis of DNA microarray gene expression data on Brahman and Brahman composite steers fed high-, medium-, and low-quality diets

J Anim Sci, August 1, 2003; 81(8): 1900 - 1910.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[HOME](#)


E. Mongodin, J. Finan, M. W. Climo, A. Rosato, S. Gill, and G. L. Archer

Microarray Transcription Analysis of Clinical

Staphylococcus aureus Isolates Resistant to Vancomycin

J. Bacteriol., August 1, 2003; 185(15): 4638 - 4643.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Microbiology [▶ HOME](#)

J. E. Dove, K. Yasukawa, C. R. Tinsley, and X. Nassif Production of the signalling molecule, autoinducer-2, by *Neisseria meningitidis*: lack of evidence for a concerted transcriptional response

Microbiology, July 1, 2003; 149(7): 1859 - 1869.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



THE NEW ENGLAND JOURNAL OF MEDICINE [▶ HOME](#)

M. Sarwal, M.-S. Chua, N. Kambham, S.-C. Hsieh, T. Satterwhite, M. Masek, and O. Salvatierra Jr.

Molecular Heterogeneity in Acute Renal Allograft Rejection Identified by DNA Microarray Profiling

N. Engl. J. Med., July 10, 2003; 349(2): 125 - 138.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics [▶ HOME](#)

S. Welle, A. I. Brooks, J. M. Delehanty, N. Needler, and C. A. Thornton

Gene expression profile of aging in human muscle

Physiol Genomics, July 7, 2003; 14(2): 149 - 159.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Physiological Genomics [▶ HOME](#)

S. Keay, F. Seillier-Moisewitsch, C.-O. Zhang, T. C. Chai, and J. Zhang

Changes in human bladder epithelial cell gene expression associated with interstitial cystitis or antiproliferative factor treatment

Physiol Genomics, July 7, 2003; 14(2): 107 - 115.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



ANNALS of the New York Academy of Sciences [▶ HOME](#)

R. M. MILLER, C. CASACELI, L. CHEN, and H. J. FEDEROFF

Gene Expression Analysis of the MPTP-Lesioned Substantia Nigra in Mice

Ann. N.Y. Acad. Sci., June 1, 2003; 991(1): 315 - 318.

[\[Full Text\]](#) [\[PDF\]](#)



Journal of Virology

► HOME

K. Lohr, O. Hartmann, H. Schafer, and M. Dobbelstein
Mutual Interference of Adenovirus Infection and myc Expression

J. Virol., July 15, 2003; 77(14): 7936 - 7944.

[Abstract] [Full Text] [PDF]



Nucleic Acids Research

► HOME

N. M. Luscombe, T. E. Royce, P. Bertone, N. Echols, C. E. Horak, J. T. Chang, M. Snyder, and M. Gerstein
ExpressYourself: a modular platform for processing and visualizing microarray data

Nucleic Acids Res., July 1, 2003; 31(13): 3477 - 3482.

[Abstract] [Full Text] [PDF]



JBC Online

► HOME

J. R. Brown, M. M. Fuster, T. Whisenant, and J. D. Esko
Expression Patterns of {alpha}2,3-Sialyltransferases and {alpha}1,3-Fucosyltransferases Determine the Mode of Sialyl Lewis X Inhibition by Disaccharide Decoys

J. Biol. Chem., June 27, 2003; 278(26): 23352 - 23359.

[Abstract] [Full Text] [PDF]



The American Journal of Pathology

► HOME

D. E. Hansel, A. Rahman, M. Hidalgo, P. J. Thuluvath, K. D. Lillemoe, R. Shulick, J.-L. Ku, J.-G. Park, K. Miyazaki, R. Ashfaq, I. I. Wistuba, R. Varma, L. Hawthorne, J. Geradts, P. Argani, and A. Maitra

Identification of Novel Cellular Targets in Biliary Tract Cancers Using Global Gene Expression Technology

Am. J. Pathol., July 1, 2003; 163(1): 217 - 229.

[Abstract] [Full Text] [PDF]



Plant Physiology


► HOME

J. O. Borevitz and M. Nordborg

The Impact of Genomics on the Study of Natural Variation in Arabidopsis

Plant Physiology, June 1, 2003; 132(2): 718 - 725.

[Full Text] [PDF]



Proceedings of the National Academy of Sciences

▶ HOME

J.-T. Chi, H. Y. Chang, N. N. Wang, D. S. Chang, N. Dunphy,
and P. O. Brown

**Genomewide view of gene silencing by small
interfering RNAs**

PNAS, May 27, 2003; 100(11): 6343 - 6346.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology


▶ HOME

S. Perkowski, J. Sun, S. Singhal, J. Santiago, G. D. Leikauf,
and S. M. Albelda

**Gene Expression Profiling of the Early Pulmonary
Response to Hyperoxia in Mice**

Am. J. Respir. Cell Mol. Biol., June 1, 2003; 28(6): 682 -
696.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

R. M. Berka, X. Cui, and C. Yanofsky

**Genomewide transcriptional changes associated with
genetic alterations and nutritional supplementation
affecting tryptophan metabolism in *Bacillus subtilis***

PNAS, May 13, 2003; 100(10): 5682 - 5687.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

▶ HOME

M. Ho, E. Yang, G. Matcuk, D. Deng, N. Sampas, A. Tsalenko,
R. Tabibiazar, Y. Zhang, M. Chen, S. Talbi, Y. D. Ho, J. Wang,
P. S. Tsao, A. Ben-Dor, Z. Yakhini, L. Bruhn, and T.
Quertermous

**Identification of endothelial cell genes by combined
database mining and microarray analysis**

Physiol Genomics, May 13, 2003; 13(3): 249 - 262.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

▶ HOME

S. Rome, K. Clement, R. Rabasa-Lhoret, E. Loizon, C. Poitou,
G. S. Barsh, J.-P. Riou, M. Laville, and H. Vidal

**Microarray Profiling of Human Skeletal Muscle Reveals
That Insulin Regulates ~800 Genes during a
Hyperinsulinemic Clamp**

J. Biol. Chem., May 16, 2003; 278(20): 18063 - 18068.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood


HOME

J. Shaughnessy, J. Jacobson, J. Sawyer, J. McCoy, A. Fassas, F. Zhan, K. Bumm, J. Epstein, E. Anaissie, S. Jagannath, D. Vesole, D. Siegel, R. Desikan, N. Munshi, A. Badros, E. Tian, M. Zangari, G. Tricot, J. Crowley, and B. Barlogie

Continuous absence of metaphase-defined cytogenetic abnormalities, especially of chromosome 13 and hypodiploidy, ensures long-term survival in multiple myeloma treated with Total Therapy I: interpretation in the context of global gene expression

Blood, May 15, 2003; 101(10): 3849 - 3856.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Nutrition

HOME

R. J. Cousins, R. K. Blanchard, J. B. Moore, L. Cui, C. L. Green, J. P. Liuzzi, J. Cao, and J. A. Bobo

Regulation of Zinc Metabolism and Genomic Outcomes

J. Nutr., May 1, 2003; 133(5): 1521S - 1526.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Chemistry


HOME

T. C.W. Poon, T.-T. Yip, A. T.C. Chan, C. Yip, V. Yip, T. S.K. Mok, C. C.Y. Lee, T. W.T. Leung, S. K.W. Ho, and P. J. Johnson

Comprehensive Proteomic Profiling Identifies Serum Proteomic Signatures for Detection of Hepatocellular Carcinoma and Its Subtypes

Clin. Chem., May 1, 2003; 49(5): 752 - 760.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



the Journal of Molecular Diagnostics

HOME

C. Cheadle, M. P. Vawter, W. J. Freed, and K. G. Becker

Analysis of Microarray Data Using Z Score Transformation

Mol. Diagn., May 1, 2003; 5(2): 73 - 81.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Investigative Ophthalmology & Visual Science

HOME


S. Goswami, N. L. Sheets, J. Zavadil, B. K. Chauhan, E. P. Bottinger, V. N. Reddy, M. Kantorow, and A. Cvekl

Spectrum and Range of Oxidative Stress Responses of Human Lens Epithelial Cells to H₂O₂ Insult

Invest. Ophthalmol. Vis. Sci., May 1, 2003; 44(5): 2084 -

2093.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Infection and Immunity


[HOME](#)

L. J. Thompson, D. S. Merrell, B. A. Neilan, H. Mitchell, A. Lee, and S. Falkow

Gene Expression Profiling of *Helicobacter pylori* Reveals a Growth-Phase-Dependent Switch in Virulence Gene Expression

Infect. Immun., May 1, 2003; 71(5): 2643 - 2655.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[HOME](#)

D. Sanoudou, J. N. Haslett, A. T. Kho, S. Guo, H. T. Gazda, S. A. Greenberg, H. G. W. Lidov, I. S. Kohane, L. M. Kunkel, and A. H. Beggs

Expression profiling reveals altered satellite cell numbers and glycolytic enzyme transcription in nemaline myopathy muscle

PNAS, April 15, 2003; 100(8): 4666 - 4671.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF THE NATIONAL CANCER INSTITUTE

[HOME](#)

G. J. Gordon, R. V. Jensen, L.-L. Hsiao, S. R. Gullans, J. E. Blumenstock, W. G. Richards, M. T. Jaklitsch, D. J. Sugarbaker, and R. Bueno

Using Gene Expression Ratios to Predict Outcome Among Patients With Mesothelioma

J Natl Cancer Inst, April 16, 2003; 95(8): 598 - 605.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

[HOME](#)

B. Enjalbert, A. Nantel, and M. Whiteway

Stress-induced Gene Expression in *Candida albicans*: Absence of a General Stress Response

Mol. Biol. Cell, April 1, 2003; 14(4): 1460 - 1467.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular & Cellular PROTEOMICS

[HOME](#)

G. Chen, H. Wang, T. G. Gharib, C.-C. Huang, D. G. Thomas, K. A. Shedden, R. Kuick, J. M. G. Taylor, S. L. R. Kardia, D. E. Misek, T. J. Giordano, M. D. Iannettoni, M. B. Orringer, S. M. Hanash, and D. G. Beer

Overexpression of Oncoprotein 18 Correlates with Poor Differentiation in Lung Adenocarcinomas

Mol. Cell. Proteomics, February 1, 2003; 2(2): 107 - 116.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH

[HOME](#)

L. R. Nunes, Y. B. Rosato, N. H. Muto, G. M. Yanai, V. S. da Silva, D. B. Leite, E. R. Goncalves, A. A. de Souza, H. D. Coletta-Filho, M. A. Machado, S. A. Lopes, and R. Costa de Oliveira

Microarray Analyses of *Xylella fastidiosa* Provide Evidence of Coordinated Transcription Control of Laterally Transferred Elements

Genome Res., April 1, 2003; 13(4): 570 - 578.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[HOME](#)

M. Mizuki, J. Schwable, C. Steur, C. Choudhary, S. Agrawal, B. Sargin, B. Steffen, I. Matsumura, Y. Kanakura, F. D. Bohmer, C. Muller-Tidow, W. E. Berdel, and H. Serve

Suppression of myeloid transcription factors and induction of STAT response genes by AML-specific Flt3 mutations

Blood, April 15, 2003; 101(8): 3164 - 3173.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Neuroscience

[HOME](#)

R. C. Elliott, M. F. Miles, and D. H. Lowenstein

Overlapping Microarray Profiles of Dentate Gyrus Gene Expression during Development- and Epilepsy-Associated Neurogenesis and Axon Outgrowth

J. Neurosci., March 15, 2003; 23(6): 2218 - 2227.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



MOLECULAR ENDOCRINOLOGY

[HOME](#)

M. W. M. Yao, H. Lim, D. J. Schust, S. E. Choe, A. Farago, Y. Ding, S. Michaud, G. M. Church, and R. L. Maas

Gene Expression Profiling Reveals Progesterone-Mediated Cell Cycle and Immunoregulatory Roles of Hoxa-10 in the Preimplantation Uterus

Mol. Endocrinol., April 1, 2003; 17(4): 610 - 627.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

► HOME

W. Yu, N. R. Murray, C. Weems, L. Chen, H. Guo, R. Ethridge, J. D. Ceci, B. M. Evers, E. A. Thompson, and A. P. Fields

Role of Cyclooxygenase 2 in Protein Kinase C beta II-mediated Colon Carcinogenesis

J. Biol. Chem., March 28, 2003; 278(13): 11167 - 11174.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of PATHOLOGY

► HOME

C. A. Iacobuzio-Donahue, A. Maitra, M. Olsen, A. W. Lowe, N. T. Van Heek, C. Rosty, K. Walter, N. Sato, A. Parker, R. Ashfaq, E. Jaffee, B. Ryu, J. Jones, J. R. Eshleman, C. J. Yeo, J. L. Cameron, S. E. Kern, R. H. Hruban, P. O. Brown, and M. Goggins

Exploration of Global Gene Expression Patterns in Pancreatic Adenocarcinoma Using cDNA Microarrays

Am. J. Pathol., April 1, 2003; 162(4): 1151 - 1162.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

S. Beltran, E. Blanco, F. Serras, B. Perez-Villamil, R. Guigo, S. Artavanis-Tsakonas, and M. Corominas

Transcriptional network controlled by the trithorax-group gene ash2 in Drosophila melanogaster

PNAS, March 18, 2003; 100(6): 3293 - 3298.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

► HOME

H. Zhang, K.-H. Pan, and S. N. Cohen

Senescence-specific gene expression fingerprints reveal cell-type-dependent physical clustering of up-regulated chromosomal loci

PNAS, March 18, 2003; 100(6): 3251 - 3256.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH

► HOME

J. O. Borevitz, D. Liang, D. Plouffe, H.-S. Chang, T. Zhu, D. Weigel, C. C. Berry, E. Winzeler, and J. Chory

Large-Scale Identification of Single-Feature Polymorphisms in Complex Genomes

Genome Res., March 1, 2003; 13(3): 513 - 523.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The American Journal of PATHOLOGY

▶ HOME

J. P.T. Higgins, R. Shinghal, H. Gill, J. H. Reese, M. Terris, R. J. Cohen, M. Fero, J. R. Pollack, M. van de Rijn, and J. D. Brooks

Gene Expression Patterns in Renal Cell Carcinoma Assessed by Complementary DNA Microarray

Am. J. Pathol., March 1, 2003; 162(3): 925 - 932.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

T.-W. Chun, J. S. Justement, R. A. Lempicki, J. Yang, G. Dennis Jr., C. W. Hallahan, C. Sanford, P. Pandya, S. Liu, M. McLaughlin, L. A. Ehler, S. Moir, and A. S. Fauci

Gene expression and viral production in latently infected, resting CD4+ T cells in viremic versus aviremic HIV-infected individuals

PNAS, February 18, 2003; 100(4): 1908 - 1913.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

A. R. Whitney, M. Diehn, S. J. Popper, A. A. Alizadeh, J. C. Boldrick, D. A. Relman, and P. O. Brown

Individuality and variation in gene expression patterns in human blood

PNAS, February 18, 2003; 100(4): 1896 - 1901.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Plant Physiology


▶ HOME

L.-X. Yu and T. L. Setter

Comparative Transcriptional Profiling of Placenta and Endosperm in Developing Maize Kernels in Response to Water Deficit

Plant Physiology, February 1, 2003; 131(2): 568 - 582.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

A. Mueller, J. O'Rourke, J. Grimm, K. Guillemin, M. F. Dixon, A. Lee, and S. Falkow

Distinct gene expression profiles characterize the histopathological stages of disease in Helicobacter-induced mucosa-associated lymphoid tissue lymphoma

PNAS, February 4, 2003; 100(3): 1292 - 1297.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

▶ HOME

T. F. Cooper, D. E. Rozen, and R. E. Lenski
Parallel changes in gene expression after 20,000 generations of evolution in Escherichiacoli

PNAS, February 4, 2003; 100(3): 1072 - 1077.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



RECENT PROGRESS IN HORMONE RESEARCH

▶ HOME

J. A. Copland, P. J. Davies, G. L. Shipley, C. G. Wood, B. A. Luxon, and R. J. Urban

The Use of DNA Microarrays to Assess Clinical Samples: The Transition from Bedside to Bench to Bedside

Recent Prog. Horm. Res., January 1, 2003; 58(1): 25 - 53.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



RECENT PROGRESS IN HORMONE RESEARCH

▶ HOME

N.M. Svrakic, O. Nestic, M.R.K. Dasu, D. Herndon, and J.R. Perez-Polo

Statistical Approach to DNA Chip Analysis

Recent Prog. Horm. Res., January 1, 2003; 58(1): 75 - 93.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Nucleic Acids Research

▶ HOME

L. Smith, P. Underhill, C. Pritchard, Z. Tymowska-Lalanne, S. Abdul-Hussein, H. Hilton, L. Winchester, D. Williams, T. Freeman, S. Webb, and A. Greenfield

Single primer amplification (SPA) of cDNA for microarray expression analysis

Nucleic Acids Res., February 1, 2003; 31(3): e9 - 9.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

▶ HOME

V. M. Boer, J. H. de Winder, J. T. Pronk, and M. D. W. Piper
The Genome-wide Transcriptional Responses of Saccharomyces cerevisiae Grown on Glucose in Aerobic Chemostat Cultures Limited for Carbon, Nitrogen, Phosphorus, or Sulfur

J. Biol. Chem., January 31, 2003; 278(5): 3265 - 3274.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology

HOME

V. V. Yurovsky

Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand Enhances Collagen Production by Human Lung Fibroblasts

Am. J. Respir. Cell Mol. Biol., February 1, 2003; 28(2): 225 - 231.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Biology of the Cell

HOME

D. Chen, W. M. Toone, J. Mata, R. Lyne, G. Burns, K. Kivinen, A. Brazma, N. Jones, and J. Bahler

Global Transcriptional Responses of Fission Yeast to Environmental Stress

Mol. Biol. Cell, January 1, 2003; 14(1): 214 - 229.

[\[Abstract\]](#) [\[Full Text\]](#)



Physiological Genomics

HOME

M. Steenman, Y.-W. Chen, M. Le Cunff, G. Lamirault, A. Varro, E. Hoffman, and J. J. Leger

Transcriptomal analysis of failing and nonfailing human hearts

Physiol Genomics, January 15, 2003; 12(2): 97 - 112.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

HOME

A. M. Bronikowski, P. A. Carter, T. J. Morgan, T. Garland Jr, N. Ung, T. D. Pugh, R. Weindruch, and T. A. Prolla

Lifelong voluntary exercise in the mouse prevents age-related alterations in gene expression in the heart

Physiol Genomics, January 15, 2003; 12(2): 129 - 138.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

HOME

A. L. Astier, R. Xu, M. Svoboda, E. Hinds, O. Munoz, R. de Beaumont, C. D. Crean, T. Gabig, and A. S. Freedman

Temporal gene expression profile of human precursor B leukemia cells induced by adhesion receptor: identification of pathways regulating B-cell survival

Blood, February 1, 2003; 101(3): 1118 - 1127.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

blood

HOME



I. S. Lossos, A. A. Alizadeh, R. Rajapaksa, R. Tibshirani, and R. Levy

HGAL is a novel interleukin-4-inducible gene that strongly predicts survival in diffuse large B-cell lymphoma

Blood, January 15, 2003; 101(2): 433 - 440.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Journal of Virology [▶ HOME](#)

J. O. Jones and A. M. Arvin

Microarray Analysis of Host Cell Gene Transcription in Response to Varicella-Zoster Virus Infection of Human T Cells and Fibroblasts In Vitro and SCIDhu Skin Xenografts In Vivo

J. Virol., December 20, 2002; 77(2): 1268 - 1280.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular Cancer Therapeutics [▶ HOME](#)

T. R. Sutter, X.-R. He, P. Dimitrov, L. Xu, G. Narasimhan, E. O. George, C. H. Sutter, C. Grubbs, R. Savory, M. Stephan-Gueldner, D. Kreder, M. J. Taylor, R. Lubet, T. A. Patterson, and T. W. Kensler

Multiple Comparisons Model-based Clustering and Ternary Pattern Tree Numerical Display of Gene Response to Treatment: Procedure and Application to the Preclinical Evaluation of Chemopreventive Agents

Mol. Cancer Ther., December 1, 2002; 1(14): 1283 - 1292.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Antimicrobial Agents and Chemotherapy [▶ HOME](#)

**R. W. Hong, M. Shchepetov, J. N. Weiser, and P. H. Axelsen
Transcriptional Profile of the Escherichia coli Response to the Antimicrobial Insect Peptide Cecropin A**

Antimicrob. Agents Chemother., January 1, 2003; 47(1): 1 - 6.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



The Journal of Immunology [▶ HOME](#)

**R. L. Hengel, V. Thaker, M. V. Pavlick, J. A. Metcalf, G. Dennis Jr., J. Yang, R. A. Lempicki, I. Sereti, and H. C. Lane
Cutting Edge: L-Selectin (CD62L) Expression Distinguishes Small Resting Memory CD4+ T Cells That Preferentially Respond to Recall Antigen**

J. Immunol., January 1, 2003; 170(1): 28 - 32.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology


[▶ HOME](#)

J. D. Helmann, M. F. W. Wu, A. Gaballa, P. A. Kobel, M. M. Morshedi, P. Fawcett, and C. Paddon

The Global Transcriptional Response of *Bacillus subtilis* to Peroxide Stress Is Coordinated by Three Transcription Factors

J. Bacteriol., January 1, 2003; 185(1): 243 - 253.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Arteriosclerosis, Thrombosis, and Vascular Biology

[▶ HOME](#)

R. L. Geary, J. M. Wong, A. Rossini, S. M. Schwartz, and L. D. Adams

Expression Profiling Identifies 147 Genes Contributing to a Unique Primate Neointimal Smooth Muscle Cell Phenotype

Arterioscler. Thromb. Vasc. Biol., December 1, 2002; 22(12): 2010 - 2016.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Endocrinology

[▶ HOME](#)

S.-T. Pang, K. Dillner, X. Wu, A. Pousette, G. Norstedt, and A. Flores-Morales

Gene Expression Profiling of Androgen Deficiency Predicts a Pathway of Prostate Apoptosis that Involves Genes Related to Oxidative Stress

Endocrinology, December 1, 2002; 143(12): 4897 - 4906.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



HEMATOLOGY

[▶ HOME](#)

K. C. Anderson, J. D. Shaughnessy Jr., B. Barlogie, J.-L. Harousseau, and G. D. Roodman

Multiple Myeloma

Hematology, January 1, 2002; 2002(1): 214 - 240.

[\[Abstract\]](#) [\[Full Text\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

K. Guillemin, N. R. Salama, L. S. Tompkins, and S. Falkow
Cag pathogenicity island-specific responses of gastric epithelial cells to *Helicobacter pylori* infection

PNAS, November 12, 2002; 99(23): 15136 - 15141.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J. N. Haslett, D. Sanoudou, A. T. Kho, R. R. Bennett, S. A. Greenberg, I. S. Kohane, A. H. Beggs, and L. M. Kunkel
Gene expression comparison of biopsies from Duchenne muscular dystrophy (DMD) and normal skeletal muscle

PNAS, November 12, 2002; 99(23): 15000 - 15005.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[HOME](#)

M. L. Paustian, B. J. May, D. Cao, D. Boley, and V. Kapur
Transcriptional Response of Pasteurella multocida to Defined Iron Sources

J. Bacteriol., December 1, 2002; 184(23): 6714 - 6720.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




The American Journal of Pathology

[HOME](#)

A. Dash, I. P. Maine, S. Varambally, R. Shen, A. M. Chinnaiyan, and M. A. Rubin
Changes in Differential Gene Expression because of Warm Ischemia Time of Radical Prostatectomy Specimens

Am. J. Pathol., November 1, 2002; 161(5): 1743 - 1748.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

Y.-Y. E. Chuang, Y. Chen, Gadisetti, V. R. Chandramouli, J. A. Cook, D. Coffin, M.-H. Tsai, W. DeGraff, H. Yan, S. Zhao, A. Russo, E. T. Liu, and J. B. Mitchell

Gene Expression after Treatment with Hydrogen Peroxide, Menadione, or t-Butyl Hydroperoxide in Breast Cancer Cells

Cancer Res., November 1, 2002; 62(21): 6246 - 6254.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[HOME](#)

W.-K. Hofmann, S. de Vos, M. Komor, D. Hoelzer, W. Wachsman, and H. P. Koeffler

Characterization of gene expression of CD34+ cells from normal and myelodysplastic bone marrow

Blood, November 15, 2002; 100(10): 3553 - 3560.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



MOLECULAR ENDOCRINOLOGY

[HOME](#)

R. A. Miller, Y. Chang, A. T. Galecki, K. Al-Regaiey, J. J. Kopchick, and A. Bartke

Gene Expression Patterns in Calorically Restricted Mice: Partial Overlap with Long-Lived Mutant Mice

Mol. Endocrinol., November 1, 2002; 16(11): 2657 - 2666.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



the Journals of gerontology BIOLOGICAL SCIENCES AND MEDICAL SCIENCES

[HOME](#)

E. Wang, C. Lacelle, S. Xu, X. Zhao, and M. Hou

Designer Microarrays: From Soup To Nuts

J. Gerontol. A Biol. Sci. Med. Sci., November 1, 2002; 57 (11): B400 - 405.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online


[HOME](#)

M. Bernaudin, Y. Tang, M. Reilly, E. Petit, and F. R. Sharp

Brain Genomic Response following Hypoxia and Re-oxygenation in the Neonatal Rat. IDENTIFICATION OF GENES THAT MIGHT CONTRIBUTE TO HYPOXIA-INDUCED ISCHEMIC TOLERANCE

J. Biol. Chem., October 18, 2002; 277(42): 39728 - 39738.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[HOME](#)

R. M. Williams, M. Primig, B. K. Washburn, E. A. Winzeler, M. Bellis, C. Sarrauste de Menthier, R. W. Davis, and R. E. Esposito

The Ume6 regulon coordinates metabolic and meiotic gene expression in yeast

PNAS, October 15, 2002; 99(21): 13431 - 13436.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation Research

[HOME](#)

S. A. Cook and A. Rosenzweig


DNA Microarrays: Implications for Cardiovascular Medicine

Circ. Res., October 4, 2002; 91(7): 559 - 564.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Proceedings of the National Academy of Sciences


[HOME](#)



J. P. Brody, B. A. Williams, B. J. Wold, and S. R. Quake
Significance and statistical errors in the analysis of DNA microarray data

PNAS, October 1, 2002; 99(20): 12975 - 12978.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

H. Y. Chang, J.-T. Chi, S. Dudoit, C. Bondre, M. van de Rijn, D. Botstein, and P. O. Brown

Diversity, topographic differentiation, and positional memory in human fibroblasts

PNAS, October 1, 2002; 99(20): 12877 - 12882.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[▶ HOME](#)

M. D. W. Piper, P. Daran-Lapujade, C. Bro, B. Regenberg, S. Knudsen, J. Nielsen, and J. T. Pronk

Reproducibility of Oligonucleotide Microarray Transcriptome Analyses. AN INTERLABORATORY COMPARISON USING CHEMOSTAT CULTURES OF SACCHAROMYCES CEREVISIAE

J. Biol. Chem., October 4, 2002; 277(40): 37001 - 37008.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[▶ HOME](#)

F. Cong, J. Tang, B. J. Hwang, B. Q. Vuong, G. Chu, and S. P. Goff

Interaction between UV-damaged DNA Binding Activity Proteins and the c-Abl Tyrosine Kinase

J. Biol. Chem., September 20, 2002; 277(38): 34870 - 34878.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[▶ HOME](#)

A. Weber and K. Jung

Profiling Early Osmostress-Dependent Gene Expression in Escherichia coli Using DNA Macroarrays

J. Bacteriol., October 1, 2002; 184(19): 5502 - 5507.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Human Molecular Genetics

[▶ HOME](#)

A. D. Strand, J. M. Olson, and C. Kooperberg

Estimating the statistical significance of gene



expression changes observed with oligonucleotide arrays

Hum. Mol. Genet., September 15, 2002; 11(19): 2207 - 2221.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[HOME](#)

C. Kooperberg, S. Sipione, M. LeBlanc, A. D. Strand, E. Cattaneo, and J. M. Olson

Evaluating test statistics to select interesting genes in microarray experiments

Hum. Mol. Genet., September 15, 2002; 11(19): 2223 - 2232.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Physiological Genomics

[HOME](#)

M. S. Halfon and A. M. Michelson

Exploring genetic regulatory networks in metazoan development: methods and models

Physiol Genomics, September 3, 2002; 10(3): 131 - 143.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



blood

[HOME](#)

J. O. Claudio, E. Masih-Khan, H. Tang, J. Goncalves, M. Voralia, Z. H. Li, V. Nadeem, E. Cukerman, O. Francisco-Pabalan, C. C. Liew, J. R. Woodgett, and A. K. Stewart

A molecular compendium of genes expressed in multiple myeloma

Blood, September 15, 2002; 100(6): 2175 - 2186.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Am. J. Physiol. Cell Physiol.

[HOME](#)

K. Rouger, M. Le Cunff, M. Steenman, M.-C. Potier, N. Gibelin, C. A. Dechesne, and J. J. Leger

Global/temporal gene expression in diaphragm and hindlimb muscles of dystrophin-deficient (mdx) mice

Am J Physiol Cell Physiol, September 1, 2002; 283(3): C773 - 784.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Human Molecular Genetics

[HOME](#)

X. L. Xu, J. M. Olson, and L. P. Zhao

A regression-based method to identify differentially



expressed genes in microarray time course studies and its application in an inducible Huntington's disease transgenic model

Hum. Mol. Genet., August 15, 2002; 11(17): 1977 - 1985.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF **Respiratory Cell and Molecular Biology**

[▶ HOME](#)

N. Kaminski and N. Friedman

Practical Approaches to Analyzing Results of Microarray Experiments

Am. J. Respir. Cell Mol. Biol., August 1, 2002; 27(2): 125 - 132.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[▶ HOME](#)

D. R. Rhodes, T. R. Barrette, M. A. Rubin, D. Ghosh, and A. M. Chinnaiyan

Meta-Analysis of Microarrays: Interstudy Validation of Gene Expression Profiles Reveals Pathway Dysregulation in Prostate Cancer

Cancer Res., August 1, 2002; 62(15): 4427 - 4433.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Clinical Chemistry

[▶ HOME](#)

P. F. Macgregor and J. A. Squire

Application of Microarrays to the Analysis of Gene Expression in Cancer

Clin. Chem., August 1, 2002; 48(8): 1170 - 1177.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[▶ HOME](#)

C. Schoch, A. Kohlmann, S. Schnittger, B. Brors, M. Dugas, S. Mergenthaler, W. Kern, W. Hiddemann, R. Eils, and T. Haferlach

Acute myeloid leukemias with reciprocal rearrangements can be distinguished by specific gene expression profiles

PNAS, July 23, 2002; 99(15): 10008 - 10013.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Journal of Applied Physiology

[▶ HOME](#)

B. S. Tseng, P. Zhao, J. S. Pattison, S. E. Gordon, J. A. Granchelli, R. W. Madsen, L. C. Folk, E. P. Hoffman, and F.




W. Booth

Regenerated mdx mouse skeletal muscle shows differential mRNA expression

J Appl Physiol, August 1, 2002; 93(2): 537 - 545.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[▶ HOME](#)

C. Cicala, J. Arthos, S. M. Selig, G. Dennis Jr., D. A. Hosack, D. Van Ryk, M. L. Spangler, T. D. Steenbeke, P. Khazanie, N. Gupta, J. Yang, M. Daucher, R. A. Lempicki, and A. S. Fauci

HIV envelope induces a cascade of cell signals in non-proliferating target cells that favor virus replication

PNAS, July 9, 2002; 99(14): 9380 - 9385.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of the American Society of Nephrology

[▶ HOME](#)

M. Kretzler, C. D. Cohen, P. Doran, A. Henger, S. Madden, E. F. Grone, P. J. Nelson, D. Schlondorff, and H.-J. Grone

Repuncturing the Renal Biopsy: Strategies for Molecular Diagnosis in Nephrology

J. Am. Soc. Nephrol., July 1, 2002; 13(7): 1961 - 1972.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences


[▶ HOME](#)

G. W. Birrell, J. A. Brown, H. I. Wu, G. Giaever, A. M. Chu, R. W. Davis, and J. M. Brown

Transcriptional response of *Saccharomyces cerevisiae* to DNA-damaging agents does not identify the genes that protect against these agents

PNAS, June 25, 2002; 99(13): 8778 - 8783.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[▶ HOME](#)

M. A. Fisher, B. B. Plikaytis, and T. M. Shinnick

Microarray Analysis of the *Mycobacterium tuberculosis* Transcriptional Response to the Acidic Conditions Found in Phagosomes

J. Bacteriol., July 15, 2002; 184(14): 4025 - 4032.


[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

ANNALS of the New York Academy of Sciences

[▶ HOME](#)

K. GOHIL and L. PACKER

Bioflavonoid-Rich Botanical Extracts Show Antioxidant



and Gene Regulatory Activity

Ann. N.Y. Acad. Sci., May 1, 2002; 957(1): 70 - 77.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

Y. Xu, F. M. Selaru, J. Yin, T. T. Zou, V. Shustova, Y. Mori, F. Sato, T. C. Liu, A. Olaru, S. Wang, M. C. Kimos, K. Perry, K. Desai, B. D. Greenwald, M. J. Krasna, D. Shibata, J. M. Abraham, and S. J. Meltzer

Artificial Neural Networks and Gene Filtering Distinguish Between Global Gene Expression Profiles of Barrett's Esophagus and Esophageal Cancer

Cancer Res., June 1, 2002; 62(12): 3493 - 3497.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Bacteriology

[HOME](#)

M. L. Paustian, B. J. May, and V. Kapur

Transcriptional Response of *Pasteurella multocida* to Nutrient Limitation

J. Bacteriol., July 1, 2002; 184(13): 3734 - 3739.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF DENTAL RESEARCH

[HOME](#)

X. Han and S. Amar

Identification of Genes Differentially Expressed in Cultured Human Periodontal Ligament Fibroblasts vs. Human Gingival Fibroblasts by DNA Microarray Analysis

J. Dent. Res., June 1, 2002; 81(6): 399 - 405.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



EUKARYOTIC CELL

[HOME](#)

M. D. Cleary, U. Singh, I. J. Blader, J. L. Brewer, and J. C. Boothroyd

Toxoplasma gondii Asexual Development: Identification of Developmentally Regulated Genes and Distinct Patterns of Gene Expression

Eukaryot. Cell, June 1, 2002; 1(3): 329 - 340.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

MOLECULAR ENDOCRINOLOGY

[HOME](#)

A. Flores-Morales, H. Gullberg, L. Fernandez, N. Stahlberg, N. H. Lee, B. Vennstrom, and G. Norstedt



Patterns of Liver Gene Expression Governed by TR {beta}

Mol. Endocrinol., June 1, 2002; 16(6): 1257 - 1268.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Cancer Research

[HOME](#)

H. Shi, P. S. Yan, C.-M. Chen, F. Rahmatpanah, C. Lofton-Day, C. W. Caldwell, and T. H.-M. Huang

Expressed CpG Island Sequence Tag Microarray for Dual Screening of DNA Hypermethylation and Gene Silencing in Cancer Cells

Cancer Res., June 1, 2002; 62(11): 3214 - 3220.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Endocrinology

[HOME](#)

J. A. Mong, C. Krebs, and D. W. Pfaff

Perspective: Microarrays and Differential Display PCR--Tools for Studying Transcript Levels of Genes in Neuroendocrine Systems

Endocrinology, June 1, 2002; 143(6): 2002 - 2006.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

R. Tibshirani, T. Hastie, B. Narasimhan, and G. Chu Diagnosis of multiple cancer types by shrunken centroids of gene expression

PNAS, May 14, 2002; 99(10): 6567 - 6572.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Molecular & Cellular PROTEOMICS

[HOME](#)

G. Chen, T. G. Gharib, C.-C. Huang, J. M. G. Taylor, D. E. Misek, S. L. R. Kardia, T. J. Giordano, M. D. Iannettoni, M. B. Orringer, S. M. Hanash, and D. G. Beer

Discordant Protein and mRNA Expression in Lung Adenocarcinomas

Mol. Cell. Proteomics, April 1, 2002; 1(4): 304 - 313.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

A. I. Brooks, C. S. Stein, S. M. Hughes, J. Heth, P. M. McCray Jr., S. L. Sauter, J. C. Johnston, D. A. Cory-Slechta, H. J. Federoff, and B. L. Davidson

From the Cover: Functional correction of established

**central nervous system deficits in an animal model of
lysosomal storage disease with feline
immunodeficiency virus-based vectors**

PNAS, April 30, 2002; 99(9): 6216 - 6221.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



AMERICAN JOURNAL OF Respiratory Cell and Molecular Biology [▶ HOME](#)

I. G. Luzina, S. P. Atamas, R. Wise, F. M. Wigley, H. Q. Xiao,
and B. White

**Gene Expression in Bronchoalveolar Lavage Cells from
Scleroderma Patients**

Am. J. Respir. Cell Mol. Biol., May 1, 2002; 26(5): 549 - 557.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




JBC Online [▶ HOME](#)

S. R. Schwarze, S. E. DePrimo, L. M. Grabert, V. X. Fu, J. D.
Brooks, and D. F. Jarrard

**Novel Pathways Associated with Bypassing Cellular
Senescence in Human Prostate Epithelial Cells**

J. Biol. Chem., April 26, 2002; 277(17): 14877 - 14883.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences [▶ HOME](#)

Q. Zhao, A. Kho, A. M. Kenney, D.-i. Yuk, I. Kohane, and D.
H. Rowitch

**Identification of genes expressed with temporal-
spatial restriction to developing cerebellar neuron
precursors by a functional genomic approach**

PNAS, April 16, 2002; 99(8): 5704 - 5709.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JOURNAL OF CLINICAL ONCOLOGY [▶ HOME](#)

S. Ramaswamy and T. R. Golub

DNA Microarrays in Clinical Oncology

J. Clin. Oncol., April 1, 2002; 20(7): 1932 - 1941.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Nucleic Acids Research [▶ HOME](#)

Y. Cohen, M. Dardalhon, and D. Aeverbeck

**Homologous recombination is essential for RAD51 up-
regulation in Saccharomyces cerevisiae following DNA
crosslinking damage**

Nucleic Acids Res., March 1, 2002; 30(5): 1224 - 1232.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

K.-H. Pan, C.-J. Lih, and S. N. Cohen

Analysis of DNA microarrays using algorithms that employ rule-based expert knowledge

PNAS, February 19, 2002; 99(4): 2118 - 2123.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH

[HOME](#)

K. Clement, N. Viguerie, M. Diehn, A. Alizadeh, P. Barbe, C. Thalamas, J. D. Storey, P. O. Brown, G. S. Barsh, and D. Langin

In Vivo Regulation of Human Skeletal Muscle Gene Expression by Thyroid Hormone

Genome Res., February 1, 2002; 12(2): 281 - 291.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Human Molecular Genetics

[HOME](#)

J. D. Porter, S. Khanna, H. J. Kaminski, J. S. Rao, A. P. Merriam, C. R. Richmonds, P. Leahy, J. Li, W. Guo, and F. H. Andrade

A chronic inflammatory response dominates the skeletal muscle molecular signature in dystrophin-deficient mdx mice

Hum. Mol. Genet., February 1, 2002; 11(3): 263 - 272.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



GENOME RESEARCH


[HOME](#)

J. Theilhaber, T. Connolly, S. Roman-Roman, S. Bushnell, A. Jackson, K. Call, T. Garcia, and R. Baron

Finding Genes in the C2C12 Osteogenic Pathway by k-Nearest-Neighbor Classification of Expression Data

Genome Res., January 1, 2002; 12(1): 165 - 176.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Nutrition

[HOME](#)

J. B. Moore, R. K. Blanchard, W. T. McCormack, and R. J. Cousins

cDNA Array Analysis Identifies Thymic LCK as Upregulated in Moderate Murine Zinc Deficiency before T-Lymphocyte Population Changes

J. Nutr., December 1, 2001; 131(12): 3189 - 3196.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

R. K. Blanchard, J. B. Moore, C. L. Green, and R. J. Cousins

Inaugural Article: Modulation of intestinal gene expression by dietary zinc status: Effectiveness of cDNA arrays for expression profiling of a single nutrient deficiency

PNAS, November 20, 2001; 98(24): 13507 - 13513.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



JBC Online

[HOME](#)

B. S. Finlin, C.-L. Gau, G. A. Murphy, H. Shao, T. Kimel, R. S. Seitz, Y.-F. Chiu, D. Botstein, P. O. Brown, C. J. Der, F. Tamanoi, D. A. Andres, and C. M. Perou

RERG Is a Novel ras-related, Estrogen-regulated and Growth-inhibitory Gene in Breast Cancer

J. Biol. Chem., November 9, 2001; 276(45): 42259 - 42267.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

N. N. Khodarev, J. O. Park, J. Yu, N. Gupta, E. Nodzenski, B. Roizman, and R. R. Weichselbaum

Dose-dependent and independent temporal patterns of gene responses to ionizing radiation in normal and tumor cells and tumor xenografts

PNAS, October 23, 2001; 98(22): 12665 - 12670.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

G. W. Birrell, G. Giaever, A. M. Chu, R. W. Davis, and J. M. Brown

A genome-wide screen in *Saccharomyces cerevisiae* for genes affecting UV radiation sensitivity

PNAS, October 23, 2001; 98(22): 12608 - 12613.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

J. D. Porter, S. Khanna, H. J. Kaminski, J. S. Rao, A. P. Merriam, C. R. Richmonds, P. Leahy, J. Li, and F. H. Andrade

Extraocular muscle is defined by a fundamentally distinct gene expression profile

PNAS, October 9, 2001; 98(21): 12062 - 12067.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

T. Sorlie, C. M. Perou, R. Tibshirani, T. Aas, S. Geisler, H. Johnsen, T. Hastie, M. B. Eisen, M. van de Rijn, S. S. Jeffrey, T. Thorsen, H. Quist, J. C. Matese, P. O. Brown, D. Botstein, P. E. Lonning, and A.-L. Borresen-Dale

Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications

PNAS, September 11, 2001; 98(19): 10869 - 10874.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Science


[HOME](#)

E. E. M. Furlong, E. C. Andersen, B. Null, K. P. White, and M. P. Scott

Patterns of Gene Expression During Drosophila Mesoderm Development

Science, August 31, 2001; 293(5535): 1629 - 1633.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Circulation Research

[HOME](#)

J. S. Rao and M. Bond

Microarrays : Managing the Data Deluge

Circ. Res., June 22, 2001; 88(12): 1226 - 1227.

[\[Full Text\]](#) [\[PDF\]](#)



Proceedings of the National Academy of Sciences

[HOME](#)

C. S. Detweiler, D. B. Cunanan, and S. Falkow

Host microarray analysis reveals a role for the Salmonella response regulator phoP in human macrophage cell death

PNAS, May 8, 2001; 98(10): 5850 - 5855.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

This Article

▸ [Abstract](#) FREE

▸ [Full Text \(PDF\)](#)

▸ [An erratum has been published](#)

▸ [An erratum has been published](#)

▸ [Alert me when this article is cited](#)

▸

Alert me if a correction is posted

‣ **Citation Map**

Services

‣ **Similar articles in this journal**

‣ **Similar articles in ISI Web of Science**

‣ **Similar articles in PubMed**

‣ **Alert me to new issues of the journal**

‣ **Add to My File Cabinet**

‣ **Download to citation manager**

‣ **Search for citing articles in:
ISI Web of Science (1076)**

‣ **Request Copyright Permission**

PubMed

‣ **PubMed Citation**


‣ **Articles by Tusher, V. G.**

‣ **Articles by Chu, G.**

[Current Issue](#) | [Archives](#) | [Online Submission](#) | [Info for Authors](#) | [Editorial Board](#) | [About](#)
[Subscribe](#) | [Advertise](#) | [Contact](#) | [Site Map](#)

[Copyright © 2001 by the National Academy of Sciences](#)

ATTACHMENT 2



SAM: Significance Analysis of Microarrays

**Supervised learning software
for genomic expression data mining**

News

NEW New release 2.20, Oct 4, 2005. SAM now provides sample size assessment-estimates of FDR, FNR, type I error and power for different sample sizes.

"A simple method for assessing sample sizes in microarray experiments" (pdf) .

NEW Major New Release: Version 2.0. June 6, 2005. Now version 2.11----- Aug 24, 2005. All users should upgrade to this version. SAM now handles time course data, does non-parametric tests and pattern discovery, It also reports local false discovery rates and miss rates.

A discussion and announcement group for all SAM-related discussions and announcements has been created. See <http://groups.yahoo.com/group/sam-software>.

Features

- Developed at Stanford University Labs: based on recent paper of Tusher, Tibshirani and Chu (2001):
"Significance analysis of microarrays applied to the ionizing radiation response" (ps file). (pdf version). PNAS 2001 98: 5116-5121, (Apr 24).
"Raw data"
- Correlates gene expression data to a wide variety of clinical parameters including treatment, diagnosis categories, survival time and time trends
- Provides estimate of False Discovery Rate for multiple testing
- Convenient Excel Add-in
- Works with data from both cDNA and oligo microarrays. Can also be applied to protein expression data and SNP chip data.
- Patent Pending for SAM technology

- SAM uses the FDR and q-value method presented in Storey (2002) A direct approach to false discovery rates. J. Roy. Stat. Soc. Ser. B, 64:479-498;

Local false discovery rates proposed in Efron, B., Tibshirani, R., Storey, JD, and Tusher, V. (2001). Empirical Bayes Analysis of a Microarray Experiment, JASA, 96, 1151-1160 and Efron and Tibshirani, "Microarrays, Empirical Bayes Methods, and False Discovery Rates" Genet. Epidemiol. 2002 Jun;23(1):70-86;

and *Miss rates*--- Jon Taylor, Rob Tibshirani and Brad Efron. The "Miss rate" for the analysis of gene expression data; Biostatistics 2005 6 (1):111-117.

- [List of features](#)
- [R package samr](#)
- [Sample screens](#)
- [Frequently Asked Questions](#)
- ["Samster" tool](#)
- [Euan Ashley's heatmap builder](#)
- Related links: [PAM package for microarray classification](#);
[CGH-Miner package for CGH data](#);
[PPC package for protein mass spec classification](#)
[Superpc package for microarray prediction](#);

Obtaining SAM


- Academic users can download SAM by going directly to the [registration page](#). **Please note that this is the full version!**
- Non academic users should first register via the [registration page](#). An evaluation version (limited to 500 genes) can be downloaded directly from that page.

If you are a commercial user and wish to obtain a complete version of SAM, proceed to the [SAM resource](#) at the [Office of Technology and Licensing](#). The SAM contact is [Sara Nakashima](#)

(sara.nakashima@stanford.edu) at the Office of Technology and Licensing,
Phone: (650) 725-9117.

Please do not contact Sara Nakashima about downloading, technical questions etc. All she handles is commercial licensing!

- Returning users (those who have already registered) who want to download the software again can proceed directly to the Academic Download Page or the Non-Academic Download Page. You will need the registration information that you received via email.



ATTACHMENT 3

ACADEMY REPORTS ONLINE!

Signup for PNAS Online eTocs

Get notified by email when new content goes online

[Info for Authors](#) | [Editorial Board](#) | [About](#) | [Subscribe](#) | [Advertise](#) | [Contact](#) | [Site Map](#)

PNAS

Proceedings of the National Academy of Sciences of the United States of America

[Current Issue](#)

[Archives](#)

[Online Submission](#)

GO

[advanced search >>](#)

**Published online before print September 2, 2005,
10.1073/pnas.0506230102**

PNAS | September 20, 2005 | vol. 102 | no. 38 | 13550-13555

OPEN ACCESS ARTICLE

**From The Cover
GENETICS**

An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival

**Lance D. Miller^{*,†}, Johanna Smeds[‡],
Joshy George^{*}, Vinsensius B. Vega^{*},
Liza Vergara^{*}, Alexander Ploner[§],
Yudi Pawitan[§], Per Hall[§], Sigrid Klaar[‡],
Edison T. Liu^{*,†} and Jonas Bergh[‡]**

^{*} Genome Institute of Singapore, 60 Biopolis Street,
#02-01, Singapore 138672; [†] Department of
Oncology and Pathology, Radiumhemmet,
Karolinska Institute and Hospital, S-17176

Stockholm, Sweden; and [§] Department of Medical Epidemiology and Biostatistics,

This Article

Free via Open Access: OA

- ▶ [OA Abstract](#)
- ▶ [Figures Only](#)
- ▶ [Full Text \(PDF\)](#)
- ▶ [Supporting Information](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)
- ▶ [Citation Map](#)

Services

- ▶ [Related articles in PNAS](#)
- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Add to My File Cabinet](#)
- ▶ [Download to citation manager](#)
- ▶ [Request Copyright Permission](#)

Google Scholar

- ▶ [Articles by Miller, L. D.](#)
- ▶ [Articles by Bergh, J.](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Miller, L. D.](#)
- ▶ [Articles by Bergh, J.](#)

Karolinska Institutet, 17177 Stockholm, Sweden

Communicated by Raymond L. White, University of California at San Francisco, Emeryville, CA, July 26, 2005 (received for review February 20, 2005)

► Abstract

Perturbations of the p53 pathway are associated with more aggressive and therapeutically refractory tumors. However, molecular assessment of p53 status, by using sequence analysis and immunohistochemistry, are incomplete assessors of p53 functional effects. We posited that the transcriptional fingerprint is a more definitive downstream indicator of p53 function. Herein, we analyzed transcript profiles of 251 p53-sequenced primary breast tumors and identified a clinically embedded 32-gene expression signature that distinguishes p53-mutant and wild-type tumors of different histologies and outperforms sequence-based assessments of p53 in predicting prognosis and therapeutic response. Moreover, the p53 signature identified a subset of aggressive tumors absent of sequence mutations in p53 yet exhibiting expression characteristics consistent with p53 deficiency because of attenuated p53 transcript levels. Our results show the primary importance of p53 functional status in predicting clinical breast cancer behavior.

microarray | expression analysis | tumor profiling | class prediction

▲ Top
• Abstract
▼ Methods
▼ Results
▼ Discussion
▼ References

The p53 tumor suppressor is a critical regulator of tissue homeostasis, and its inactivation at the gene or protein level confers cellular properties conducive for oncogenesis and cancer progression. Mutations in p53 occur in >50% of human cancers ([1](#), [2](#)), and the mutational status of p53 is prognostic in many malignancies ([3](#)). In breast cancer, p53 mutations are associated with worse overall and disease-free survival, independent of other risk factors ([4](#)), and have been implicated in resistance to anticancer therapies ([5-11](#)). These observations, however, have been

inconsistent ([12](#), [13](#)), owing, in part, to the variable accuracy of the methods to ascertain p53 status, variation in disease severity attributable to the different forms of p53 mutation, and studies of insufficient size ([8](#), [11](#), [14](#)). Further confounding the association between p53 status and patient risk is the growing number of alternative molecular mechanisms (e.g., MDM2) that compromise p53 function.

In this study, we explored the possibility that a gene-expression signature, derived from differences between p53 mutant (mt) and wild-type (wt) breast tumors, could provide a more accurate measure of the functional configuration of p53, thereby improving its prognostic utility. Using oligonucleotide microarrays covering >30,000 genes, we analyzed the global transcript levels of 251 primary invasive breast tumors for which we have detailed information on p53 status, as determined by cDNA sequencing ([6](#)) and pursued a validation strategy of intersecting alternative array data sets. We found that, in most cases, tumors with mt and wt p53 can readily be distinguished by their expression profiles and that a 32-gene p53 signature is consistently associated with patient survival in different patient subsets, independent of other risk factors, and is a superior prognostic and predictive indicator, compared with p53 mutation status alone.

► **Methods**

Patients and Specimens. Frozen tissue was collected from 315 consecutively presented primary breast cancers representing 65% of all those resected in Uppsala County, Sweden, from January 1, 1987 to December 31, 1989 ([6](#)). Of these tissues, 251 were comprised predominantly of diseased tissue, were sequenced for p53 ([6](#)), and yielded sufficient RNA for array analysis. Clinicopathological variables measured at diagnosis were obtained from patient records and are described in detail in *Supporting Materials and*

▲ Top
▲ Abstract
• Methods
▼ Results
▼ Discussion
▼ References

Methods, which is published as supporting information on the PNAS web site. This microarray study was approved by the ethical committee at the Karolinska Institute, Stockholm, Sweden.

Expression Profiling. Total RNA was extracted from samples by using RNEasy Mini kit (Qiagen, Hilden, Germany) and evaluated on a 2100 Bioanalyzer (Agilent Technologies). *In vitro* transcription products were prepared from 2-5 μ g of total RNA, hybridized to the Affymetrix U133 A and B arrays and washed and scanned according to the manufacturer's instructions.

Microarray Data Processing. Raw data were normalized by using the global mean method. Probe-set signal values were natural log transformed and scaled by adjusting the mean intensity to a target signal value of log 500. Samples with suboptimal average signal intensities (i.e., scaling factors >3.5) or GAPDH 3'/5' ratios >3.5 were relabeled and rehybridized on new arrays. If visible artifacts were observed, the same cRNA was rehybridized on new chips.

Class Prediction. For gene selection, we fit a linear model to the expression data with expression level as the response and p53 status, estrogen-receptor (ER) status, and grade status as the predictor variables. As an initial filter, we excluded genes with a *P* value for model fit >0.001 and ranked genes in decreasing order of the absolute value of the p53 status coefficient. For class prediction, we evaluated several supervised learning methods, including diagonal linear discriminant analysis (15), *k* nearest neighbors (16), and support vector machines (17), as described in *Supporting Materials and Methods*.

Data Analysis. For all hierarchical cluster analyses, log expression values of each gene were mean centered, and genes and tumors were clustered by using Pearson correlation and average linkage (CLUSTER and TREEVIEW software, <http://rana.lbl.gov/EisenSoftware.htm>).

The Kaplan-Meier estimate was used to compute survival curves, and the P value of the likelihood-ratio test was used to assess statistical significance of the hazard ratios. All patients with contralateral or bilateral cancers were omitted, and patients who died of their cancer 10 years after diagnosis were systematically censored.

For association tests, the χ^2 test was used, unless the number of events was <5 in any category, in which case Fisher's exact test was used.

Cox regression was used to confirm the prognostic significance of the p53 classifier in multivariate analyses. The initial model, comprising all conventional predictors, and p53 mutation status and the p53 signature as competing measures of p53 activity, was simplified by using a stepwise model-selection procedure based on the Akaike information criterion. Remaining predictors were assessed by likelihood-ratio test.

Independent Datasets. The Sørbye *et al.* (18) and Chen *et al.* (19) data and clinical annotations were obtained from the Stanford microarray database by using filtering parameters as described by the authors. The Ma *et al.* (20) "whole tumor" data set was downloaded from the Gene Expression Omnibus with accession no. GSE1379 [NCBI GEO], and each array was mean centered. The van't Veer *et al.* (21) data and survival annotation were accessed through the Rosetta Inpharmatics publications archive. All IMAGE clone IDs or GenBank accession nos. of array probes were mapped to UniGene build no. 167.

► Results

P53 Mutant and WT Tumors Are Molecularly Distinct. Transcript profiles of 251 primary breast tumors were assessed by using Affymetrix U133 oligonucleotide microarrays. Previously, cDNA sequence analysis revealed that 58 of these tumors had p53 mutations

▲	Top
▲	Abstract
▲	Methods
•	Results
▼	Discussion
▼	References

resulting in protein-level changes, whereas the remaining 193 tumors were p53 wt (6). By unsupervised hierarchical cluster analysis, we found that p53 mt and wt tumors are distinguished by pervasive molecular differences. With the top 2,000 most variably expressed genes (selected independent of p53 status), >80% of the p53 mt tumors clustered into one branch and >70% of the p53 wts into the other ($P = 5.6 \times 10^{-13}$; see Fig. 5, which is published as [supporting information](#) on the PNAS web site). Importantly, this separation remained highly significant ($P < 2 \times 10^{-12}$) across a range of gene panels from the top 5,000 genes with highest variance to the top 125 (see Table 1, which is published as [supporting information](#) on the PNAS web site). This separation was most heavily influenced by three predominant gene clusters comprising genes involved in immune response, proliferation, and estrogen response (Fig. 5). Univariate analysis by statistical analysis of microarrays (SAM) (22) identified 6,545 Affymetrix probe sets representing $\approx 5,290$ distinct genes whose expression patterns distinguished p53 mt and wt tumors with a false discovery rate (q value) <1% and d score (modified t statistic) >2.0 (see Table 2, which is published as [supporting information](#) on the PNAS web site), further illuminating the extensive nature of the molecular variation underlying p53 status. Topping the list of genes most highly expressed in p53 mt tumors were those with roles in cell cycle and proliferation, consistent with the observation that wt p53 has a negative regulatory effect on cell-cycle genes. The genes more highly expressed in the p53 wt tumors included uncharacterized genes, signaling molecules and transcription factors, transcriptional targets of p53, and estrogen-inducible genes.

The p53 status was also correlated with two other clinical parameters, ER status and tumor grade (Fig. 5). Within the p53 mt-rich cluster, we observed 89% of ER-negative tumors ($P = 1.9 \times 10^{-10}$), 79% of grade III tumors ($P = 3.8 \times 10^{-11}$), and only 14% of grade I tumors ($P = 2.5 \times 10^{-7}$). The finding that p53 mutant tumors are correlated with ER negativity and grade III status is consistent with previous reports that p53 mutations associate with ER negativity and high tumor grade (23).


A Gene Expression Classifier Predicts p53 Status in Independent Breast and Liver Cancer Data Sets. We considered the possibility that the differential expression observed between p53 mt and wt tumors might, to some extent, reflect changes in the operational configuration of the p53 pathway. We reasoned that some p53 wt tumors would be p53 deficient through mechanisms other than p53 mutation, such as MDM2 amplification or p14/ARF deletion and, thus, possess expression profiles more akin to p53 mt tumors with dysfunctional p53. To explore this possibility, we fitted a multivariate linear regression model (i.e., linear modelfit) (24) that allowed us to rank genes by their correlation with p53 status, while controlling for histologic grade and ER status. As a result, many cell-cycle genes correlated with p53 status by univariate analysis were no longer well associated (see Fig. 6, which is published as supporting information on the PNAS web site), suggesting that the transcriptional profiles of most cell-cycle genes are more related to histologic grade than to p53 status.

For class discrimination, we evaluated several linear learning methods including: diagonal linear discriminant analysis (DLDA) (15), *k*-nearest neighbors (*k*NN) (16), and support vector machines (SVM) (17). In each case, the optimal gene classifier was obtained by leave-one-out cross validation, where the linear model-fit procedure was iteratively applied to all samples minus the left-out sample. The resulting prediction accuracies were highly similar, ranging from 84.9% to 85.7% (see *Supporting Materials and Methods*). Interestingly, 20 tumors were consistently "misclassified" by all three methods (8 wt and 12 mt), indicating a surprising degree of concordance among misclassified tumors. DLDA showed the highest sensitivity for detecting p53 mutants (i.e., 79% sensitivity compared with 53% for both *k*NN and SVM) and was therefore selected for further analysis. By DLDA, the optimal classifier was comprised of 32 genes, whereby 26 of the wt tumors were misclassified as mutant-like, and 12 mutants were misclassified as wt-like (Fig. 1A).

To evaluate the performance of the classifier genes (referred to hereafter as the p53

signature genes) as a clinical discriminator of p53 status, we accessed two publicly available cDNA microarray data sets where p53 mutational status was known: a breast cancer study by Sørli *et al.* (18) and a liver cancer study by Chen *et al.* (19). In the Sørli data set, 69 breast tumors had been sequenced for p53 mutations. Of our p53 signature genes, 28 mapped to established UniGene IDs, and more than half of these 28 genes were represented on the Sørli *et al.* microarray. However, only nine were found to correspond to cDNA probes having expression measurements present in >50% of tumors, where the tumors possessed measurements for >50% of genes (resulting in a subset of 44 well sampled tumors). Because the classification rules could not be directly applied, we used this 9-gene subset of the p53 signature to hierarchically cluster the tumors in an unsupervised manner. Fig. 1B shows a significant separation of p53 mt and wt tumors: 77% of mutants clustered into one branch, and 77% of wts clustered into the other ($P = 0.0003$). By Monte Carlo simulations, we estimated the probability that a randomly selected nine-gene subset could cluster the samples with equivalent or better significance was $P = 0.008$, thus reaffirming the robust discriminative power of the p53 signature genes.

In the Chen *et al.* liver cancer data set (38), p53 protein levels had been ascertained by immunohistochemistry (IHC). Eight of our signature genes could be mapped to all 59 tumors assayed for p53, with each gene having data present in >90% of all tumors and where each tumor contained data for >50% of the genes. We observed that even this eight-gene subset was able to cluster the liver cancers into two primary clusters significantly correlated with p53 levels: 87% of the IHC-positive (predicted mts) in one cluster, and 61% of the predicted wts in the other ($P = 0.00035$) (Fig. 1C). Again, the probability of this clustering occurring by random chance was $P = 0.009$ by Monte Carlo P value estimation. Taken together, these observations suggest that the genes comprising the p53 signature are robust in their ability to classify not only breast tumors but also liver cancers according to their p53 mutational status and, therefore, may have generalizable utility in predicting p53 status in a range of cancer types.



View larger version

(93K):


[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 1. The p53 signature is associated with p53 status in independent data sets. Clustergrams are oriented as outlined in Fig. 5. (A) Expression profiles of the Uppsala tumors segregated by the 32-gene signature. Unigene symbols and GenBank IDs are listed to the right. (B) P53 mt and wt breast tumors from Sørle *et al.* (18) were clustered by using a nine-gene subset of the p53 signature. (C) P53 mt and wt liver tumors (predicted by immunohistochemistry) from Chen *et al.* (19) were clustered by using an eight-gene subset of the p53 signature. Green dendrogram branches denote tumors with the wt-like configuration; red branches indicate those with mt-like profiles. Probe IMAGE clone IDs from the original studies are listed. Black bars denote mt p53 status.

Transcript Analysis of p53 Pathway Genes Corroborates Tumor

Classifications. We hypothesized that the p53 expression signature may better reflect the relative intactness of p53 function in the tumor than sequence mutation status alone, implying that p53 sequence-wt tumors "misclassified" as mt-like may, in fact, be p53 deficient by other means. First, we considered the possibility that p53 deficiency could result from reduced p53 transcript levels. We compared the transcript levels of p53 among the different tumor classes (Fig. 2). We observed that the overall expression level of p53 was significantly reduced in the 26 wt tumors with mt-like signatures (referred to henceforth as the "26 mt-like" tumors), compared with the remaining 167 wt tumors classified as wt-like ($P = 1.8 \times 10^{-4}$), strongly suggesting that reduced p53 transcripts can result in biological consequences *in vivo*.



View larger version

Fig. 2. Transcript levels of p53 and its transcriptional targets are consistent with classification results. Expression levels of p53-pathway-relevant genes were examined in different tumor subgroups. The

(37K):
[\[in this window\]](#)
[\[in a new window\]](#)

four tumor subgroups are defined as follows: (i) p53 mt tumors classified as mt-like ($n = 46$), (ii) p53 wt tumors classified as mt-like ($n = 26$), (iii) p53 wt tumors classified as wt-like ($n = 167$), and (iv) p53 mt tumors classified as wt-like ($n = 12$). Differences in transcript levels were determined by t test and are shown in a summary table to the right; P values >0.05 are shown in gray.

We further hypothesized that known transcriptional targets of p53 would show altered transcription in p53-deficient tumors. Indeed, a number of p53 target genes demonstrated expression patterns consistent with a mutant p53 status (Fig. 2). The *TP53*-inducible genes *TP53INP1*, *SEMA3B*, *PMAIP1* (*NOXA*), *FDXR*, *CCNG1*, and *LRDD*, which all contain functional p53-binding sites in their promoters, showed significantly lower expression in the 26 mt-like tumors, compared with the other wt (all at $P < 0.05$). In a consistent manner, all but one of these genes were also significantly reduced in the p53 mt tumors, compared with all wt tumors. Furthermore, in all but two cases, these genes showed significantly higher expression in the set of 12 sequence-mt tumors classified as wt-like when compared with the other mts, suggesting that the p53 mutations in these 12 tumors may have a more benign effect, with respect to p53 functionality. *CHEK1* and *CHEK2* are both upstream effectors of p53 function known to be transcriptionally repressed by p53. Significantly, their mRNA levels were elevated in both the p53 mt and p53 mt-like classes. Again, the 12 mts classified as wt-like showed a reversed pattern, i.e., displaying significantly lower expression of these genes, compared with the other 46 p53 mutants. Together, these observations suggest that the "misclassified" tumors more correctly reflect the active/inactive status of the p53 pathway and are consistent with the notion that reduced p53 levels in breast tumors result in downstream transcriptional changes similar to those found in p53 mutations.

Of note, the canonical marker of p53 activity, *CDKN1A* (p21/WAF1), was only

moderately higher in p53 wt tumors, compared with those with sequence mutations ($P = 0.02$), and not significantly lower in the 26 mt-like tumors, compared with the other wts ($P = 0.09$; data not shown). Furthermore, the known p53-inducible genes *PERP*, *BAX*, and *SFN* (14-3-3 sigma) were, paradoxically, all expressed at higher levels in the p53 mutants and the 26 mt-like tumors rather than the expected lower levels (Fig. 2). These observations may reflect cross-talk among different transcriptional regulators in the consensus of primary tissues, as compared with dynamic changes in single cell lines. For example, the p53 target genes *p21* and *BAX* are also directly regulated by the breast cancer oncogene, *c-Myc*, in a manner independent of, and antagonistic to, p53 (25, 26). The regulation of p53 target genes by alternative transcriptional modifiers acting independently of p53 or in the context of p53 deficiency (e.g., *PERP*, *BAX*, and *SFN*) may have implications for p53 tumor-suppressor activity.

We next asked whether the mutational spectrum of p53 in our tumors could explain the different functional consequences, as measured by the expression profiles. Of the 46 p53 mt tumors correctly classified as mts, 43% (20 of 46) possessed "severe" mutations, defined as insertions ($n = 2$), deletions ($n = 11$), and stop codons ($n = 7$) resulting in frame shifts and truncations, whereas in the 12 p53 mutants classified as wt-like by the expression signature, only 1 contained a severe mutation, a 3-bp insertion in the DNA-binding domain, resulting in the in-frame addition of a glycine residue. Notably, this difference was statistically significant at $P = 0.02$. Using the IARC TP53 mutation database (ITMD) (27), we cross-compared the missense point mutations (mpms) in each tumor group with the ITMD's index of 418 mutants previously analyzed for dominant-negative function. Only 1 of the 11 mpms among the 12 wt-like mutants had been demonstrated previously to have dominant-negative activity, compared with 12 of 27 within the mt-like group ($P = 0.039$). Together, these data suggest that, at the sequence level, the 12 p53 mutants classified as wt-like may, in fact, represent p53 mutant forms that have less biological effect.

The p53 Signature Predicts Outcome Better Than p53 Mutation Status Alone.

We next asked whether the p53 signature could predict disease-specific survival in the patients of the Uppsala cohort. The classifier separated patients into low and high risk groups with a much higher statistical significance than the sequence-based p53 status alone ($P = 0.0006$ versus $P = 0.01$, respectively) (Fig. 3 *A* and *B*). More interestingly, when the classifier was tested on the subset of women with wt p53 by sequence, we again observed a significant separation of patients by survival ($P = 0.02$; Fig. 3*C*), indicating that women with p53 sequence-wt tumors, yet exhibiting the mt-like expression signature, have a greater likelihood of dying from their cancer. Fig. 3*D* shows that the survival curve for this tumor type is highly similar to that of p53 mt tumors classified as mt-like (blue and green curves, respectively), whereas the 12 individuals with p53 mt tumors classified as wt-like do not have significantly unique outcomes.




Fig. 3. The p53 classifier has greater prognostic significance than p53 mutation status alone. Kaplan-Meier survival plots for disease-specific survival are shown for patients classified according to p53 mutation status (*A* and *E*), the p53 classifier (*B*, *C*, and *F*), or both (*D*). All patients were assessed in *A*, *B*, and *D*. Only the patients with p53 wt tumors were assessed in *C*. Sixty-seven ER⁺, hormone-treated (TAM) patients were assessed in *E* and

View larger version *F*.

(41K):

[in this window]

[in a new window]


To further test the clinical utility of the p53 signature, we analyzed its prognostic performance on therapy-specific treatment groups. In a subpopulation of the Uppsala cohort consisting of 67 ER⁺ patients who received only adjuvant hormonal therapy

after surgery, the signature was a significant predictor of disease-specific survival ($P = 0.05$), whereas p53 mutation status alone was not ($P = 0.4$) (Fig. 3 *E* and *F*).

Importantly, by multivariate Cox regression analysis, the p53 classifier remained significantly associated with survival in the hormone-treated group ($P = 0.02$), the complete cohort ($P = 0.02$), and the p53 wt group ($P = 0.002$), even when controlling for the classical predictors (ER and progesterone receptor) and prognostic factors (lymph node status, Elston grade, tumor size, and patient age), whereas the p53 mutation status, as determined by sequencing, did not. This demonstrates that the expression classifier is more directly prognostic of patient survival than is p53 mutation status alone.

The p53 Signature Predicts Outcome in Independent Therapy-Specific Data

Sets. We next assessed the prognostic capability of the p53 signature genes in therapy-specific cohorts by using independent microarray data sets from the public domain (Fig. 4; and see Fig. 7, which is published as supporting information on the PNAS web site). First, we evaluated whether the signature genes were prognostic of tumor recurrence in the Ma *et al.* (20) data set of 60 breast tumors derived from patients treated with postoperative radiation and adjuvant tamoxifen monotherapy. In this cohort, patients with and without recurrent disease were matched with respect to tumor grade and tumor node metastasis stage. Twenty-two of the p53 signature genes mapped to 27 probes on the Ma *et al.* spotted oligonucleotide array. Hierarchical cluster analysis with these genes revealed two to three primary tumor clusters with expression profiles that resembled the mt-like and wt-like configurations (Fig. 4*A*). Using these tumor clusters to define patient survival groups, we analyzed disease-free survival (DFS) by the Kaplan-Meier estimate. As shown in Fig. 4 *B* and *C*, the clusters were significantly associated with tumor recurrence [$P = 0.01$ (two clusters, C1 and C2) and $P = 0.005$ (three clusters, C1, C2, and C3)]. Thus, concordant results in two independent studies suggest that functional p53 deficiency, as assessed by an expression readout, is predictive of outcome to hormonal therapy.



**View larger
version (34K):**
[\[in this window\]](#)
[\[in a new window\]](#)

Fig. 4. The p53 signature predicts survival in independent clinically diverse data sets. (A) Tumor dendrogram from clustering 60 tumors and 22 genes (27 probes) from Ma *et al.* (20). (B and C) Patient subgroups determined by the primary tumor branches (C1-C4) were analyzed for correlations with DFS. (D) Tumor dendrogram from clustering 76 tumors and 9 genes from Sørlie *et al.* (18). Patient subgroups defined by the primary tumor branches (C1 and C2) were analyzed for correlations with disease-specific survival (DSS) (E) and DFS (F). (G) Tumor dendrogram from clustering 97 tumors and 21 genes (25 probes) from van't Veer *et al.* (21). (H and I) Primary tumor clusters (C1-C5) defined patient subgroups for DFS analysis. Red branches denote tumors with the p53 mt-like signature; black branches identify those with the wt-like signature. Black triangles indicate patients who relapsed within 5 years. See Fig. 7 for gene heat maps and probe IDs.

To examine the prognostic performance of the p53 signature genes in patients treated with systemic chemotherapy, we used the Sørlie *et al.* cDNA microarray data set. The majority of patients (>80%) in the Sørlie study received weekly doxorubicin or 5FU and mitomycin and were comprised mostly of late-stage patients (10, 11). Here, the nine-gene partial signature that could distinguish mt and wt tumors with 77% accuracy, was used to hierarchically cluster 76 well sampled tumors with corresponding treatment and survival data (Fig. 4D). Again, we observed the tumors cluster into two primary branches with expression patterns characteristic of the wt-like and mt-like configurations. Survival analysis resulted in a highly significant difference in outcome between patients with mt-like and wt-like tumors [$P = 7.5 \times 10^{-5}$ (disease-specific survival) and $P = 5.0 \times 10^{-5}$ (DFS)]; Fig. 4E and F) despite the small number of genes used. Notably, Fig. 4E predicts a remarkable 5-year 90% survival rate for the 31 p53 wt-like patients, compared with a 35% probability of 5-year survival for the 44 p53 mt-like patients.

Next, we tested the performance of the signature genes on a set of 97 early stage tumors (T1/T2, N0), from patients <55 years of age at diagnosis and treated by radiotherapy alone (21). From our 32-gene signature, we were able to map 25 probes corresponding to 21 signature genes to all 97 tumors with outcome information. Unsupervised clustering revealed two primary and four secondary tumor clusters (Fig. 4G) that could significantly discern patients based on time to distant metastasis within a 5-year period [Fig. 4 H and I; $P = 0.0006$ (two clusters, C1 and C2) and $P = 0.001$ (four clusters, C1, C2, C3, and C4)]. Notably, of the 24 tumors in cluster 1 (C1) that bear the molecular configuration of p53 mt-like tumors, 75% belonged to patients who developed a distant metastasis within 5 years, compared with 26% of 34 patients with tumors comprising C4 (which most closely resemble the p53 wt-like signature). These findings indicate that the p53 signature is also prognostic of recurrence in early stage, locally treated breast cancer.

The p53 Signature Genes Are Not Canonical p53 Targets. To gain some mechanistic insights, we examined the functional annotations of the signature genes for clues to explain their correlations with p53 status and patient outcome. We found that none of the signature genes are known transcriptional targets of p53, nor have they been previously implicated in the p53 pathway. Moreover, promoter analysis revealed no evidence of p53-binding sites. Of the characterized genes, a number are associated with cell growth and proliferation (*MYBL2*, *TFF1*, *BRRN1*, *CHAD*, *SCGB3A1*, *DACH*, and *CDCA8*), transcription (*LAF4*, *NY-BR-1*, *DACH*, and *MYBL2*), ion transport (*CACNG4*, *CY-BRD1*, and *LRP2*), and breast cancer biology (*SCGB3A1*, *TFF1*, *STC2*, *NY-BR-1*, and *AGR2*). Interestingly, *MYBL2*, which was transcriptionally up-regulated in the p53 mt-like tumors, is a growth-promoting transcription factor structurally related to the *c-MYB* oncogene. *MYBL2* maps to a chromosomal region frequently amplified in breast cancer (20q13) and has previously been reported to be overexpressed in breast cancer cell lines and sporadic ovarian carcinomas (28, 29). *SCGB3A1* (*HIN1*), which we observed to be down-regulated in the p53 mt-like tumors, is a putative tumor-

suppressor gene that can inhibit breast cancer cell growth when overexpressed and has been found to be transcriptionally silenced by promoter hypermethylation in early stages of breast tumorigenesis (30). Thus, some of the p53 signature genes may contribute mechanistically to the poor prognosis associated with the p53 mt-like tumors.

► Discussion

Breast cancers are characterized by multiple genetic alterations that, together, comprise the genotype that dictates tumor behavior. It is therefore reasonable that the compilation of genetic changes is a better indicator of clinical behavior than a single gene. Herein, we show that an expression signature, deduced from differences in the molecular configurations of p53 wt and mt tumors, predicts for p53 functional inactivation in primary breast cancers and provides a more accurate and useful measure of p53 clinical functionality than p53 mutation status alone. We show that, in independent data sets of both breast and liver cancers and regardless of other clinical features, subsets of the p53 signature can predict p53 status with significant accuracy. As a predictor of disease-specific survival, we found that the signature significantly outperformed p53 mutation status in a large patient cohort with heterogeneous treatment. Importantly, the p53 signature could significantly distinguish patients having more or less benefit from specific systemic adjuvant therapies and locoregional radiotherapy. Recently, Ma *et al.* identified by microarray analysis two genes (*HOXB13* and *IL17RB*) whose expression ratio was predictive of tamoxifen response. Notably, we found that these genes were also predictive of disease-specific survival in the 67 Uppsala patients treated with tamoxifen monotherapy ($P < 0.01$; data not shown). However, these genes were not prognostic of recurrence in the van't Veer data set, nor were the van't Veer 70 genes prognostic of recurrence in the Ma data set (20),

- ▲ [Top](#)
- ▲ [Abstract](#)
- ▲ [Methods](#)
- ▲ [Results](#)
- [Discussion](#)
- ▼ [References](#)

suggesting that tumor stage and/or therapeutic context is an important determinant of the prognostic capacity of some genes. In contrast, we demonstrate that the p53 signature genes are robustly prognostic of survival and recurrence in both early and late stage disease and in different therapeutic settings.

Although the p53 pathway may be compromised at some level in most human cancers, our analysis of genes involved in the p53 pathway suggests that the p53 expression signature defines some operational configuration of this pathway in breast tumors (more so than p53 mutation status alone) that impacts patient survival and therapeutic response. Recent evidence suggests that tumor sensitivity to some anti-cancer agents may depend largely on the relative intactness of p53-dependent mechanisms of apoptosis ([7](#), [8](#), [10](#), [11](#)) and that taxols (microtubule stabilizers), in particular, may have greater efficacy against p53-mt breast tumors than anthracycline-based (genotoxic) compounds ([9](#)). Whether the p53 classifier genes identified here are involved in some aspect of this p53 function or will have robust clinical utility as a predictor of therapeutic response warrants further investigation.

Other studies have elucidated gene expression signatures prognostic of breast cancer outcomes ([21](#), [31](#)). Although a 21-gene subset of our p53 signature could significantly distinguish patients with recurrent and nonrecurrent disease in the van't Veer study ([21](#)), none of these genes were found to overlap with the 231 genes identified as prognostic discriminators in the van't Veer set; and only one of the classifier genes, *MYBL2*, was found in the Sotiriou 485 survival-correlated genes ([31](#)). Similarly, none of the p53 signature genes were found in the top 25 relapse-associated genes reported by Ma *et al.* Thus, the p53 signature genes identified here represent a previously unrecognized prognostic cassette.

In cancer, it is clear that not all p53 mutations have equal effects; some simply confer loss of function, whereas others have a dominant-negative effect (such as

transdominant suppression of wt p53 or oncogenic gain of function), whereas still others show only a partial loss of function where, for example, only a fraction of p53 target genes are dysregulated ([32](#), [33](#)). For these reasons, no single molecular assessment of p53 status appears to provide an absolute indication of the complete p53 function. Although the p53 classification method developed here seeks to categorize all tumors as either p53-deficient or not, it is likely that intermediate types exist with partial p53 functionality, distinguished by expression patterns that fall between those of the predominant mt-like and wt-like classes. Further investigation will be required to resolve the biological and clinical implications of such intermediate tumor classes.

► **Acknowledgements**

We thank A. Ramasamy and K. R. Govindarajan for developing the database for cross-study analyses and other analytical tools used herein and Professor Hans Nordgren for Elston-grading of the tumor material. This study was supported by the Singapore Agency for Science Technology and Research and by grants from the Swedish Cancer Society, the Stockholm Cancer Society, and the King Gustav Fifth Jubilee Fund.

► **Footnotes**

Freely available online through the PNAS open access option.

Abbreviations: DFS, disease-free survival; ER, estrogen receptor; mt, mutant; wt, wild type.

[†] To whom correspondence may be addressed. E-mail: millerl@gis.a-star.edu.sg or liue@gis.a-star.edu.sg.

► References

▲ [Top](#)
 ▲ [Abstract](#)
 ▲ [Methods](#)
 ▲ [Results](#)
 ▲ [Discussion](#)
 • [References](#)

1. Hollstein, M., Sidransky, D., Vogelstein, B. & Harris, C. C. (1991) *Science* **253**, 49-53.[\[ISI\]](#)[\[Medline\]](#)
2. Levine, A. J., Momand, J. & Finlay, C. A. (1991) *Nature* **351**, 453-456.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
3. Peller, S. (1998) *Semin. Cancer Biol.* **8**, 379-387.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
4. Pharoah, P. D., Day, N. E. & Caldas, C. (1999) *Br. J. Cancer* **80**, 1968-1973.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
5. Lowe, S. W., Bodis, S., McClatchey, A., Remington, L., Ruley, H. E., Fisher, D. E., Housman, D. E. & Jacks, T. (1994) *Science* **266**, 807-810.[\[ISI\]](#)[\[Medline\]](#)
6. Bergh, J., Norberg, T., Sjogren, S., Lindgren, A. & Holmberg, L. (1995) *Nat. Med.* **1**, 1029-1034.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
7. Aas, T., Borresen, A. L., Geisler, S., Smith-Sorensen, B., Johnsen, H., Varhaug, J. E., Akslen, L. A. & Lonning, P. E. (1996) *Nat. Med.* **2**, 811-814.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
8. Berns, E. M., Foekens, J. A., Vossen, R., Look, M. P., Devilee, P., Henzen-Logmans, S. C., van Staveren, I. L., van Putten, W. L., Inganas, M., Meijer-van Gelder, M. E., *et al.* (2000) *Cancer Res.* **60**, 2155-2162.[\[Abstract/Free Full Text\]](#)
9. Kandioler-Eckersberger, D., Ludwig, C., Rudas, M., Kappel, S., Janschek, E., Wenzel, C., Schlagbauer-Wadl, H., Mittlbock, M., Gnant, M., Steger, G. & Jakesz, R. (2000) *Clin. Cancer Res.* **6**, 50-56.[\[Abstract/Free Full Text\]](#)
10. Geisler, S., Borresen-Dale, A. L., Johnsen, H., Aas, T., Geisler, J., Akslen, L. A., Anker, G. & Lonning, P. E. (2003) *Clin. Cancer Res.* **9**, 5582-5588.[\[Abstract/Free Full Text\]](#)
11. Geisler, S., Lonning, P. E., Aas, T., Johnsen, H., Fluge, O., Haugen, D. F., Lillehaug, J. R., Akslen, L. A. & Borresen-Dale, A. L. (2001) *Cancer Res.* **61**, 2505-2512.[\[Abstract/Free Full Text\]](#)
12. Archer, S. G., Eliopoulos, A., Spandidos, D., Barnes, D., Ellis, I. O., Blamey, R. W., Nicholson, R. I. & Robertson, J. F. (1995) *Br. J. Cancer* **72**, 1259-1266.[\[ISI\]](#)[\[Medline\]](#)
13. Rozan, S., Vincent-Salomon, A., Zafrani, B., Validire, P., De Cremoux, P., Bernoux, A., Nieruchalski, M., Fourquet, A., Clough, K., Dieras, V., *et al.* (1998) *Int. J. Cancer* **79**, 27-33.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)

14. Borresen-Dale, A. L. (2003) *Hum. Mutat.* **21**, 292-300.[CrossRef][ISI][Medline]
15. Dudoit, S., Frilyand, J. & Speed, T. P. (2002) *J. Am. Stat. Assoc.* **97**, 77-87.
[CrossRef][ISI]
16. Ripley, B. D. (1996) in *Pattern Recognition and Neural Networks* (Cambridge Univ. Press, Cambridge, U.K.), pp. 181-212.
17. Vapnik, V. (1995) *The Nature of Statistical Learning Theory* (Springer, New York).
18. Sørlie, T., Tibshirani, R., Parker, J., Hastie, T., Marron, J. S., Nobel, A., Deng, S., Johnsen, H., Pesich, R., Geisler, S., *et al.* (2003) *Proc. Natl. Acad. Sci. USA* **100**, 8418-8423.[Abstract/Free Full Text]
19. Chen, X., Cheung, S. T., So, S., Fan, S. T., Barry, C., Higgins, J., Lai, K. M., Ji, J., Dudoit, S., Ng, I. O., *et al.* (2002) *Mol. Biol. Cell* **13**, 1929-1939.
[Abstract/Free Full Text]
20. Ma, X. J., Wang, Z., Ryan, P. D., Isakoff, S. J., Barmettler, A., Fuller, A., Muir, B., Mohapatra, G., Salunga, R., Tuggle, J. T., *et al.* (2004) *Cancer Cell* **5**, 607-616.
[CrossRef][ISI][Medline]
21. van 't Veer, L. J., Dai, H., van de Vijver, M. J., He, Y. D., Hart, A. A., Mao, M., Peterse, H. L., van der Kooy, K., Marton, M. J., Witteveen, A. T., *et al.* (2002) *Nature* **415**, 530-536.[CrossRef][ISI][Medline]
22. Tusher, V. G., Tibshirani, R. & Chu, G. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 5116-5121.[Abstract/Free Full Text]
23. Cattoretti, G., Rilke, F., Andreola, S., D'Amato, L. & Delia, D. (1988) *Int. J. Cancer* **41**, 178-183.[ISI][Medline]
24. Draper, N. R. & Smith, H. (1998) in *Applied Regression Analysis* (Wiley, New York), 3rd Ed, pp. 217-234.
25. Mitchell, K. O., Ricci, M. S., Miyashita, T., Dicker, D. T., Jin, Z., Reed, J. C. & El-Deiry, W. S. (2000) *Cancer Res.* **60**, 6318-6325.[Abstract/Free Full Text]
26. Ceballos, E., Delgado, M. D., Gutierrez, P., Richard, C., Muller, D., Eilers, M., Ehinger, M., Gullberg, U. & Leon, J. (2000) *Oncogene* **19**, 2194-2204.[CrossRef][ISI][Medline]
27. Hainaut, P., Hernandez, T., Robinson, A., Rodriguez-Tome, P., Flores, T., Hollstein, M., Harris, C. C. & Montesano, R. (1998) *Nucleic Acids Res.* **26**, 205-213.[Abstract/Free Full Text]
28. Forozan, F., Mahlamaki, E. H., Monni, O., Chen, Y., Veldman, R., Jiang, Y., Gooden, G. C., Ethier, S. P., Kallioniemi, A. & Kallioniemi, O. P. (2000) *Cancer Res.* **60**, 4519-4525.[Abstract/Free Full Text]
29. Tanner, M. M., Grenman, S., Koul, A., Johannsson, O., Meltzer, P., Pejovic, T., Borg, A. & Isola, J. J. (2000) *Clin. Cancer Res.* **6**, 1833-1839.

[\[Abstract/Free Full Text\]](#)

30. Krop, I. E., Sgroi, D., Porter, D. A., Lunetta, K. L., LeVangie, R., Seth, P., Kaelin, C. M., Rhei, E., Bosenberg, M., Schnitt, S., *et al.* (2001) *Proc. Natl. Acad. Sci. USA* **98**, 9796-9801.[\[Abstract/Free Full Text\]](#)
31. Sotiriou, C., Neo, S. Y., McShane, L. M., Korn, E. L., Long, P. M., Jazaeri, A., Martiat, P., Fox, S. B., Harris, A. L. & Liu, E. T. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 10393-10398.[\[Abstract/Free Full Text\]](#)
32. Epstein, C. B., Attiyeh, E. F., Hobson, D. A., Silver, A. L., Broach, J. R. & Levine, A. J. (1998) *Oncogene* **16**, 2115-2122.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)
33. Monti, P., Campomenosi, P., Ciribilli, Y., Iannone, R., Inga, A., Abbondandolo, A., Resnick, M. A. & Fronza, G. (2002) *Oncogene* **21**, 1641-1648.[\[CrossRef\]](#)[\[ISI\]](#)[\[Medline\]](#)

Related articles in PNAS:

This Week in PNAS

PNAS 2005 102: 13353-13354. [\[Extract\]](#) [\[Full Text\]](#)

This Article

Free via Open Access: OA

- ▶ [OA Abstract](#)
- ▶ [Figures Only](#)
- ▶ [Full Text \(PDF\)](#)
- ▶ [Supporting Information](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)
- ▶ [Citation Map](#)

Services

- ▶ [Related articles in PNAS](#)
- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Add to My File Cabinet](#)
- ▶ [Download to citation manager](#)
- ▶ [Request Copyright Permission](#)
- ▶ [Articles by Miller, L. D.](#)
- ▶ [Articles by Bergh, J.](#)

PubMed

- ▶ **PubMed Citation**
- ▶ **Articles by Miller, L. D.**
- ▶ **Articles by Bergh, J.**

[Current Issue](#) | [Archives](#) | [Online Submission](#) | [Info for Authors](#) | [Editorial Board](#) | [About](#)
[Subscribe](#) | [Advertise](#) | [Contact](#) | [Site Map](#)

[Copyright © 2005 by the National Academy of Sciences](#)

ATTACHMENT 4

ACADEMY REPORTS ONLINE!

Sign up for PNAS Online eTocs

Get notified by email when new content goes on-line

[Info for Authors](#) | [Editorial Board](#) | [About](#) | [Subscribe](#) | [Advertise](#) | [Contact](#) | [Site Map](#)

PNAS

Proceedings of the National Academy of Sciences of the United States of America

[Current Issue](#)

[Archives](#)

[Online Submission](#)

GO

[advanced search >>](#)

**Published online before print August 15, 2005,
10.1073/pnas.0505446102**

PNAS | August 23, 2005 | vol. 102 | no. 34 | 12200-12205

OPEN ACCESS ARTICLE

MICROBIOLOGY

Ancestral antibiotic resistance in *Mycobacterium tuberculosis*

Rowan P. Morris^{*,†}, Liem Nguyen^{†,‡},
John Gatfield[†], Kevin Visconti^{*},
Kien Nguyen^{†,§}, Dirk Schnappinger[¶],
Sabine Ehrt[¶], Yang Liu^{*}, Leonid Heifets^{||},
Jean Pieters[†], Gary Schoolnik^{*} and
Charles J. Thompson^{†,‡,**}

^{*} Departments of Medicine and of Microbiology and Immunology, Division of Infectious Diseases and Geographic Medicine, Stanford Medical School, Stanford, CA 94305; [†] Biocentre, University of Basel, Klingelbergstrasse 50/70, CH-4056 Basel, Switzerland; [‡] University of British Columbia, Life Sciences Centre, 2350 Health Sciences Mall,

Vancouver, BC, Canada V6T 1Z3; [¶] Department of Microbiology and Immunology, Weill Graduate School of Medical Sciences, Cornell University, New York, NY 10021; and

This Article

Free via Open Access: OA

- ▶ [OA Abstract](#)
- ▶ [Figures Only](#)
- ▶ [Full Text \(PDF\)](#)
- ▶ [Supporting Information](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)
- ▶ [Citation Map](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Add to My File Cabinet](#)
- ▶ [Download to citation manager](#)
- ▶ [Cited by other online articles](#)
- ▶ [Request Copyright Permission](#)

Google Scholar

- ▶ [Articles by Morris, R. P.](#)
- ▶ [Articles by Thompson, C. J.](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Morris, R. P.](#)
- ▶ [Articles by Thompson, C. J.](#)

¹¹National Jewish Medical and Research Center, Denver, CO 80206

Communicated by Werner Arber, University of Basel, Basel, Switzerland, June 29, 2005
(received for review March 2, 2005)

► Abstract

Chemotherapeutic options to treat tuberculosis are severely restricted by the intrinsic resistance of *Mycobacterium tuberculosis* to the majority of clinically applied antibiotics. Such resistance is partially provided by the low permeability of their unique cell envelope. Here we describe a complementary system that coordinates resistance to drugs that have penetrated the envelope, allowing mycobacteria to tolerate diverse classes of antibiotics that inhibit cytoplasmic targets. This system depends on *whiB7*, a gene that pathogenic *Mycobacterium* shares with *Streptomyces*, a phylogenetically related genus known as the source of diverse antibiotics. In *M. tuberculosis*, *whiB7* is induced by subinhibitory concentrations of antibiotics (erythromycin, tetracycline, and streptomycin) and *whiB7* null mutants (*Streptomyces* and *Mycobacterium*) are hypersusceptible to antibiotics *in vitro*. *M. tuberculosis* is also antibiotic sensitive within a monocyte model system. In addition to antibiotics, *whiB7* is induced by exposure to fatty acids that pathogenic *Mycobacterium* species may accumulate internally or encounter within eukaryotic hosts during infection. Gene expression profiling analyses demonstrate that *whiB7* transcription determines drug resistance by activating expression of a regulon including genes involved in ribosomal protection and antibiotic efflux. Components of the *whiB7* system may serve as attractive targets for the identification of inhibitors that render *M. tuberculosis* or multidrug-resistant derivatives more antibiotic-sensitive.

▲ Top
• Abstract
▼ Methods
▼ Results
▼ Discussion
▼ References

multidrug resistance | *Streptomyces* | WhiB | microarray | gene expression

The World Health Organization has estimated that between 2000 and 2020, nearly one billion people will be newly infected, 200 million people will get sick, and 35 million will die from tuberculosis (TB) (1). It is the remarkable antibiotic tolerance of the infectious agent *Mycobacterium tuberculosis* to many commonly used broad-spectrum antibiotics that limits chemotherapeutic options and is the root cause of all treatment failure (2). Tolerance may reflect physiological adaptations that occur within the host, perhaps including an undefined developmental or physiological state that underlies persistent infection (3). As a result, patients must be treated with multiple antibiotics for 6-12 months. Patient noncompliance or inadequate drug dosage favors the sequential acquisition of mutations providing resistance and the emergence of multidrug-resistant *M. tuberculosis* strains. In contrast to acquired resistance, intrinsic resistance in *Mycobacterium* has largely been attributed to its impermeable mycolic acid-containing cell envelope (4, 5) that is not found in many other Actinomycetes including *Streptomyces*. However, Jarlier and Nikaido (4) have also pointed out that this permeability barrier is insufficient to fully explain the high levels of drug resistance in *Mycobacterium*, suggesting that there must be synergistic systems effective against drugs that penetrate this barrier. Indeed, several mycobacterial genes not involved in outer envelope assembly confer resistance to specific, broad-spectrum antibiotics (6-8).

Although the best-known *Mycobacterium* species are pathogenic, most are ubiquitous environmental saprophytes belonging to the Actinomycete taxon (9). The taxon also include *Streptomyces* species, filamentous bacteria known for their extraordinary capacity to produce thousands of diverse antibiotics as a part of a developmental program leading to sporulation. Antibiotic biosynthetic genes are found in clusters that typically include the corresponding resistance genes to provide self-protection (10). However, as in other bacteria, genes scattered throughout the genome that may have alternative physiological roles can also confer antibiotic resistance (11). Intuitively, the protective activity of these resistance genes should be a prerequisite for the evolution

of antibiotic biosynthetic pathways.

Indeed, at sublethal concentrations, antibiotics can induce a wide variety of genes, many not known to provide antibiotic resistance (12-14). The expression of some of these antibiotic-induced genes is under the control of stress inducible systems that respond to (13) or lead to (15) decreases in growth rate. The underlying control elements that affect antibiotic resistance include general stress-responsive sigma factors (16) and transcriptional activator proteins of the AraC family (MarA, SoxR, and Rob) (17), as well as genetic systems that provide for more specific adaptation to DNA damage (15) or oxidative stress (8). Such systems are commonly found in diverse bacterial groups and typically modulate antibiotic resistance within a rather narrow concentration range (8, 17). Here we describe a multidrug-resistance system that apparently evolved in the ancestors of antibiotic producing bacteria, which has been retained in saprophytes and pathogens belonging to the Actinomycete taxon.

► **Methods**

Media and Strains. *Streptomyces lividans* 1326 was grown in the nutrient-rich liquid media YEME and cultivated on NE solid media (18).

The slow growing mycobacteria *Mycobacterium bovis* bacillus

Calmette-Guérin, *M. tuberculosis* H37Rv, and the clinical *M.*

tuberculosis isolate 1254 were propagated in 7H9 media (19), supplemented with 10% ADS (5% BSA/2% dextrose/0.8% sodium chloride).

Plasmid Constructions and Mutant Analyses. Annotated *whiB7* ORFs were deleted in the genomes of *S. lividans* and *Streptomyces coelicolor* (nucleotide coordinates 5,647,587-5,648,293; <http://jic-bioinfo.bbsrc.ac.uk/streptomyces/ScoDB>), *M. tuberculosis* H37Rv (nucleotide coordinates 3,568,405-3,568,801; <http://genolist.pasteur.fr/TubercuList>), and *M. bovis* bacillus Calmette-Guérin

▲ Top
▲ Abstract
• Methods
▼ Results
▼ Discussion
▼ References


(nucleotide coordinates 3,523,346-3,522,950; <http://genolist.pasteur.fr/BovList>) as described *Supporting Text* and *Data Set*, which are published as supporting information on the PNAS web site. Corresponding ORFs were expressed from vector promoters.

Mycobacterial Survival in Monocytes. Resting or activated J774 monocytes were grown in DMEM/FCS. Monocytes were exposed to *Mycobacterium bovis* bacillus Calmette-Guérin and the corresponding *whiB7* mutant at a multiplicity of one. Activation was achieved by 16-h exposure to 500 units/ml IFN- γ followed by 4-h exposure to both IFN- γ and 1 μ g/ml LPS. The monocytes were washed twice with PBS and incubated for 45 min at 37°C/5% CO₂ with amikacin (200 μ g/ml). Cells were again washed twice in PBS and incubated in DMEM/FCS. Survival was determined at the indicated incubation times by bacterial incorporation of tritiated uracil followed by liquid scintillation counting (20). For antibiotic susceptibility testing, the infected monocytes were incubated in the presence of indicated spectinomycin concentrations for 48 h before permeabilization and mycobacterial labeling (see *Supporting Text* for details).

Microarray Expression Profiling and Analysis. Labeling of RNA and hybridizations to 70-mer oligonucleotide-based microarrays (Operon) was performed as described (21). Microarrays were scanned by using GenePix 4000A (Axon Instruments). Fluorescence intensities of the two channels at each spot were quantified by using the SCANALYZE software (<http://rana.lbl.gov/EisenSoftware.htm>). After data for each array were normalized (21), expression ratios were averaged from two biological replicates for antibiotic-induced cultures or from three cultures for the mid-log comparison, and with two microarrays for each of the biological replicates. Data from each experimental condition was analyzed separately by using SIGNIFICANCE ANALYSIS OF MICROARRAYS (22) with a false discovery ratio $\leq 0.3\%$. Significantly regulated genes for all experimental conditions were combined to generate a data set containing 2,879 genes. Within this

list, gene expression data could be present for one experimental condition and absent from another. To aid hierarchical clustering, these genes were then filtered to include those that were present across 95% of the 25 experimental conditions (total number of rows in Fig. 3 *b-d*) and a differential expression >2-fold under at least three of these conditions. The resulting 880 filtered genes were organized according to their expression profiles by average linkage clustering using GENESIS software (<http://genome.tugraz.at/Software/GenesisCenter.html>).

Analysis of Mycobacterial RNA with Quantitative Real-Time RT-PCR. Real time PCR to confirm microarray analysis of *in vitro* grown cultures was performed by using SYBR green (Applied Biosystems). A standard curve was generated for the relative quantification of all genes, and a control reaction lacking reverse transcriptase was performed for every RNA sample. The major housekeeping sigma factor gene *sigA* was used to normalize mRNA levels. Gene induction values were calibrated by comparison with the reference RNA isolated for each experiment.



View larger version

(95K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 1. Identification of *whiB7*, a gene providing intrinsic antibiotic resistance in *Streptomyces*. (*a*) Antibiotic susceptibility of the wild-type *S. lividans* strain (*Left*) and the spontaneous *whiB7* mutant RM1 (*Right*). Etest strips were applied to seeded spores, and minimal inhibition concentration values were read from the scale ($\mu\text{g/ml}$) at the point of intersection between inhibition ellipse edge and the strip. Upper, erythromycin; right, tetracycline; lower, rifampicin; left, quinupristin/dalfopristin. (*b*) *S. lividans* RM1 was engineered to allow thiostrepton-inducible expression of *whiB7* by using the expression plasmid pIJ8600. Seeded spores of *S. lividans* wild-type/pIJ8600 (*Left*), the *whiB7* mutant RM1/pIJ8600 (*Center*), or RM1/pIJ8600::*whiB7* (*Right*) were exposed to radial gradients by discs containing 100 μg of oleandomycin

(left), chlorotetracycline (right), or thiostrepton (top). The high frequency of suppressor colonies in the RM1/pIJ8600::*whiB7* culture are presumed to be promoter-up mutants that allow unregulated *whiB7* expression. Tetracycline induced synthesis of a red pigment (likely to be the antibiotic undecylprodigiosin) in both wild-type *S. lividans* and the *whiB7* mutant.


► Results

Identification of a Multidrug Resistance Regulator in

***Streptomyces*.** We isolated a spontaneous mutant of *S. lividans*, RM1, that was hypersensitive to a diverse array of chemically and functionally unrelated clinical antibiotics that it does not synthesize, including chloramphenicol, fusidic acid, imipenem, lincosamides (clindamycin and lincomycin), macrolides (erythromycin, oleandomycin, and spiramycin), rifampicin, streptogramins (pristinamycin and virginiamycin), and tetracycline (see Fig. 5, which is published as [supporting information](#) on the PNAS web site). Sensitivities of the mutant and wild-type parent were quantified by using Etest diffusion strips to compare their minimal inhibition concentrations to four structurally and functionally distinct classes of antibiotics. RM1 was 600-, 400-, 25-, and 40-fold more sensitive to erythromycin, tetracycline, rifampicin, and the pristinamycin derivatives quinupristin/dalfopristin, respectively (Fig. 1*a*). The mutant displayed large decreases in intrinsic antibiotic resistance, but its susceptibility to a variety of other toxic, nonantibiotic stresses, including detergents, antiseptics, and oxidative stress inducers, was unchanged (Fig. 6, which is published as [supporting information](#) on the PNAS web site). Standard cloning, sequencing, and site-directed mutagenesis experiments (described in *Supporting Text*) identified the gene responsible for this multidrug resistance as *whiB7* in both *S. lividans* (GenBank accession no. AF205848 [[GenBank](#)]) and *S. coelicolor* genomes

▲	Top
▲	Abstract
▲	Methods
•	Results
▼	Discussion
▼	References

(Fig. 2) (23). Sequence comparison of the wild-type locus to that of RM1 revealed a frame shift in the *whiB7* gene resulting from the insertion of a cytosine at nucleotide position 239. This locus was not linked to any recognizable antibiotic biosynthetic cluster. *whiB7* encodes a 122-aa protein related to *whiB*, a putative *S. coelicolor* transcriptional regulatory gene (24). In *M. tuberculosis*, a WhiB-like protein (WhiB3) may act as a transcriptional regulator by binding and modulating the activity of RpoV (SigA), the principle sigma factor in *M. tuberculosis* (25). To further demonstrate the correlation of *whiB7* transcriptional activity with antibiotic resistance, the *S. lividans* mutant RM1 was engineered to allow inducible expression (26) of a wild-type copy of *whiB7*. Fig. 1*b* shows that the circular zones of inhibition caused by diffusion of tetracycline (an aromatic polyketide) or oleandomycin (a macrolide) were distorted and reduced by a radial gradient of the inducer (thiostrepton), indicating higher levels of resistance associated with increased *whiB7* transcription.



View larger version

(27K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 2. The orthologous *whiB7* loci of *Streptomyces* and *Mycobacterium*. (a) Alignment of the WhiB7 proteins from both *Mycobacterium tuberculosis* (WhiB7-tub) and *leprae* (WhiB7-lep) with the *Streptomyces* WhiB7 and the prototypic family member WhiB from *Streptomyces coelicolor*. Four absolutely conserved cysteine residues and a tryptophan-containing/glycine-rich motif are conserved throughout the WhiB family (circled). An A/T-Hook DNA binding consensus sequence is found only in WhiB7 paralogs. ~, N-terminal sequence not shown. (b) Gene organization of the *whiB7* genomic region. Shaded block arrows represent conserved ORFs.

Members of the *whiB* gene lineage, including *whiB7* of *S. lividans* (alternatively named *wblC*), are restricted to the Actinomycetes (27); BLAST searches did not identify orthologs in any other published bacterial genome sequences. The prototype of this


gene family, *whiB*, was identified as a developmental gene in *Streptomyces* species that is essential for the differentiation of mycelium into pigmented spores (*white*) (28). The family signature is defined by four absolutely conserved cysteines that form an oxygen sensitive iron sulfur cluster (ref. 29 and L.N., P. Jensen, R.P.M., M. Folcher, S. Durr, S. Grzesiek, and C.J.T., unpublished data), and a tryptophan within a glycine-rich sequence (Fig. 2a). In addition, *whiB7* paralogs also encode a C-terminal "A/T Hook" domain that is known to bind AT-rich DNA sequences. Although the *M. tuberculosis* genome encodes seven *whiB*-like genes (*whiB1-7*), both homology and synteny predict a minimal core of five orthologous *whiB*-like genes common to *M. tuberculosis*, *Mycobacterium leprae*, and *Streptomyces* species (*whiB1-4* and *-7*). TBLASTN searches of the 201 completed bacterial genomes identified *whiB7* orthologs in all species of *Streptomyces* (*S. coelicolor* and *Streptomyces averimidilus*), *Mycobacterium* (*tuberculosis* H37Rv, *tuberculosis* CDC1551, *bovis* AF2122/97, *leprae* TN, *avium* subsp. *paratuberculosis*), and *Nocardia* (*farcinica* IFN10152). The role of the streptomycete *whiB7* gene in determining broad spectrum drug resistance, predicted that it might play a similar role in pathogenic *M. tuberculosis*.

A *whiB7* Ortholog Controls Multidrug Resistance in *M. tuberculosis*. Intrinsic resistance in *M. tuberculosis* could be partially due to a *whiB7* ortholog that is able to provide resistance to antibiotics that have penetrated the cell envelope and entered the cytoplasm. To test this hypothesis, we constructed a gene replacement mutant in *M. tuberculosis*, strain H37Rv. The mutant grew normally, but was defective in its resistance (Table 1) to a variety of antibiotics including macrolides, a lincosamide, and an aminoglycoside. The *whiB7* gene was cloned into the integrative vector pMV361 to provide expression from a strong constitutive promoter (hsp60). Integration of this plasmid (pRPM251) into the chromosome of the *M. tuberculosis whiB7* mutant restored normal, or slightly elevated levels of antibiotic resistance (Table 1). Multidrug sensitivity also resulted from disruption of the *whiB7* gene of the fast growing saprophytic *Mycobacterium smegmatis* (R.P.M. and C.J.T., unpublished results).

View this table: Table 1. Multidrug resistance in *M. tuberculosis*
[\[in this window\]](#) **determined by *whiB7***
[\[in a new window\]](#)

Induction of the *M. tuberculosis whiB7* Gene by Multiple Antibiotics and Fatty Acids. Antibiotic resistance genes typically confer resistance to one class of antibiotic and are specifically activated by the corresponding drugs. Experiments were carried out to determine whether the broad spectrum of resistance conferred by *whiB7* might be controlled by regulatory systems that are responsive to dissimilar drugs (30). Microarray transcript profiling of all annotated *M. tuberculosis* genes was used to monitor expression in response to three chemically distinct classes of common antibiotics: the frontline antimycobacterial drug streptomycin, as well as erythromycin and tetracycline (Fig. 3*b*). *M. tuberculosis* 1254 cultures were treated with five concentrations of each antibiotic, spanning three orders of magnitude (0.5-100 $\mu\text{g/ml}$) including the minimal inhibition concentration. Expression was assayed 15 min after exposure to maximize detection of genes whose regulation most directly reflected *whiB7* activity, rather than downstream pleiotropic effects. *whiB7* expression was significantly induced by subinhibitory concentrations of both erythromycin (1.0 $\mu\text{g/ml}$) and tetracycline (0.5 $\mu\text{g/ml}$) and also higher levels of streptomycin (25 $\mu\text{g/ml}$). After longer exposure (24 h), concentrations of streptomycin as low as 1 $\mu\text{g/ml}$ induced *whiB7* (Fig. 3*c*). The levels of induction were dose dependent for all three antibiotics (Fig. 3*b*). Activation of *whiB7* transcription by tetracycline (1 $\mu\text{g/ml}$) was confirmed by quantitative RT-PCR showing that *whiB7* RNA levels progressively increased ≈ 70 -fold during 24 h of exposure (Fig. 3*e*).

Fig. 3. Identification of antibiotic resistance genes as parts of the *M. tuberculosis whiB7* regulon. Significantly altered gene



**View larger
version (21K):**

[\[in this window\]](#)

[\[in a new window\]](#)

expression ratios from all experiments were averaged, \log_2 transformed, and clustered according to the displayed color code. Red and blue indicate higher and lower gene expression, respectively, in experimental samples, in comparison to the reference. Black represents no difference. (a) Genes of the *whiB7* regulon. Rv and ORF numbers designate annotated *M. tuberculosis* genes, and arrows indicate contiguous genes. (b) Treatment (15 min) of *M. tuberculosis* 1254 with 0.5, 1.0, 5, 25 and 100 $\mu\text{g/ml}$ of erythromycin, streptomycin, and tetracycline. (c) Extended antibiotic treatment for 2 and 24 h (1 $\mu\text{g/ml}$). (d) *M. tuberculosis* H37Rv was used as reference and compared to a *whiB7* null mutant (WhiB7 KO) and a H37Rv strain engineered to overexpress *whiB7* (WhiB7 OV). Cells were assayed at an optical density of 0.4 at 600 nm. (e) *whiB7* induction by fatty acids. Quantitative RT-PCR-determined induction factors of *whiB7* expression. Primary y axis: *M. tuberculosis* grown in MDG fed 50 μM palmitic acid (purple) or its unsaturated form, oleic acid (gray). Secondary y axis: induction to prolonged exposure of tetracycline (connected dots) ($\approx 2 \mu\text{M}$, 1 $\mu\text{g/ml}$).

Many antibiotics, including erythromycin and tetracycline, are based on polyketides, fatty acid-like molecules with carbon backbones synthesized by enzyme complexes similar to fatty acid synthase. Like antibiotics, many fatty acids are known to suppress growth of diverse bacteria (31), including *Mycobacterium spp.* (32). Palmitic acid, as well as an unsaturated derivative, oleic acid, were likewise tested for their abilities to induce *whiB7* transcription by quantitative RT-PCR. Although both fatty acids activated *whiB7* transcription, the palmitic acid response was more rapid and achieved higher levels of induction. Although induction kinetics were concentration dependent, at least for the antibiotics tested, higher concentrations of externally applied palmitic acid were needed and lower levels of maximal induction were achieved (3- to 4-fold compared to 70-fold for tetracycline).

In conclusion, *whiB7* expression was progressively induced at the transcriptional level

by sublethal concentrations of antibiotics and fatty acids. Up-regulation of *whiB7* expression may be required for the induction of other genes that could plausibly provide antibiotic resistance. These observations suggested that *whiB7* encoded a regulator whose transcriptional induction activated a regulon providing intrinsic antibiotic resistance.

Identification of Genes in the *whiB7* Regulon by Microarray Analyses. To determine whether the induction of *whiB7* was correlated with the expression of genes associated with antibiotic resistance, microarray expression profiles of mid-log phase cultures of the *whiB7* deletion mutant and a strain overexpressing *whiB7* were compared to parental strain *M. tuberculosis* H37Rv (Fig. 3d). These global analyses (details not presented) showed that *whiB7* was the only gene induced initially, after exposure to minimal concentrations of antibiotic (0.5 mg/ml tetracycline for 15 min, for example). Thus, *whiB7* represented a primary regulatory gene whose expression was followed by transcription of other genes in its regulon. Average distance hierarchical clustering identified 12 significantly regulated genes (SAM false discovery rate $\leq 0.3\%$) whose expression profile appeared to be influenced by antibiotic exposure and the activity of *whiB7* (Fig. 3). The *whiB7*-dependent set of eight transcripts includes three genes that may provide intrinsic antibiotic resistance: *tap* (Rv1258c), encoding an efflux pump that confers low-level resistance to aminoglycosides and tetracycline (33); an unstudied ORF encoding a putative macrolide transporter (Rv1473) with an ATP-binding cassette; and *erm* (Rv1988), homologous to ribosomal methyltransferases and conferring MLS (macrolide, lincosamide, and streptogramin) resistance by modification of 23S rRNA (7, 34). Although the *whiB7* regulon may include unrecognized antibiotic resistance determinants, other functions were also suggested. These include *eis* (Rv2416C), a putative acetyl-transferase providing enhanced survival within macrophages, Rv0263C, a putative carboxylase catalyzing urea degradation, and *cut2*, a putative cutinase/lipase that is reported to be exposed on the outside of the cell membrane and potentially able to release fatty acids from external lipids (35). These

possible functions of other genes in the putative *whiB7* regulon, not known to be antibiotic resistance determinants, require further investigation. Some may play roles in bacterial physiology, a recognized but not well understood determinant of antibiotic resistance (36).

Quantitative RT-PCR was used to independently confirm the induction data for genes within the *whiB7* regulon, including Rv1258, Rv1473, and Rv1988. Furthermore, primers targeting an intergenic sequence upstream of *whiB7* showed that *whiB7* was transcriptionally coupled to the small upstream unannotated ORF, ORFD0316 (Fig. 7, which is published as [supporting information](#) on the PNAS web site). Strains engineered to constitutively express *whiB7* in trans (*whiB7* OV) were associated with elevated levels of ORFD0316 transcription, and ORFD0316 was down-regulated in the *whiB7* mutant, suggesting that *whiB7* positively autoregulates its own transcription.


Multidrug Resistance in a Monocyte Model System. The physiology of *Mycobacterium* growing in laboratory cultures is much different from their natural state during host cell infection (36). To investigate whether *whiB7* controls survival or antibiotic resistance in a eukaryotic cellular environment, we monitored the intracellular survival of *M. bovis* bacillus Calmette-Guérin and a constructed isogenic *whiB7* mutant harbored within untreated or spectinomycin-treated J774, a monocyte-like cell line most commonly used for antibiotic sensitivity testing. In the absence of antibiotic, the wild type and *whiB7* mutant had similar survival curves in resting or IFN- γ -activated J774 during the first 72 h of infection (Fig. 4a). Compared to liquid cultures, both bacillus Calmette-Guérin wild type and the *whiB7* mutant were more sensitive to spectinomycin in J774. However, more importantly, in resting J774, the *whiB7* mutant was >10-fold more sensitive to spectinomycin (as reflected by the concentration of antibiotic needed to reduce transcription by 50%; Fig. 4b).

► Discussion

▲ Top
▲ Abstract
▲ Methods
▲ Results
• Discussion
▼ References

The fact that the *whiB7* gene and its multidrug resistance phenotype have been retained in most Actinomycetes, including those that have not been continuously exposed to antibiotics in the environment, provides important insights into the evolution and biological function of "antibiotic resistance" genes and their regulatory systems. There are 14 genes belonging to the *whiB* family in the sequenced genome of *S. coelicolor* (23). Site-directed mutagenesis of this set of genes (B. Gust and K. Chater, personal communication) has shown that, like *whiB7*, many do not have obvious sporulation (white) defects under standard growth conditions and that the multidrug sensitive phenotype is a unique characteristic of *whiB7* mutants (L.N. and C.J.T., unpublished results). Studies of *whiB* paralogs in *Mycobacterium* species have shown that the *M. smegmatis whiB2* gene (also called *whmD*) is essential (37) and that another, *whiB3*, plays a role in virulence in some model systems (25). Here we focus on the ability of *whiB7* to determine multiple antibiotic resistances in Actinomycetes and suggest that, in mycobacterial species, it acts synergistically with a rather impermeable cell envelope to provide high levels of intrinsic resistance.

whiB7 is notably different from systems reported in other bacteria that allow adaptation to a variety of different nonspecific stress conditions and may incidentally provide multiantibiotic resistance. *whiB7* does not confer resistance to antiseptics, but rather to antibiotics having specific targets (see Figs. 5 and 6). *whiB7* function is also unique in that it confers relatively high levels of resistance: in the *S. lividans* mutant, antibiotic sensitivity increased by orders of magnitude; this is distinct from the general stress adaptive system, which confers much lower levels of multidrug resistance (*mar*) in enteric bacteria (17, 38) by using any one of three transcriptional activators having highly redundant functions.



**View larger
version (15K):**
[\[in this window\]](#)
[\[in a new window\]](#)

Fig. 4. The phenotype of the *whiB7* mutation in J774, a monocyte-derived cell line. Mycobacterial RNA transcription, an indicator of survival, was assayed by incorporation of tritiated uracil. Shown are mean values (\pm SD) of six to eight determinations. Resting (REST) or activated (ACT) J774 monocytes growing in microtiter plates were exposed to *M. bovis* bacillus Calmette-Guérin and the corresponding *whiB7* mutant. (a) RNA synthesis of internalized WT and *whiB7* mutant cells decreased at the same rate in resting or activated monocytes, presumably because of bactericidal activity of the monocytes. In contrast, growth did occur in the control experiment where the bacteria were cultured in the same medium without monocytes (data not shown). (b) For antibiotic susceptibility testing, infected monocytes were incubated with medium containing the indicated concentrations of spectinomycin; % transcriptional activity is the percentage of incorporation rates in spectinomycin-treated vs. untreated bacteria. The same results were obtained in infected monocytes treated with amikacin after infection to confirm that incorporation rates reflected internalized mycobacteria (data not shown).

whiB7 is a putative transcriptional activator that is induced by antibiotics and controls the expression of at least two documented antibiotic resistance genes. The presence of these structural genes and corresponding regulatory systems in *Mycobacterium* suggests that this system provides selective advantage. The retention of the multiple antibiotic-responsive regulatory system controlling *M. tuberculosis whiB7* provides circumstantial evidence that toxic metabolites of various structures may have played a key role in directing the early evolution of the regulon to provide antibiotic resistance. The *whiB7* gene, as well as 5 of its 10 *M. tuberculosis* target genes (Fig. 3a), are present in *M. leprae* (Rv1473, Rv1257c, Rv1258c, Rv0263, and Rv2725), whose genome has undergone dramatic reduction during evolution within metazoan (presumably mammalian) hosts. The presence of the functionally conserved *whiB7* locus in all *Streptomyces* and *Mycobacterium* spp. (also including saprophytic *M.*

smegmatis) genomes now sequenced records its origin in their presumed soil dwelling ancestor. Although it is not clear why this capacity should be retained by *M. tuberculosis* and *M. leprae*, long after their progenitor left the antibiotic containing soil, some of these genes may have been adapted to protect the microbe against compounds of the mammalian immune system. The *whiB7* system was active in a monocyte model system; mutant was more sensitive to spectinomycin in J774 (Fig. 4b).

This evolutionary retention of *whiB7*, along with the observation that antibiotics with different structures activate it, implies a common endogenous inducer made by actinomycetes in response to antibiotics. Indeed, sublethal concentrations of some antibiotics induce synthesis of other secondary metabolites as demonstrated in *Streptomyces* (Figs. 1B and 5) that may also be autotoxic. Although *Mycobacterium* species are not recognized as antibiotic producers, they do have a remarkably large repertoire of polyketide biosynthesis gene clusters (39), some of which may encode biosynthetic pathways for autotoxic compounds.

Fatty acids, serving as precursors for diverse lipids and for the assembly of biological membranes, are nevertheless toxic to a wide variety of bacteria. Unlike most other bacteria, and for reasons that are not well understood, Actinomycetes commonly synthesize and accumulate extremely large amounts (20-80% of their biomass) of triacyl glycerols (40). This includes the primary precursor of complex lipids, palmitic acid, along with several unsaturated fatty acid derivatives, oleic, linoleic (unpublished data) and arachidonic (unpublished data) acids. All induced *whiB7*, with palmitic acid being the most active (Fig. 3e). The fact that the *whiB7* regulon, including antibiotic resistance genes, can be activated by palmitic acid has important implications for mycobacterial chemotherapy. Palmitic acid has been found in mycobacterial cytosol, and is considered to be a major source of carbon used by *M. tuberculosis* in the mammalian macrophage (41). It is also the principle fatty acid found in animal tissues

and serum. Therefore, the *whiB7* regulon may be induced when *M. tuberculosis* enters macrophages or other lipid rich cells, organs, or tissues, thereby allowing mycobacteria to more effectively resist some chemotherapeutic strategies, sheltered in specific areas of the body.

The intrinsic resistance of *M. tuberculosis* to antibiotics during *in vivo* growth and persistence underlies the need for protracted therapy for tuberculosis (2). Knowledge of such inducible intrinsic mycobacterial systems could generate derivatives of antibiotics that might circumvent detection by *whiB7* regulators or perhaps *WhiB7* inhibitors that augment conventional therapies by inactivating groups of genes that confer intrinsic resistance. Such developments could not only open up a powerful repertoire of currently redundant clinical antibiotics in the treatment of tuberculosis but also reduce the problematic duration of chemotherapy.

► Acknowledgements

We thank Keith Chater and Berthold Gust (John Innes Centre, Norwich, U.K.) for providing us with a collection of *S. coelicolor* mutants having mutations in genes belonging to the *whiB* family. This work was supported by the Swiss National Science Foundation Grants 4049-69384 (to C.J.T. and J.P.), a World Health Organization grant (to J.P.), and National Institutes of Health Grant AI44826 (to G.S.).

► Footnotes

Author contributions: R.P.M., L.N., G.S., and C.J.T. designed research; R.P.M., L.N., J.G., K.N., D.S., S.E., and Y.L. performed research; R.P.M., L.N., J.G., K.V., L.H., J.P., G.S., and C.J.T. contributed new reagents/analytic tools; R.P.M., L.N., J.G., K.V., J.P., G.S., and C.J.T. analyzed data; and R.P.M., L.N., G.S., and C.J.T. wrote the paper.

Freely available online through the PNAS open access option.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. AF205848 [[GenBank](#)]).

§ Present address: Cubist Pharmaceuticals, 65 Hayden Avenue, Lexington, MA 02421. ▀

** To whom correspondence should be addressed. E-mail: charles.thompson@ubc.ca .

© 2005 by [The National Academy of Sciences of the USA](#)

► References

1. Dye, C., Scheele, S., Dolin, P., Pathania, V. & Raviglione, M. C. (1999) *J. Am. Med. Assoc.* **282**, 677-686.[[Abstract/Free Full Text](#)]
2. McKinney, J. D. (2000) *Nat. Med.* **6**, 1330-1333.[[CrossRef](#)][[ISI](#)][[Medline](#)]
3. Parrish, N. M., Dick, J. D. & Bishai, W. R. (1998) *Trends Microbiol.* **6**, 107-112.[[CrossRef](#)][[ISI](#)][[Medline](#)]
4. Jarlier, V. & Nikaido, H. (1994) *FEMS Microbiol. Lett.* **123**, 11-18.[[CrossRef](#)][[ISI](#)][[Medline](#)]
5. Nguyen, L., Chinnapapagari, S. & Thompson, C. J. (2005) *J. Bacteriol.*, in press.
6. Ainsa, J. A., Blokpoel, M. C., Otal, I., Young, D. B., De Smet, K. A. & Martin, C. (1998) *J. Bacteriol.* **180**, 5836-5843.[[Abstract/Free Full Text](#)]
7. Doucet-Populaire, F., Buriankova, K., Weiser, J. & Pernodet, J. L. (2002) *Curr. Drug Targets Infect Disord.* **2**, 355-370.[[CrossRef](#)][[Medline](#)]
8. Rawat, M., Newton, G. L., Ko, M., Martinez, G. J., Fahey, R. C. & Av-Gay, Y. (2002) *Antimicrob. Agents Chemother.* **46**, 3348-3355.[[Abstract/Free Full Text](#)]
9. Cosma, C. L., Sherman, D. R. & Ramakrishnan, L. (2003) *Annu. Rev. Microbiol.* **57**, 641-676.[[CrossRef](#)][[ISI](#)][[Medline](#)]
10. Davies, J. (1994) *Science* **264**, 375-382.[[ISI](#)][[Medline](#)]
11. Folcher, M., Morris, R. P., Dale, G., Salah-Bey-Hocini, K., Viollier, P. H. & Thompson, C. J. (2001) *J. Biol. Chem.* **276**, 1479-1485.[[Abstract/Free Full Text](#)]
12. Goh, E. B., Yim, G., Tsui, W., McClure, J., Surette, M. G. & Davies, J. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 17025-17030.[[Abstract/Free Full Text](#)]

▲ Top
▲ Abstract
▲ Methods
▲ Results
▲ Discussion
• References

13. Novotna, J., Vohradsky, J., Berndt, P., Gramajo, H., Langen, H., Li, X. M., Minas, W., Orsaria, L., Roeder, D. & Thompson, C. J. (2003) *Mol. Microbiol.* **48**, 1289-1303.[CrossRef][ISI][Medline]
14. Murakami, T., Holt, T. G. & Thompson, C. J. (1989) *J. Bacteriol.* **171**, 1459-1466. [ISI][Medline]
15. Miller, C., Thomsen, L. E., Gaggero, C., Mosseri, R., Ingmer, H. & Cohen, S. N. (2004) *Science* **305**, 1629-1631.[Abstract/Free Full Text]
16. Wu, S., de Lencastre, H. & Tomasz, A. (1996) *J. Bacteriol.* **178**, 6036-6042. [Abstract/Free Full Text]
17. Barbosa, T. M. & Levy, S. B. (2000) *J. Bacteriol.* **182**, 3467-3474. [Abstract/Free Full Text]
18. Kieser, T., Bibb, M. B., Buttner, M. J., Chater, K. F. & Hopwood, D. A. (2000) *Practical Streptomyces Genetics* (The John Innes Foundation, Norwich, U.K.).
19. Braunstein, M., Bardarov, S. & Jacobs, W. J. (2002) *Methods Enzymol.* **358**, 67-99.[ISI][Medline]
20. Walburger, A., Koul, A., Ferrari, G., Nguyen, L., Prescianotto-Baschong, C., Huygen, K., Klebl, B., Thompson, C., Bacher, G. & Pieters, J. (2004) *Science* **304**, 1800-1804.[Abstract/Free Full Text]
21. Schnappinger, D., Ehrt, S., Voskuil, M. I., Liu, Y., Mangan, J. A., Monahan, I. M., Dolganov, G., Efron, B., Butcher, P. D., Nathan, C. & Schoolnik, G. K. (2003) *J. Exp. Med.* **198**, 693-704.[Abstract/Free Full Text]
22. Tusher, V. G., Tibshirani, R. & Chu, G. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 5116-5121.[Abstract/Free Full Text]
23. Bentley, S. D., Chater, K. F., Cerdeno-Tarraga, A. M., Challis, G. L., Thomson, N. R., James, K. D., Harris, D. E., Quail, M. A., Kieser, H., Harper, D., *et al.* (2002) *Nature* **417**, 141-147.[CrossRef][ISI][Medline]
24. Davis, N. K. & Chater, K. F. (1992) *Mol. Gen. Genet* **232**, 351-358.[ISI][Medline]
25. Steyn, A. J., Collins, D. M., Hondalus, M. K., Jacobs, W. R., Jr., Kawakami, R. P. & Bloom, B. R. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 3147-3152. [Abstract/Free Full Text]
26. Sun, J., Kelemen, G. H., Fernandez-Abalos, J. M. & Bibb, M. J. (1999) *Microbiology* **145**, 2221-2227.[ISI][Medline]
27. Soliveri, J. A., Gomez, J., Bishai, W. R. & Chater, K. F. (2000) *Microbiology* **146**, 333-343.[ISI][Medline]
28. Chater, K. F. (2001) *Curr. Opin. Microbiol.* **4**, 667-673.[CrossRef][ISI][Medline]
29. Jakimowicz, P., Cheesman, M. R., Bishai, W. R., Chater, K. F., Thomson, A. J. & Buttner, M. J. (2004) *J. Biol. Chem.* **280**, 8309-8315.[CrossRef][ISI][Medline]

30. Salah-Bey, K., Blanc, V. & Thompson, C. J. (1995) *Mol. Microbiol.* **17**, 1001-1012.
[CrossRef][ISI][Medline]
31. Saito, H., Tomioka, H. & Yoneyama, T. (1984) *Antimicrob. Agents Chemother.* **26**, 164-169.[ISI][Medline]
32. Kanetsuna, F. (1985) *Microbiol. Immunol.* **29**, 127-141.[ISI][Medline]
33. De Rossi, E., Arrigo, P., Bellinzoni, M., Silva, P. A., Martin, C., Ainsa, J. A., Gugliera, P. & Riccardi, G. (2002) *Mol. Med.* **8**, 714-724.[ISI][Medline]
34. Buriankova, K., Doucet-Populaire, F., Dorson, O., Gondran, A., Ghnassia, J. C., Weiser, J. & Pernodet, J. L. (2004) *Antimicrob. Agents Chemother.* **48**, 143-150.
[Abstract/Free Full Text]
35. Martinez, C., Nicolas, A., van Tilbeurgh, H., Egloff, M. P., Cudrey, C., Verger, R. & Cambillau, C. (1994) *Biochemistry* **33**, 83-89.[CrossRef][ISI][Medline]
36. Nathan, C. (2004) *Nature* **431**, 899-902.[CrossRef][ISI][Medline]
37. Gomez, J. E. & Bishai, W. R. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 8554-8559.
[Abstract/Free Full Text]
38. Hèachler, H., Cohen, S. P. & Levy, S. B. (1991) *J. Bacteriol.* **173**, 5532-5538.[ISI][Medline]
39. Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V., Eiglmeier, K., Gas, S., Barry, C. E., III, et al. (1998) *Nature* **393**, 537-544.
[CrossRef][ISI][Medline]
40. Alvarez, H. M. & Steinbuchel, A. (2002) *Appl. Microbiol. Biotechnol.* **60**, 367-376.
[CrossRef][ISI][Medline]
41. Wheeler, P. R. (1994) in *Tuberculosis: Pathogenesis, Protection and Control*, ed. Bloom, B. R. (Am. Soc. Microbiol. Press, Washington, DC), pp. 353-385.

**This article has been cited by other articles in
HighWire Press-hosted journals:** ([Search Google Scholar for Other
Citing Articles](#))



Journal of Bacteriology [HOME](#)

L. Nguyen, S. Chinnapapagari, and C. J. Thompson
**FbpA-Dependent Biosynthesis of Trehalose Dimycolate
Is Required for the Intrinsic Multidrug Resistance, Cell
Wall Structure, and Colonial Morphology of
*Mycobacterium smegmatis***

ATTACHMENT 5

CUSTOMER NO. 36257

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Virginia Goss TUSHER et al.
Title: Significance Analysis of Microarrays
Application No.: 09/811,762 Filing Date: March 19, 2001
Examiner: Lori A. CLOW Group Art Unit: 1631
Docket No.: STAN.058US1 Conf. No.: 8102

Mail Stop Petitions
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

**DECLARATION OF GILBERT CHU IN SUPPORT OF
AMENDMENT TO OVERCOME 35 U.S.C. § 101 REJECTION**

Sir/Madam:

I, Dr. Gilbert Chu, hereby declare as follows:

1. I am one of the inventors of the above-identified application,
2. SAM software that is referred to in the attached Bulletin "SAM: Significance Analysis Microarrays: Supervised learning software for genomic expression data mining" has been widely licensed to the public since 2001.
3. Two other co-authors and I published the article "*Significance analysis of microarrays applied to the ionizing radiation response*," Virginia Goss Tusher, et al., Proceedings of the National Academy of Sciences of the United States of America (PNAS), which was published April 24, 2001, Volume 98, No. 9, pages 5116-5121. This article is referred to herein as the "SAM article."

Attorney Docket No.: STAN.058US1

Application No.: 09/811,762

4. Attached is an article entitled "An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival," by Lance Miller et al., published in PNAS, September 20, 2005, Volume 102, No. 38, pages 13550-13555. As stated on page 6 of this article: "Univariate analysis by statistical analysis of microarrays (SAM) (22) identified 6,545 Affymetrix probe sets representing $\approx 5,290$ distinct genes whose expression patterns distinguished P53 mt and wt tumors with a false discovery rate (q value) $< 1\%$ and d score (modified t statistic) > 2.0 ..., further illuminating the extensive nature of the molecular variation underlying p53 status." Reference to "statistical analysis of microarrays (SAM) (22)" in the Miller article refers to the SAM article.

5. Attached is an article entitled "Ancestral antibiotic resistance in Mycobacterium tuberculosis," by Rowan P. Morris, et al., published in PNAS, August 23, 2005, Volume 102, No. 34, pages 12200-12205. As described on page 5 of this article by Morris et al., monocytes were infected by mycobacterium and activated. Labeling of RNA and hybridizations were performed. Then, "data from each experimental condition was analyzed separately by using significance analysis of microarrays (22) with a false discovery ratio $\leq 0.3\%$." The reference to "significance analysis of microarrays (22)" is to the SAM article.

6. All of the independent Claims of the above-identified application include the limitation that an expected value of the parameter is derived and compared to an observed or calculated value of the parameter, where the expected value is indicative of the extent of variations in the parameter introduced by the process by which the data (called associated values in the claims) are acquired. Claims 1, 44 and 58 also contain the limitation that the parameters of the plurality of genes be adjusted so that variables related to the parameters are substantially independent of variations of scatter values or average associated values of the genes over the sets, said scatter values defined by standard deviation of the associated values in the sets.

7. I believe that the analyses using SAM in the above quotes from the Miller and Morris articles employ the two claim limitations of paragraph 6 above through the use of SAM software.

8. I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that

these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Respectfully submitted,

Dated: 7/20/06



Gilbert Chu